

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 May 2003 (15.05.2003)

PCT

(10) International Publication Number
WO 03/039225 A2

- (51) International Patent Classification: Not classified
- (21) International Application Number: PCT/EP02/12449
- (22) International Filing Date:
7 November 2002 (07.11.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/331,045 7 November 2001 (07.11.2001) US
10/050,902 18 January 2002 (18.01.2002) US
PCT/IB02/00166 21 January 2002 (21.01.2002) IB
60/396,635 19 July 2002 (19.07.2002) US
- (71) Applicant (for all designated States except US): **CYTOS BIOTECHNOLOGY AG** [CH/CH]; Wagisstrasse 25, CH-8952 Zurich (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **BACHMANN, Martin** [CH/CH]; Goldackerweg 8, CH-8472 Seuzach (CH). **MAURER, Patrik** [DE/CH]; Richenbergstrasse 38, CH-8400 Winterthur (CH). **SPOHN, Gunther** [DE/CH]; Schulstrasse 36, CH-8952 Schlieren (CH).
- (74) Agent: **WICHMANN, Hendrik**; Isenbruck Bösl Hörschler Wichmann Huhn, Postfach 86 08 80, 81635 München (DE).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTIGEN ARRAYS FOR TREATMENT OF BONE DISEASE

(57) Abstract: The present invention is related to the fields of molecular biology, virology, immunology and medicine. The invention provides a composition comprising an ordered and repetitive antigen or antigenic determinant array, and in particular a RANKL protein, RANKL fragment or RANKL peptide-VLP-array. More specifically, the invention provides a composition comprising a virus-like particle and at least one RANKL protein, RANKL fragment or RANKL peptide bound thereto. The invention also provides a process for producing the conjugates and the ordered and repetitive arrays, respectively. The compositions of the invention are useful in the production of vaccines for the treatment of bone diseases and as a pharmaccine to prevent or cure bone diseases and to efficiently induce immune responses, in particular antibody responses. Furthermore, the compositions of the invention are particularly useful to efficiently induce self-specific immune responses within the indicated context.

BEST AVAILABLE COPY

ANTIGEN ARRAYS FOR TREATMENT OF BONE DISEASE

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] The present invention is related to the fields of molecular biology, virology, immunology and medicine. The invention provides a composition comprising an ordered and repetitive antigen or antigenic determinant array, and in particular a RANKL protein, RANKL fragment or RANKL peptide-array. More specifically, the invention provides a composition comprising a virus-like particle and at least one RANKL protein, RANKL fragment or RANKL peptide bound thereto. The invention also provides a process for producing the conjugates and the ordered and repetitive arrays, respectively. The compositions of the invention are useful in the production of vaccines for the treatment of bone diseases and as a pharmaccine to prevent or cure bone diseases and to efficiently induce immune responses, in particular antibody responses. Furthermore, the compositions of the invention are particularly useful to efficiently induce self-specific immune responses within the indicated context.

Related Art

[0002] Living bone is permanently turned over by balanced and coordinated remodeling processes. Primarily two cell types contribute to this remodeling: osteoblasts are essential for the formation of bone while osteoclasts promote dissolution of bone matrix and solubilization of the hydroxyapatite. In young individuals with growing bone, the rate of bone formation exceeds the rate of bone resorption, while in older individuals the rate of resorption can exceed formation and result in a net loss of bone mineral density and/or bone mass. In the latter case the bone strength is weakened and leads to an increased risk of fracture as well as slow or incomplete repair of broken bones. Multiple

- 2 -

conditions in humans are known to be associated with an imbalance in bone remodeling.

[0003] Recently three proteins have been described that are crucially involved in the formation of osteoclasts from hematopoietic precursor cells and regulation of bone remodeling. RANKL (Receptor activator of NFkB Ligand) which is also known as TNFSF11 (Tumor necrosis factor superfamily member 11), TRANCE (TNF-related activation induced cytokine), ODF (Osteoclast differentiation factor) or OPGL (Osteoprotegerin ligand) is a transmembrane protein of 245 amino acids that forms homotrimers. Part of the extracellular region of RANKL can be shed by a TACE-like protease. In addition, splice variants lacking the transmembrane region have been described. The shed part of RANKL contains the domain that is highly homologous to the TNF- α (Lum, L., *et al.*, *J. Biol. Chem.* 274: 13613-13618 (2000)).

[0004] Processes how to produce RANKL protein and RANKL fragments have been disclosed in WO 9846751, US 5,843,678, WO 98259958, US 6,242,586, WO 9828426, US 6,242,213, WO 9929865, JP 2000102390 and WO 0015807.

[0005] RANKL interacts with a transmembrane molecule on osteoclasts, termed RANK (Receptor activator of NFkB). This interaction leads to activation of the osteoclast precursor and ends in the formation of active, bone-resorbing osteoclasts. In vivo, a soluble decoy receptor termed osteoprotegerin, is involved in the regulation of osteoclastogenesis by its ability to bind to RANKL and inhibit the interaction of RANKL with its receptor RANK. This inhibition leads to a suppression of osteoclastogenesis and thus provides a means to stop excessive bone resorption. The interaction of RANKL with its receptor RANK can be suppressed by recombinant osteoprotegerin and by a soluble RANK-Fc fusion protein. In accordance with these findings, RANKL- and RANK-deficient mice develop osteopetrosis while RANKL-overexpressing transgenic mice as well as osteoprotegerin-deficient mice develop osteoporosis (Kong YY., *et al.*, *Nature* 397:315-322 (1999), Kim, N., *et al.*, *Proc. Natl. Acad. Sci USA* 97:10905-10910 (2000),

- 3 -

Dougall, B., et al., Proc. Natl. Acad. Sci USA 97:1566-1571 (1999), Bucay, N., et al., Genes Dev. 12: 1260-1268 (1998)).

[0006] The importance of the RANKL-RANK-osteoprotegerin system is further confirmed in an rodent model for osteoporosis induced by estrogen-deficiency. Recombinant osteoprotegerin completely abolished ovariectomy-induced bone loss (Simonet, W.S., et al. Cell 89:309-319 (1997).

[0007] In an adjuvant-induced arthritis model osteoprotegerin injection was able to prevent bone loss and cartilage destruction, but not inflammation (paw swelling). Beside its expression on stromal cells RANKL is also expressed on T cells, and RANK is found on antigen-presenting cells. It is assumed that during an arthritic reaction activated T cells with enhanced RANKL expression mediate an increase in osteoclastogenesis and subsequent bone loss. The interaction of RANKL with RANK also enhances the longevity and adjuvant properties of dendritic cells (Kong Y.Y., et al., Nature 402:304-309 (1999)).

[0008] Alveolar bone destruction and subsequent tooth loss is observed in periodontal infections. In vivo inhibition of RANKL function with osteoprotegerin diminished alveolar bone destruction and reduced the number of periodontal osteoclasts after microbial challenge (Teng, Y.T.A., et al., J. Clin. Invest. 106:R59-R67 (2000).

[0009] Bone tumors and certain tumor metastases are characterized by increased bone resorption due to an increased osteoclastogenesis (Hofbauer, L.C. and Heufelder A.E., J. Clin Endocrin. Met. 85:2355-2363 (2000). Osteoprotegerin was shown to inhibit prostate-cancer induced osteoclastogenesis and prevent prostate tumor growth in the bone of mice (Zhang Y., et al., J. Clin. Invest. 107:1219-1220 (2001). It also diminished advanced bone cancer pain in mice (Luger N.M., et al., Cancer Res. 61:4038-4047 (2001)). Multiple myeloma is a B cell malignancy characterized by the accumulation of plasma cells in the bone marrow and the development of osteolytic bone disease. In mouse models for multiple myeloma injection of osteoprotegerin or RANK-Fc fusion protein prevented the development of

- 4 -

lytic bone lesions and interfered with myeloma progression (Pearse RN., et al., Proc. Natl. Acad. Sci USA 98:11581-11586 (2001)).

[0010] Central to the etiology of aseptic loosening of prosthetic implants is periprosthetic osteolysis at the bone-implant interface, which is caused by wear-debris-induced inflammation. Fibroblast-like synoviocytes, transfected with osteoprotegerin, were able to prevent wear debris induced osteoclastogenesis in a mouse model (Gouter J.J., et al., J. Orthop. Res. 202:169-173 (2002)).

[0011] Vascular calcification is found with high clinical incidence in the osteoporotic patient population. An involvement of the RANKL-RANK-osteoprotegerin system is demonstrated by the finding that osteoprotegerin-deficient mice showed arterial calcification which could be reversed by recombinant osteoprotegerin (Min, H., et al., J. Exp. Med. 192:463-474 (2000)).

[0012] All these finding point to a crucial importance of the RANKL-RANK-osteoprotegerin system in regulation bone resorption in a variety of pathological conditions. So far, inhibition of bone loss has been mainly shown by injection of recombinant osteoprotegerin or a RANK-Fc fusion protein. Conceptually, immunization of an animal with RANKL should allow the production of RANKL-specific antibodies which, by binding to the RANK binding site or steric inhibition, should interfere with osteoclastogenesis.

[0013] However, so far no vaccination with a RANKL protein or peptide has been reported. Moreover, there has been no evidence that vaccines might be effective for protection against bone diseases, in particular, since it is usually difficult to induce antibody responses to self-molecules by conventional vaccination.

[0014] One way to improve the efficiency of vaccination is to increase the degree of repetitiveness of the antigen applied. Unlike isolated proteins, viruses induce prompt and efficient immune responses in the absence of any adjuvants both with and without T -cell help (Bachmann and Zinkernagel, *Ann. Rev. Immunol.* 15:235-270 (1991)). Although viruses often consist of

- 5 -

few proteins, they are able to trigger much stronger immune responses than their isolated components. For B-cell responses, it is known that one crucial factor for the immunogenicity of viruses is the repetitiveness and order of surface epitopes. Many viruses exhibit a quasi-crystalline surface that displays a regular array of epitopes which efficiently crosslinks epitope-specific immunoglobulins on B cells (Bachmann and Zinkernagel, *Immunol. Today* 17:553-558 (1996)). This crosslinking of surface immunoglobulins on B cells is a strong activation signal that directly induces cell-cycle progression and the production of IgM antibodies. Further, such triggered B cells are able to activate T helper cells, which in turn induce a switch from IgM to IgG antibody production in B cells and the generation of long-lived B cell memory - the goal of any vaccination (Bachmann and Zinkernagel, *Ann. Rev. Immunol.* 15:235-270 (1997)). Viral structure is even linked to the generation of anti-antibodies in autoimmune disease and as a part of the natural response to pathogens (see Fehr, T., *et al.*, *J Exp. Med.* 185:1785-1792 (1997)). Thus, antibodies presented by a highly organized viral surface are able to induce strong anti-antibody responses.

[0015] As indicated, however, the immune system usually fails to produce antibodies against self-derived structures. For soluble antigens present at low concentrations, this is due to tolerance at the Th cell level. Under these conditions, coupling the self-antigen to a carrier that can deliver T help may break tolerance. For soluble proteins present at high concentrations or membrane proteins at low concentration, B and Th cells may be tolerant. However, B cell tolerance may be reversible (anergy) and can be broken by administration of the antigen in a highly organized fashion coupled to a foreign carrier (Bachmann and Zinkernagel, *Ann. Rev. Immunol.* 15:235-270 (1997)).

- 6 -

BRIEF SUMMARY OF THE INVENTION

[0016] We have now found that RANKL proteins, RANKL fragments or RANKL peptides, which are bound to a core particle having a structure with an inherent repetitive organization, and hereby in particular to virus-like-particles (VLPs) and subunits of VLPs, respectively, leading to highly ordered and repetitive conjugates represent potent immunogens for the induction of antibodies specific for RANKL. The antibodies are able to block and neutralize, respectively, the interaction of RANKL with its receptor RANK. Therefore, the present invention provides a therapeutic mean for the treatment of bone diseases, which is based on an ordered and repetitive RANKL-core particle array, and in particular a VLP-RANKL-conjugate and -array, respectively. This therapeutic is able to induce high titers of anti-RANKL antibodies in a vaccinated animal.

[0017] The present invention, thus, provides for a composition comprising: (a) a core particle with at least one first attachment site; and (b) at least one antigen or antigenic determinant with at least one second attachment site, wherein said antigen or antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide, and wherein said second attachment site being selected from the group consisting of (i) an attachment site not naturally occurring with said antigen or antigenic determinant; and (ii) an attachment site naturally occurring with said antigen or antigenic determinant, wherein said second attachment site is capable of association to said first attachment site; and wherein said antigen or antigenic determinant and said core particle interact through said association to form an ordered and repetitive antigen array. Preferred embodiments of core particles suitable for use in the present invention are a virus, a virus-like particle, a bacteriophage, a bacterial pilus or flagella or any other core particle having an inherent repetitive structure capable of forming an ordered and repetitive antigen array in accordance with the present invention.

- 7 -

[0018] More specifically, the invention provides a composition comprising an ordered and repetitive antigen or antigenic determinant array, and hereby in particular RANKL protein-, RANKL fragment- or RANKL peptide-VLP conjugates. More specifically, the invention provides a composition comprising a virus-like particle and at least one RANKL protein, RANKL fragment or RANKL peptide bound thereto. The invention also provides a process for producing the conjugates and the ordered and repetitive arrays, respectively. The compositions of the invention are useful in the production of vaccines for the treatment of bone diseases and as a pharmaccine to prevent or cure bone diseases and to efficiently induce immune responses, in particular antibody responses. Furthermore, the compositions of the invention are particularly useful to efficiently induce self-specific immune responses within the indicated context.

[0019] In the present invention, a RANKL protein, RANKL fragment or RANKL peptide is bound to a core particle and VLP, respectively, typically in an oriented manner, yielding an ordered and repetitive RANKL protein, RANKL fragment or RANKL peptide antigen array. Furthermore, the highly repetitive and organized structure of the core particles and VLPs, respectively, mediates the display of the RANKL protein, RANKL fragment or RANKL peptide in a highly ordered and repetitive fashion leading to a highly organized and repetitive antigen array. Furthermore, binding of the RANKL protein, RANKL fragment or RANKL peptide to the core particle and VLP, respectively, provides T helper cell epitopes, since the core particle and VLP is foreign to the host immunized with the core particle-RANKL protein, -RANKL fragment or -RANKL peptide array and VLP-RANKL protein, -RANKL fragment or -RANKL peptide array, respectively. Those arrays differ from prior art conjugates in their highly organized structure, dimensions, and in the repetitiveness of the antigen on the surface of the array.

[0020] In one aspect of the invention, the RANKL protein, RANKL fragment or RANKL peptide is expressed in a suitable expression host compatible with proper folding of the RANKL protein or RANKL fragment, or synthesized,

- 8 -

while the core particle and the VLP, respectively, is expressed and purified from an expression host suitable for the folding and assembly of the core particle and the VLP, respectively. RANKL protein, RANKL fragment or RANKL peptide may also be chemically synthesized. The RANKL protein, RANKL fragment or RANKL peptide array is then assembled by binding the RANKL protein, RANKL fragment or RANKL peptide to the core particle and the VLP, respectively.

[0021] In another aspect, the present invention provides for a composition comprising (a) a virus-like particle, and (b) at least one antigen or antigenic determinant, wherein said antigen or said antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide, and wherein said at least one antigen or antigenic determinant is bound to the virus-like particle.

[0022] In a further aspect, the present invention provides for a pharmaceutical composition comprising (a) the composition of claim 1 or claim 22, and (b) an acceptable pharmaceutical carrier.

[0023] In still a further aspect, the present invention provides for a vaccine composition comprising a composition comprising: (a) a core particle with at least one first attachment site; and (b) at least one antigen or antigenic determinant with at least one second attachment site, wherein said antigen or antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide, and wherein said second attachment site being selected from the group consisting of (i) an attachment site not naturally occurring with said antigen or antigenic determinant; and (ii) an attachment site naturally occurring with said antigen or antigenic determinant, wherein said second attachment site is capable of association to said first attachment site; and wherein said antigen or antigenic determinant and said core particle interact through said association to form an ordered and repetitive antigen array.

[0024] In a further aspect, the present invention provides for a vaccine composition comprising a composition, wherein said composition comprising (a) a virus-like particle; and (b) at least one antigen or antigenic determinant, wherein said antigen or said antigenic determinant is a RANKL protein,

RANKL fragment or RANKL peptide; and wherein said at least one antigen or antigenic determinant is bound to said virus-like particle.

[0025] In still a further aspect, the present invention provides for a process for producing a composition of claim 1 comprising (a) providing a virus-like particle; and (b) providing at least one antigen or antigenic determinant, wherein said antigen or said antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide; (c) combining said virus-like particle and said at least one antigen or antigenic determinant so that said at least one antigen or antigenic determinant is bound to said virus-like particle.

[0026] In still a further aspect, the present invention provides a process for producing a composition of claim 22 comprising: (a) providing a core particle with at least one first attachment site; (b) providing at least one antigen or antigenic determinant with at least one second attachment site, wherein said antigen or antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide, and wherein said second attachment site being selected from the group consisting of (i) an attachment site not naturally occurring with said antigen or antigenic determinant; and (ii) an attachment site naturally occurring with said antigen or antigenic determinant; and wherein said second attachment site is capable of association to said first attachment site; and (c) combining said core particle and said at least one antigen or antigenic determinant, wherein said antigen or antigenic determinant and said core particle interact through said association to form an ordered and repetitive antigen array.

[0027] In another aspect, the present invention provides for a method of immunization comprising administering the composition of claim 1 or claim 22 to an animal or human.

[0028] In a further aspect, the present invention provides for a use of a composition of claim 1 or claim 22 for the manufacture of a medicament for treatment of bone diseases.

[0029] In a still further aspect, the present invention provides for a use of a composition of claim 1 or claim 22 for the preparation of a medicament for the

- 10 -

therapeutic or prophylactic treatment of bone diseases, preferably of mammalian encephalopathies. Furthermore, in a still further aspect, the present invention provides for a use of a composition of claim 1 or claim 22, either in isolation or in combination with other agents, for the manufacture of a composition, vaccine, drug or medicament for therapy or prophylaxis of bone diseases, in particular mammalian encephalopathies, and/or for stimulating the mammalian immune system.

[0030] Therefore, the invention provides, in particular, vaccine compositions which are suitable for preventing and/or attenuating bone diseases or conditions related thereto. The invention further provides and immunization and vaccination methods, respectively, for preventing and/or attenuating bone diseases or conditions related thereto, in animals, and in particular in cows, sheep and cattle as well as in humans. The inventive compositions may be used prophylactically or therapeutically.

[0031] In specific embodiments, the invention provides methods for preventing and/or attenuating bone diseases or conditions related thereto which are caused or exacerbated by "self" gene products, i.e. "self antigens" as used herein. In related embodiments, the invention provides methods for inducing immunological responses in animals and individuals, respectively, which lead to the production of antibodies that prevent and/or attenuate bone diseases or conditions related thereto, which are caused or exacerbated by "self" gene products.

[0032] As would be understood by one of ordinary skill in the art, when compositions of the invention are administered to an animal or a human, they may be in a composition which contains salts, buffers, adjuvants, or other substances which are desirable for improving the efficacy of the composition. Examples of materials suitable for use in preparing pharmaceutical compositions are provided in numerous sources including *Remington's Pharmaceutical Sciences* (Osol, A, ed., Mack Publishing Co. (1990)).

[0033] Compositions of the invention are said to be "pharmacologically acceptable" if their administration can be tolerated by a recipient individual.

- 11 -

Further, the compositions of the invention will be administered in a "therapeutically effective amount" (*i.e.*, an amount that produces a desired physiological effect).

[0034] The compositions of the present invention may be administered by various methods known in the art, but will normally be administered by injection, infusion, inhalation, oral administration, or other suitable physical methods. The compositions may alternatively be administered intramuscularly, intravenously, or subcutaneously. Components of compositions for administration include sterile aqueous (*e.g.*, physiological saline) or non-aqueous solutions and suspensions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Carriers or occlusive dressings can be used to increase skin permeability and enhance antigen absorption.

[0035] Other embodiments of the present invention will be apparent to one of ordinary skill in light of what is known in the art, the following drawings and description of the invention, and the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0036] FIG. 1 shows the expression and purification of C-RANKL.

Purification of C-RANKL was analysed on a SDS gel under reducing conditions. The gel was stained with Coomassie Brilliant Blue. Molecular weights of marker proteins are given on the left margin. Lane 1: low molecular weight marker. Lanes 2 and 3: the supernatant of the cell lysates of the BL21/DE3 cells transformed with the empty vector pGEX6p1 and pGEX-RANKL, respectively, after sixteen hours of induction with IPTG 0.4 mM. Lane 4: the purified GST-PS-C-RANKL protein after GST-Trap FF column. Lane 5: the GST-Trap FF column unbound fraction. Lane 6: the purified GST-PS-C-RANKL protein after the cleavage with the PreScission protease. Lane

- 12 -

7: the unbound fraction of the GST-Trap FF column loaded with the GST-RANKL digestion, which contains the purified C-RANKL. Lane 8: the bound fraction of the GST-Trap FF column loaded with the GST-PS-C-RANKL digestion and eluted with GSH.

[0037] FIG. 2 shows the expression and purification of RANKL-C.

FIG 2A shows purification of GST-EK-RANKL-C. Proteins samples were analyzed on a SDS-PAGE under reducing conditions. The gel was stained with Coomassie Brilliant Blue. Molecular weights of marker proteins are given on the left margin. Lane 1: Prestained protein marker, broad range (New England Biolabs). Lane 2: cleared cell lysate of BL21/DE3 cells transformed with the pMod-GST-EK-mRANKL-C1 plasmid after overnight induction with 0.1 mM IPTG. Lane 3: flow through of the GST-Trap FF column loaded with the cleared lysate of lane 2. Lane 4: first wash of GST-Trap FF column. Lane 5: second wash of GST-Trap FF column. Lane 6: third wash of GST-Trap FF column. Lanes 7-15: elution fractions 1 to 9 of the GST-Trap FF column containing the purified GST-EK-RANKL-C fusion protein and a minor amount of GST-EK protein.

[0038] FIG 2B shows cleavage of GST-EK-RANKL-C with EnterokinaseMaxTM.

Digestion of GST-EK-RANKL-C was analysed on SDS-PAGE under reducing conditions. The gel was stained with Coomassie Brilliant Blue. Molecular weights of marker proteins are given on the left margin. Lane 1: Prestained protein marker, broad range (New England Biolabs). Lane 2: Purified GST-EK-RANKL-C fusion protein. Lane 3: cleavage products after 16 h incubation at 4°C with EnterokinaseMaxTM.

[0039] FIG 2C shows purification of RANKL-C.

Purification of RANKL-C after removal of GST-EK by affinity chromatography on glutathione sepharose was analysed on SDS-PAGE under

- 13 -

reducing conditions. The gel was stained with Coomassie Brilliant Blue. Molecular weights of marker proteins are given on the left margin. Lane 1: Prestained protein marker, broad range (New England Biolabs). Lane 2 and 3: cleavage products GST-EK and RANKL-C after 16 h incubation of GST-EK-RANKL-C at 4°C with EnterokinaseMaxTM. Lane 4 and 5: different amounts of the unbound fraction of the GST-Trap FF column, which contains the RANKL-C protein in high purity.

[0040] FIG 3 shows coupling of C-RANKL to Q β capsid protein.

FIG 3A shows SDS-PAGE analysis of coupling products: Proteins were analysed on 16% SDS gels under reducing conditions. The gel was stained with Coomassie Brilliant Blue. Molecular weights of marker proteins are given on the left margin. Identity of protein bands is indicated on the right margin. Lane 1: Prestained protein marker, broad range (New England Biolabs). Lane 2: derivatized Q β capsid protein. Lane 3: purified C-RANKL protein. Lane 4: C-RANKL/Q β coupling reaction.

FIG 3B and FIG 3C: Western Blot analysis of coupling products.

Proteins were run on 16% SDS gels under reducing conditions, blotted to nitrocellulose membranes and detected with anti-Q β antiserum (FIG. 3B) or anti-RANKL antibody (FIG. 3C). Molecular weights of marker proteins are given on the left margin. Identity of protein bands is indicated on the right margin. Lane 1: Prestained protein marker, broad range (New England Biolabs). Lane 2: derivatized Q β capsid protein. Lane 3: purified C-RANKL protein. Lane 4: C-RANKL/Q β coupling reaction.

[0041] FIG. 4 shows ELISA for RANKL-specific IgG in mice immunized with C-RANKL coupled to Q β .

Female Balb/c mice were vaccinated subcutaneously with 25 μ g of C-RANKL coupled to Q β in PBS on day 0, day 16 and day 64 with or without the addition of alum. Sera from days 0, 16, 23, 64, and 78 were analysed for

- 14 -

antibodies specific for RANKL. ELISA titers are expressed as the average of those sera dilutions which lead to half maximal OD420 in the ELISA assay.

[0042] FIG. 5 shows neutralizing activity of antibodies induced in mice immunized with C-RANKL coupled to Q β .

FIG 5A shows binding assay of C-RANKL and its cognate ligand RANK. ELISA plates were coated with 10 μ g/ml C-RANKL and incubated with serial dilutions of RANK-Fc fusion protein or an unrelated Fc-fusion protein. Detection of bound RANK was performed with HRP-conjugated anti-Fc antibodies.

FIG. 5B shows inhibition of C-RANKL/RANK-Fc binding by serum antibodies of mice vaccinated with C-RANKL coupled to Q β Elisa plates were coated with 10 μ g/ml C-RANKL and co-incubated with serial dilutions of mouse sera from day 78 and 1 nM RANK-Fc fusion protein. Binding of fusion protein to C-RANKL was detected with horse raddish peroxidase conjugated anti-Fc antibody.

[0043] FIG. 6A-C depict the purification of AP205 proteins for use in VLPs, as analysed by SDS PAGE and Western-blotting.

[0044] FIG. 7A-C depict electron micrographs comparing AP205 phage particles to AP205 virus like particles spontaneously assembled from recombinant protein expressed in *E. coli* and purified.

DETAILED DESCRIPTION OF THE INVENTION

[0045] Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or

- 15 -

testing of the present invention, the preferred methods and materials are hereinafter described.

1. Definitions:

[0046] Amino acid linker: An "amino acid linker", or also just termed "linker" within this specification, as used herein, either associates the antigen or antigenic determinant with the second attachment site, or more preferably, already comprises or contains the second attachment site, typically - but not necessarily - as one amino acid residue, preferably as a cysteine residue. The term "amino acid linker" as used herein, however, does not intend to imply that such an amino acid linker consists exclusively of amino acid residues, even if an amino acid linker consisting of amino acid residues is a preferred embodiment of the present invention. The amino acid residues of the amino acid linker are, preferably, composed of naturally occurring amino acids or unnatural amino acids known in the art, all-L or all-D or mixtures thereof. However, an amino acid linker comprising a molecule with a sulfhydryl group or cysteine residue is also encompassed within the invention. Such a molecule comprise preferably a C1-C6 alkyl-, cycloalkyl (C5,C6), aryl or heteroaryl moiety. However, in addition to an amino acid linker, a linker comprising preferably a C1-C6 alkyl-, cycloalkyl- (C5,C6), aryl- or heteroaryl- moiety and devoid of any amino acid(s) shall also be encompassed within the scope of the invention. Association between the antigen or antigenic determinant or optionally the second attachment site and the amino acid linker is preferably by way of at least one covalent bond, more preferably by way of at least one peptide bond.

[0047] Animal: As used herein, the term "animal" is meant to include, for example, humans, sheep, elk, deer, mule deer, minks, mammals, monkeys, horses, cattle, pigs, goats, dogs, cats, rats, mice, birds, chicken, reptiles, fish, insects and arachnids.

- 16 -

[0048] Antibody: As used herein, the term "antibody" refers to molecules which are capable of binding an epitope or antigenic determinant. The term is meant to include whole antibodies and antigen-binding fragments thereof, including single-chain antibodies. Most preferably the antibodies are human antigen binding antibody fragments and include, but are not limited to, Fab, Fab' and F(ab')₂, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising either a V_L or V_H domain. The antibodies can be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine, rabbit, goat, guinea pig, camel, horse or chicken. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulins and that do not express endogenous immunoglobulins, as described, for example, in U.S. Patent No. 5,939,598 by Kucherlapati *et al.*

[0049] Antigen: As used herein, the term "antigen" refers to a molecule capable of being bound by an antibody or a T cell receptor (TCR) if presented by MHC molecules. The term "antigen", as used herein, also encompasses T-cell epitopes. An antigen is additionally capable of being recognized by the immune system and/or being capable of inducing a humoral immune response and/or cellular immune response leading to the activation of B- and/or T-lymphocytes. This may, however, require that, at least in certain cases, the antigen contains or is linked to a Th cell epitope and is given in adjuvant. An antigen can have one or more epitopes (B- and T- epitopes). The specific reaction referred to above is meant to indicate that the antigen will preferably react, typically in a highly selective manner, with its corresponding antibody or TCR and not with the multitude of other antibodies or TCRs which may be evoked by other antigens. Antigens as used herein may also be mixtures of several individual antigens.

[0050] Antigenic determinant: As used herein, the term "antigenic determinant" is meant to refer to that portion of an antigen that is specifically

- 17 -

recognized by either B- or T-lymphocytes. B-lymphocytes responding to antigenic determinants produce antibodies, whereas T-lymphocytes respond to antigenic determinants by proliferation and establishment of effector functions critical for the mediation of cellular and/or humoral immunity.

[0051] Association: As used herein, the term "association" as it applies to the first and second attachment sites, refers to the binding of the first and second attachment sites that is preferably by way of at least one non-peptide bond. The nature of the association may be covalent, ionic, hydrophobic, polar or any combination thereof, preferably the nature of the association is covalent.

[0052] Attachment Site, First: As used herein, the phrase "first attachment site" refers to an element of non-natural or natural origin, to which the second attachment site located on the antigen or antigenic determinant may associate. The first attachment site may be a protein, a polypeptide, an amino acid, a peptide, a sugar, a polynucleotide, a natural or synthetic polymer, a secondary metabolite or compound (biotin, fluorescein, retinol, digoxigenin, metal ions, phenylmethanesulfonylfluoride), or a combination thereof, or a chemically reactive group thereof. The first attachment site is located, typically and preferably on the surface, of the core particle such as, preferably the virus-like particle. Multiple first attachment sites are present on the surface of the core and virus-like particle, respectively, typically in a repetitive configuration.

[0053] Attachment Site, Second: As used herein, the phrase "second attachment site" refers to an element associated with the antigen or antigenic determinant to which the first attachment site located on the surface of the core particle and virus-like particle, respectively, may associate. The second attachment site of the antigen or antigenic determinant may be a protein, a polypeptide, a peptide, a sugar, a polynucleotide, a natural or synthetic polymer, a secondary metabolite or compound (biotin, fluorescein, retinol, digoxigenin, metal ions, phenylmethanesulfonylfluoride), or a combination thereof, or a chemically reactive group thereof. At least one second attachment site is present on the antigen or antigenic determinant. The term "antigen or antigenic determinant with at least one second attachment site"

- 18 -

refers, therefore, to an antigen or antigenic construct comprising at least the antigen or antigenic determinant and the second attachment site. However, in particular for a second attachment site, which is of non-natural origin, i.e. not naturally occurring within the antigen or antigenic determinant, these antigen or antigenic constructs comprise an "amino acid linker".

[0054] Bone diseases: The term "bone diseases", as used herein, encompasses, in particular, conditions characterized by increased bone resorption. The term "bone diseases" includes, but is not limited, to osteoporosis in its different forms such as primary osteoporosis (such as idiopathic, postmenopausal, involutional or senile osteoporosis) and secondary osteoporosis. The latter include osteoporosis caused by glucocorticoid excess, hyperparathyroidism, hyperthyroidism, hypergonadism, hyperprolactinemia, diabetes mellitus, drug-induced osteoporosis such as the one caused by glucocorticosteroids, ethanol, dilantin, tobacco, barbiturates or heparin), disuse-induced osteoporosis and miscellaneous conditions associated with increased bone resorption such as chronic renal failure, liver disease, malabsorption syndromes, chronic obstructive lung disease, sarcoidosis. Further conditions with increased bone resorption include Paget's disease, familial expansile osteolysis, spontaneous osteolysis, bone loss associated with rheumatoid arthritis, bone resorption and tooth loss during periodontal disease, osteomyelitis, hypercalcemia of malignancy, bone tumors, cancers associated by bone metastases and bone loss caused by weightlessness as found in space flight. Those skilled in the art can recognize further conditions characterized by increased bone resorption.

[0055] Bound: As used herein, the term "bound" refers to binding or attachment that may be covalent, *e.g.*, by chemically coupling, or non-covalent, *e.g.*, ionic interactions, hydrophobic interactions, hydrogen bonds, etc. Covalent bonds can be, for example, ester, ether, phosphoester, amide, peptide, imide, carbon-sulfur bonds, carbon-phosphorus bonds, and the like. The term "bound" is broader than and includes terms such as "coupled," "fused" and "attached".

[0056] Coat protein(s): As used herein, the term "coat protein(s)" refers to the protein(s) of a bacteriophage or a RNA-phage capable of being incorporated within the capsid assembly of the bacteriophage or the RNA-phage. However, when referring to the specific gene product of the coat protein gene of RNA-phages the term "CP" is used. For example, the specific gene product of the coat protein gene of RNA-phage Q β is referred to as "Q β CP", whereas the "coat proteins" of bacteriophage Q β comprise the "Q β CP" as well as the A1 protein. The capsid of Bacteriophage Q β is composed mainly of the Q β CP, with a minor content of the A1 protein. Likewise, the VLP Q β coat protein contains mainly Q β CP, with a minor content of A1 protein.

[0057] Core particle: As used herein, the term "core particle" refers to a rigid structure with an inherent repetitive organization. A core particle as used herein may be the product of a synthetic process or the product of a biological process.

[0058] Coupled: The term "coupled", as used herein, refers to attachment by covalent bonds or by strong non-covalent interactions, typically and preferably to attachment by covalent bonds. Any method normally used by those skilled in the art for the coupling of biologically active materials can be used in the present invention.

[0059] Effective Amount: As used herein, the term "effective amount" refers to an amount necessary or sufficient to realize a desired biologic effect. An effective amount of the composition would be the amount that achieves this selected result, and such an amount could be determined as a matter of routine by a person skilled in the art. For example, an effective amount for treating an immune system deficiency could be that amount necessary to cause activation of the immune system, resulting in the development of an antigen specific immune response upon exposure to antigen. The term is also synonymous with "sufficient amount."

[0060] The effective amount for any particular application can vary depending on such factors as the disease or condition being treated, the particular composition being administered, the size of the subject, and/or the severity of

- 20 -

the disease or condition. One of ordinary skill in the art can empirically determine the effective amount of a particular composition of the present invention without necessitating undue experimentation.

[0061] Epitope: As used herein, the term "epitope" refers to continuous or discontinuous portions of a polypeptide having antigenic or immunogenic activity in an animal, preferably a mammal, and most preferably in a human. An epitope is recognized by an antibody or a T cell through its T cell receptor in the context of an MHC molecule. An "immunogenic epitope," as used herein, is defined as a portion of a polypeptide that elicits an antibody response or induces a T-cell response in an animal, as determined by any method known in the art. (See, for example, Geysen *et al.*, *Proc. Natl. Acad. Sci. USA* 81:3998-4002 (1983)). The term "antigenic epitope," as used herein, is defined as a portion of a protein to which an antibody can immunospecifically bind its antigen as determined by any method well known in the art. Immunospecific binding excludes non-specific binding but does not necessarily exclude cross-reactivity with other antigens. Antigenic epitopes need not necessarily be immunogenic. Antigenic epitopes can also be T-cell epitopes, in which case they can be bound immunospecifically by a T-cell receptor within the context of an MHC molecule.

[0062] An epitope can comprise 3 amino acids in a spatial conformation which is unique to the epitope. Generally, an epitope consists of at least about 5 such amino acids, and more usually, consists of at least about 8-10 such amino acids. If the epitope is an organic molecule, it may be as small as Nitrophenyl.

[0063] Fusion: As used herein, the term "fusion" refers to the combination of amino acid sequences of different origin in one polypeptide chain by in-frame combination of their coding nucleotide sequences. The term "fusion" explicitly encompasses internal fusions, *i.e.*, insertion of sequences of different origin within a polypeptide chain, in addition to fusion to one of its termini.

[0064] Immune response: As used herein, the term "immune response" refers to a humoral immune response and/or cellular immune response leading to the

- 21 -

activation or proliferation of B- and/or T-lymphocytes and/or antigen presenting cells. In some instances, however, the immune responses may be of low intensity and become detectable only when using at least one substance in accordance with the invention. "Immunogenic" refers to an agent used to stimulate the immune system of a living organism, so that one or more functions of the immune system are increased and directed towards the immunogenic agent. An "immunogenic polypeptide" is a polypeptide that elicits a cellular and/or humoral immune response, whether alone or linked to a carrier in the presence or absence of an adjuvant. Preferably, antigen presenting cell may be activated.

[0065] A substance which "enhances" an immune response refers to a substance in which an immune response is observed that is greater or intensified or deviated in any way with the addition of the substance when compared to the same immune response measured without the addition of the substance. For example, the lytic activity of cytotoxic T cells can be measured, *e.g.* using a ^{51}Cr release assay, in samples obtained with and without the use of the substance during immunization. The amount of the substance at which the CTL lytic activity is enhanced as compared to the CTL lytic activity without the substance is said to be an amount sufficient to enhance the immune response of the animal to the antigen. In a preferred embodiment, the immune response is enhanced by a factor of at least about 2, more preferably by a factor of about 3 or more. The amount or type of cytokines secreted may also be altered. Alternatively, the amount of antibodies induced or their subclasses may be altered.

[0066] Immunization: As used herein, the terms "immunize" or "immunization" or related terms refer to conferring the ability to mount a substantial immune response (comprising antibodies and/or cellular immunity such as effector CTL) against a target antigen or epitope. These terms do not require that complete immunity be created, but rather that an immune response be produced which is substantially greater than baseline. For example, a mammal may be considered to be immunized against a target antigen if the

- 22 -

cellular and/or humoral immune response to the target antigen occurs following the application of methods of the invention.

[0067] Natural origin: As used herein, the term "natural origin" means that the whole or parts thereof are not synthetic and exist or are produced in nature.

[0068] Non-natural: As used herein, the term generally means not from nature, more specifically, the term means from the hand of man.

[0069] Non-natural origin: As used herein, the term "non-natural origin" generally means synthetic or not from nature; more specifically, the term means from the hand of man.

[0070] Ordered and repetitive antigen or antigenic determinant array: As used herein, the term "ordered and repetitive antigen or antigenic determinant array" generally refers to a repeating pattern of antigen or antigenic determinant, characterized by a typically and preferably uniform spatial arrangement of the antigens or antigenic determinants with respect to the core particle and virus-like particle, respectively. In one embodiment of the invention, the repeating pattern may be a geometric pattern. Typical and preferred examples of suitable ordered and repetitive antigen or antigenic determinant arrays are those which possess strictly repetitive paracrystalline orders of antigens or antigenic determinants, preferably with spacings of 1 to 30 nanometers, preferably 5 to 15 nanometers.

[0071] Pili: As used herein, the term "pili" (singular being "pilus") refers to extracellular structures of bacterial cells composed of protein monomers (*e.g.*, pilin monomers) which are organized into ordered and repetitive patterns. Further, pili are structures which are involved in processes such as the attachment of bacterial cells to host cell surface receptors, inter-cellular genetic exchanges, and cell-cell recognition. Examples of pili include Type-1 pili, P-pili, F1C pili, S-pili, and 987P-pili. Additional examples of pili are set out below.

[0072] Pilus-like structure: As used herein, the phrase "pilus-like structure" refers to structures having characteristics similar to that of pili and composed of protein monomers. One example of a "pilus-like structure" is a structure

- 23 -

formed by a bacterial cell which expresses modified pilin proteins that do not form ordered and repetitive arrays that are identical to those of natural pili.

[0073] Polypeptide: As used herein, the term "polypeptide" refers to a molecule composed of monomers (amino acids) linearly linked by amide bonds (also known as peptide bonds). It indicates a molecular chain of amino acids and does not refer to a specific length of the product. Thus, peptides, dipeptides, tripeptides, oligopeptides and proteins are included within the definition of polypeptide. This term is also intended to refer to post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations, and the like. A recombinant or derived polypeptide is not necessarily translated from a designated nucleic acid sequence. It may also be generated in any manner, including chemical synthesis.

[0074] RANKL protein: The term "RANKL protein" as used herein refers to a protein encoded by a RANKL gene. Different variants of the RANKL protein may be caused by nucleotide point mutations and polymorphisms, respectively, as well as insertions, deletions and/or substitutions of one or more nucleotides, and shall be explicitly encompassed within the scope of the present invention. Further variability can be caused by posttranslational modifications, such as differentially glycosylated forms of RANKL as well as proteolytically cleaved forms of RANKL (Lum, L., *et al.*, *J. Biol. Chem.* 274: 13613-13618 (2000)). There are a number of at present known splice variants of the human RANKL gene and the RANKL gene of other species which, with the possible variants mentioned above, are also within the scope of the invention. Therefore, the term "RANKL protein", as used herein, shall also encompass the RANKL protein variants, including but not limiting to the above indicated preferred examples.

[0075] As used herein, the term "RANKL fragment" is broadly defined as any polypeptide of at least 50 amino acids length which represents part of a RANKL protein, most preferably of a folded part of RANKL, and most preferably of the extracellular part of RANKL, even more preferably of the

- 24 -

region homologous to the TNF- α . The term RANKL fragment also encompasses recombinantly produced proteins and polypeptides, respectively, corresponding to splicing isoforms and proteolytically cleaved forms of RANKL, and all variants, as described above, thereof.

[0076] As used herein, the term "RANKL peptide" is broadly defined as any peptide which represents a fraction of a RANKL protein or a RANKL fragment and containing at least two, preferably at least three, more preferably at least four, more preferably at least five, more preferably at least six consecutive amino acids of the original RANKL protein or RANKL fragment, most preferably of the extracellular part of RANKL. Moreover, the term "RANKL peptide" shall preferably encompass any fraction of said RANKL peptide, wherein said fraction may be, preferably, derived by deletion of one or more amino acids at the N and/or C terminus. The RANKL peptide can be obtained by recombinant expression in eucaryotic or procaryotic expression systems as RANKL peptide alone or as a fusion with other amino acids or proteins, e.g. to facilitate folding, expression or solubility of the RANKL peptide or to facilitate purification of the RANKL peptide. To enable coupling of RANKL peptides and subunit proteins of VLPs or capsids, at least one second attachment site may be added to the RANKL peptide. Alternatively RANKL peptides may be synthesized using methods known to the art. The term RANKL peptide as used herein shall also preferably encompass a peptide which simulates the three dimensional surface structure of RANKL. Such RANKL peptide is not necessarily derived from a continuous amino acid sequence of RANKL, but may be formed by discontinuous amino acid residues from RANKL. Such peptides may even contain amino acids which are not present in the corresponding RANKL protein.

[0077] Residue: As used herein, the term "residue" is meant to mean a specific amino acid in a polypeptide backbone or side chain.

[0078] Self antigen: As used herein, the tem "self antigen" refers to proteins encoded by the host's DNA and products generated by proteins or RNA

- 25 -

encoded by the host's DNA are defined as self. In addition, proteins that result from a combination of two or several self-molecules or that represent a fraction of a self-molecule and proteins that have a high homology two self-molecules as defined above (>95%, preferably >97%, more preferably >99%) may also be considered self.

[0079] Treatment: As used herein, the terms "treatment", "treat", "treated" or "treating" refer to prophylaxis and/or therapy. When used with respect to an infectious disease, for example, the term refers to a prophylactic treatment which increases the resistance of a subject to infection with a pathogen or, in other words, decreases the likelihood that the subject will become infected with the pathogen or will show signs of illness attributable to the infection, as well as a treatment after the subject has become infected in order to fight the infection, *e.g.*, reduce or eliminate the infection or prevent it from becoming worse. When used with respect to bone disease, the term "treatment" refers to a prophylactic or therapeutic treatment which inhibits or reduces the increased bone resorption that is associated to bone diseases.

[0080] Vaccine: As used herein, the term "vaccine" refers to a formulation which contains the composition of the present invention and which is in a form that is capable of being administered to an animal. Typically, the vaccine comprises a conventional saline or buffered aqueous solution medium in which the composition of the present invention is suspended or dissolved. In this form, the composition of the present invention can be used conveniently to prevent, ameliorate, or otherwise treat a condition. Upon introduction into a host, the vaccine is able to provoke an immune response including, but not limited to, the production of antibodies and/or cytokines and/or the activation of cytotoxic T cells, antigen presenting cells, helper T cells, dendritic cells and/or other cellular responses.

[0081] Optionally, the vaccine of the present invention additionally includes an adjuvant which can be present in either a minor or major proportion relative to the compound of the present invention. The term "adjuvant" as used herein refers to non-specific stimulators of the immune response or substances that

- 26 -

allow generation of a depot in the host which when combined with the vaccine of the present invention provide for an even more enhanced immune response. A variety of adjuvants can be used. Examples include complete and incomplete Freund's adjuvant, aluminum hydroxide and modified muramyl dipeptide.

[0082] Virus-like particle (VLP): As used herein, the term "virus-like particle" refers to a structure resembling a virus particle. Moreover, a virus-like particle in accordance with the invention is non replicative and noninfectious since it lacks all or part of the viral genome, in particular the replicative and infectious components of the viral genome. A virus-like particle in accordance with the invention may contain nucleic acid distinct from their genome. A typical and preferred embodiment of a virus-like particle in accordance with the present invention is a viral capsid such as the viral capsid of the corresponding virus, bacteriophage, or RNA-phage. The terms "viral capsid" or "capsid", as interchangeably used herein, refer to a macromolecular assembly composed of viral protein subunits. Typically and preferably, the viral protein subunits assemble into a viral capsid and capsid, respectively, having a structure with an inherent repetitive organization, wherein said structure is, typically, spherical or tubular. For example, the capsids of RNA-phages or HBcAg's have a spherical form of icosahedral symmetry. The term "capsid-like structure" as used herein, refers to a macromolecular assembly composed of viral protein subunits resembling the capsid morphology in the above defined sense but deviating from the typical symmetrical assembly while maintaining a sufficient degree of order and repetitiveness.

[0083] Virus-like particle of a bacteriophage: As used herein, the term "virus-like particle of a bacteriophage" refers to a virus-like particle resembling the structure of a bacteriophage, being non replicative and noninfectious, and lacking at least the gene or genes encoding for the replication machinery of the bacteriophage, and typically also lacking the gene or genes encoding the protein or proteins responsible for viral attachment to or entry into the host. This definition should, however, also encompass virus-like particles of

- 27 -

bacteriophages, in which the aforementioned gene or genes are still present but inactive, and, therefore, also leading to non-replicative and noninfectious virus-like particles of a bacteriophage.

[0084] VLP of RNA phage coat protein: The capsid structure formed from the self-assembly of 180 subunits of RNA phage coat protein and optionally containing host RNA is referred to as a "VLP of RNA phage coat protein". A specific example is the VLP of Q β coat protein. In this particular case, the VLP of Q β coat protein may either be assembled exclusively from Q β CP subunits (generated by expression of a Q β CP gene containing, for example, a TAA stop codon precluding any expression of the longer A1 protein through suppression, see Kozlovskaya, T.M., *et al.*, *Intervirology* 39: 9-15 (1996)), or additionally contain A1 protein subunits in the capsid assembly.

[0085] Virus particle: The term "virus particle" as used herein refers to the morphological form of a virus. In some virus types it comprises a genome surrounded by a protein capsid; others have additional structures (*e.g.*, envelopes, tails, etc.).

[0086] One, a, or an: When the terms "one," "a," or "an" are used in this disclosure, they mean "at least one" or "one or more," unless otherwise indicated.

[0087] As will be clear to those skilled in the art, certain embodiments of the invention involve the use of recombinant nucleic acid technologies such as cloning, polymerase chain reaction, the purification of DNA and RNA, the expression of recombinant proteins in prokaryotic and eukaryotic cells, etc. Such methodologies are well known to those skilled in the art and can be conveniently found in published laboratory methods manuals (*e.g.*, Sambrook, J. *et al.*, eds., *Molecular Cloning, A Laboratory Manual*, 2nd. edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); Ausubel, F. *et al.*, eds., *Current Protocols in Molecular Biology*, John H. Wiley & Sons, Inc. (1997)). Fundamental laboratory techniques for working with tissue culture cell lines (Celis, J., ed., *Cell Biology*, Academic Press, 2nd edition, (1998)) and antibody-based technologies (Harlow, E. and Lane, D.,

- 28 -

Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1988); Deutscher, M.P., "Guide to Protein Purification," *Meth. Enzymol.* 128, Academic Press San Diego (1990); Scopes, R.K., *Protein Purification Principles and Practice*, 3rd ed., Springer-Verlag, New York (1994)) are also adequately described in the literature, all of which are incorporated herein by reference.

2. Compositions and Methods for Enhancing an Immune Response

[0088] The disclosed invention provides compositions and methods for enhancing an immune response against RANKL protein, RANKL fragment or RANKL peptide in an animal. Compositions of the invention comprise, or alternatively consist of (a) a core particle with at least one first attachment site; and (b) at least one antigen or antigenic determinant with at least one second attachment site, wherein said antigen or antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide, and wherein said second attachment site being selected from the group consisting of (i) an attachment site not naturally occurring with said antigen or antigenic determinant; and (ii) an attachment site naturally occurring with said antigen or antigenic determinant, wherein said second attachment site is capable of association to said first attachment site; and wherein said antigen or antigenic determinant and said core particle interact through said association to form an ordered and repetitive antigen array. More specifically, compositions of the invention comprise, or alternatively consist of, a virus-like particle and at least one antigen or antigenic determinant, wherein the antigen or antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide, and wherein the at least one antigen or antigenic determinant is bound to the virus-like particle so as to form an ordered and repetitive antigen-VLP-array. Furthermore, the invention conveniently enables the practitioner to construct such a composition, inter alia, for treatment and/or prophylactic prevention of bone diseases characterized by increased bone resorption.

- 29 -

[0089] In one embodiment, the core particle comprises a virus, a bacterial pilus, a structure formed from bacterial pilin, a bacteriophage, a virus-like particle, a viral capsid particle or a recombinant form thereof. Any virus known in the art having an ordered and repetitive coat and/or core protein structure may be selected as a core particle of the invention; examples of suitable viruses include sindbis and other alphaviruses, rhabdoviruses (e.g. vesicular stomatitis virus), picornaviruses (e.g., human rhino virus, Aichi virus), togaviruses (e.g., rubella virus), orthomyxoviruses (e.g., Thogoto virus, Batken virus, fowl plague virus), polyomaviruses (e.g., polyomavirus BK, polyomavirus JC, avian polyomavirus BFDV), parvoviruses, rotaviruses, Norwalk virus, foot and mouth disease virus, a retrovirus, Hepatitis B virus, Tobacco mosaic virus, Flock House Virus, and human Papillomavirus, and preferably a RNA phage, bacteriophage Q β , bacteriophage R17, bacteriophage M11, bacteriophage MX1, bacteriophage NL95, bacteriophage fr, bacteriophage GA, bacteriophage SP, bacteriophage MS2, bacteriophage f2, bacteriophage PP7 (for example, see Table 1 in Bachmann, M.F. and Zinkernagel, R.M., *Immunol. Today* 17:553-558 (1996)).

[0090] In a further embodiment, the invention utilizes genetic engineering of a virus to create a fusion between an ordered and repetitive viral envelope protein and a first attachment site comprising a heterologous protein, peptide, antigenic determinant or a reactive amino acid residue of choice. Other genetic manipulations known to those in the art may be included in the construction of the inventive compositions; for example, it may be desirable to restrict the replication ability of the recombinant virus through genetic mutation. Furthermore, the virus used for the present invention is replication incompetent due to chemical or physical inactivation or, as indicated, due to lack of a replication competent genome. The viral protein selected for fusion to the first attachment site should have an organized and repetitive structure. Such an organized and repetitive structure includes paracrystalline organizations with a spacing of 5-30 nm, preferably 5-15 nm, on the surface of the virus. The creation of this type of fusion protein will result in multiple,

- 30 -

ordered and repetitive first attachment sites on the surface of the virus and reflect the normal organization of the native viral protein. As will be understood by those in the art, the first attachment site may be or be a part of any suitable protein, polypeptide, sugar, polynucleotide, peptide (amino acid), natural or synthetic polymer, a secondary metabolite or combination thereof that may serve to specifically attach the antigen or antigenic determinant leading an ordered and repetitive antigen array.

[0091] In another embodiment of the invention, the core particle is a recombinant alphavirus, and more specifically, a recombinant Sindbis virus. Alphaviruses are positive stranded RNA viruses that replicate their genomic RNA entirely in the cytoplasm of the infected cell and without a DNA intermediate (Strauss, J. and Strauss, E., *Microbiol. Rev.* 58:491-562 (1994)). Several members of the alphavirus family, Sindbis (Xiong, C. *et al.*, *Science* 243:1188-1191 (1989); Schlesinger, S., *Trends Biotechnol.* 11:18-22 (1993)), Semliki Forest Virus (SFV) (Liljeström, P. & Garoff, H., *Bio/Technology* 9:1356-1361 (1991)) and others (Davis, N.L. *et al.*, *Virology* 171:189-204 (1989)), have received considerable attention for use as virus-based expression vectors for a variety of different proteins (Lundstrom, K., *Curr. Opin. Biotechnol.* 8:578-582 (1997); Liljeström, P., *Curr. Opin. Biotechnol.* 5:495-500 (1994)) and as candidates for vaccine development. Recently, a number of patents have issued directed to the use of alphaviruses for the expression of heterologous proteins and the development of vaccines (see U.S. Patent Nos. 5,766,602; 5,792,462; 5,739,026; 5,789,245 and 5,814,482). The construction of the alphaviral core particles of the invention may be done by means generally known in the art of recombinant DNA technology, as described by the aforementioned articles, which are incorporated herein by reference.

[0092] A variety of different recombinant host cells can be utilized to produce a viral-based core particle for antigen or antigenic determinant attachment. For example, alphaviruses are known to have a wide host range; Sindbis virus infects cultured mammalian, reptilian, and amphibian cells, as well as some insect cells (Clark, H., *J. Natl. Cancer Inst.* 51:645 (1973); Leake, C., *J. Gen.*

Virol. 35:335 (1977); Stollar, V. in *THE TOGAVIRUSES*, R.W. Schlesinger, Ed., Academic Press, (1980), pp.583-621). Thus, numerous recombinant host cells can be used in the practice of the invention. BHK, COS, Vero, HeLa and CHO cells are particularly suitable for the production of heterologous proteins because they have the potential to glycosylate heterologous proteins in a manner similar to human cells (Watson, E. *et al.*, *Glycobiology* 4:227, (1994)) and can be selected (Zang, M. *et al.*, *Bio/Technology* 13:389 (1995)) or genetically engineered (Renner W. *et al.*, *Biotech. Bioeng.* 4:476 (1995); Lee K. *et al.* *Biotech. Bioeng.* 50:336 (1996)) to grow in serum-free medium, as well as in suspension.

[0093] Introduction of the polynucleotide vectors into host cells can be effected by methods described in standard laboratory manuals (*see, e.g.*, Sambrook, J. *et al.*, eds., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd. edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), Chapter 9; Ausubel, F. *et al.*, eds., *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, John H. Wiley & Sons, Inc. (1997), Chapter 16), including methods such as electroporation, DEAE-dextran mediated transfection, transfection, microinjection, cationic lipid-mediated transfection, transduction, scrape loading, ballistic introduction, and infection. Methods for the introduction of exogenous DNA sequences into host cells are discussed in Felgner, P. *et al.*, U.S. Patent No. 5,580,859.

[0094] Packaged RNA sequences can also be used to infect host cells. These packaged RNA sequences can be introduced to host cells by adding them to the culture medium. For example, the preparation of non-infective alphaviral particles is described in a number of sources, including "Sindbis Expression System", Version C (*Invitrogen* Catalog No. K750-1).

[0095] When mammalian cells are used as recombinant host cells for the production of viral-based core particles, these cells will generally be grown in tissue culture. Methods for growing cells in culture are well known in the art (*see, e.g.*, Celis, J., ed., *CELL BIOLOGY*, Academic Press, 2nd edition, (1998); Sambrook, J. *et al.*, eds., *MOLECULAR CLONING, A LABORATORY MANUAL*,

2nd. edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); Ausubel, F. *et al.*, eds., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John H. Wiley & Sons, Inc. (1997); Freshney, R., CULTURE OF ANIMAL CELLS, Alan R. Liss, Inc. (1983)).

[0096] Further examples of RNA viruses suitable for use as core particles in the present invention include, but are not limited to, the following: members of the family Reoviridae, including the genus Orthoreovirus (multiple serotypes of both mammalian and avian retroviruses), the genus Orbivirus (Bluetongue virus, Eugenangee virus, Kemerovo virus, African horse sickness virus, and Colorado Tick Fever virus), the genus Rotavirus (human rotavirus, Nebraska calf diarrhea virus, murine rotavirus, simian rotavirus, bovine or ovine rotavirus, avian rotavirus); the family Picomaviridae, including the genus Enterovirus (poliovirus, Coxsackie virus A and B, enteric cytopathic human orphan (ECHO) viruses, hepatitis A, C, D, E and G viruses, Simian enteroviruses, Murine encephalomyelitis (ME) viruses, Poliovirus muris, Bovine enteroviruses, Porcine enteroviruses, the genus Cardiovirus (Encephalomyocarditis virus (EMC), Mengovirus), the genus Rhinovirus (Human rhinoviruses including at least 113 subtypes; other rhinoviruses), the genus Aphovirus (Foot and Mouth disease (FMDV); the family Calciviridae, including Vesicular exanthema of swine virus, San Miguel sea lion virus, Feline picornavirus and Norwalk virus; the family Togaviridae, including the genus Alphavirus (Eastern equine encephalitis virus, Semliki forest virus, Sindbis virus, Chikungunya virus, O'Nyong-Nyong virus, Ross river virus, Venezuelan equine encephalitis virus, Western equine encephalitis virus), the genus Flavivirus (Mosquito borne yellow fever virus, Dengue virus, Japanese encephalitis virus, St. Louis encephalitis virus, Murray Valley encephalitis virus, West Nile virus, Kunjin virus, Central European tick borne virus, Far Eastern tick borne virus, Kyasanur forest virus, Louping III virus, Powassan virus, Omsk hemorrhagic fever virus), the genus Rubivirus (Rubella virus), the genus Pestivirus (Mucosal disease virus, Hog cholera virus, Border disease virus); the family Bunyaviridae, including the genus Bunyavirus (Bunyamwera

and related viruses, California encephalitis group viruses), the genus Phlebovirus (Sandfly fever Sicilian virus, Rift Valley fever virus), the genus Nairovirus (Crimean-Congo hemorrhagic fever virus, Nairobi sheep disease virus), and the genus Uukuvirus (Uukuniemi and related viruses); the family Orthomyxoviridae, including the genus Influenza virus (Influenza virus type A, many human subtypes); Swine influenza virus, and Avian and Equine Influenza viruses; influenza type B (many human subtypes), and influenza type C (possible separate genus); the family paramyxoviridae, including the genus Paramyxovirus (Parainfluenza virus type 1, Sendai virus, Hemadsorption virus, Parainfluenza viruses types 2 to 5, Newcastle Disease Virus, Mumps virus), the genus Morbillivirus (Measles virus, subacute sclerosing panencephalitis virus, distemper virus, Rinderpest virus), the genus Pneumovirus (respiratory syncytial virus (RSV), Bovine respiratory syncytial virus and Pneumonia virus of mice); forest virus, Sindbis virus, Chikungunya virus, O'Nyong-Nyong virus, Ross river virus, Venezuelan equine encephalitis virus, Western equine encephalitis virus), the genus Flavivirus (Mosquito borne yellow fever virus, Dengue virus, Japanese encephalitis virus, St. Louis encephalitis virus, Murray Valley encephalitis virus, West Nile virus, Kunjin virus, Central European tick borne virus, Far Eastern tick borne virus, Kyasanur forest virus, Louping III virus, Powassan virus, Omsk hemorrhagic fever virus), the genus Rubivirus (Rubella virus), the genus Pestivirus (Mucosal disease virus, Hog cholera virus, Border disease virus); the family Bunyaviridae, including the genus Bunyavirus (Bunyamwera and related viruses, California encephalitis group viruses), the genus Phlebovirus (Sandfly fever Sicilian virus, Rift Valley fever virus), the genus Nairovirus (Crimean-Congo hemorrhagic fever virus, Nairobi sheep disease virus), and the genus Uukuvirus (Uukuniemi and related viruses); the family Orthomyxoviridae, including the genus Influenza virus (Influenza virus type A, many human subtypes); Swine influenza virus, and Avian and Equine Influenza viruses; influenza type B (many human subtypes), and influenza type C (possible separate genus); the family paramyxoviridae, including the genus

- 34 -

Paramyxovirus (Parainfluenza virus type 1, Sendai virus, Hemadsorption virus, Parainfluenza viruses types 2 to 5, Newcastle Disease Virus, Mumps virus), the genus Morbillivirus (Measles virus, subacute sclerosing panencephalitis virus, distemper virus, Rinderpest virus), the genus Pneumovirus (respiratory syncytial virus (RSV), Bovine respiratory syncytial virus and Pneumonia virus of mice); the family Rhabdoviridae, including the genus Vesiculovirus (VSV), Chandipura virus, Flanders-Hart Park virus), the genus Lyssavirus (Rabies virus), fish Rhabdoviruses and, filoviruses (Marburg virus and Ebola virus); the family Arenaviridae, including Lymphocytic choriomeningitis virus (LCM), Tacaribe virus complex, and Lassa virus; the family Coronaviridae, including Infectious Bronchitis Virus (IBV), Mouse Hepatitis virus, Human enteric corona virus, and Feline infectious peritonitis (Feline coronavirus).

[0097] Illustrative DNA viruses that may be used as core particles include, but are not limited to: the family Poxviridae, including the genus Orthopoxvirus (Variola major, Variola minor, Monkey pox Vaccinia, Cowpox, Buffalopox, Rabbitpox, Ectromelia), the genus Leporipoxvirus (Myxoma, Fibroma), the genus Avipoxvirus (Fowlpox, other avian poxvirus), the genus Capripoxvirus (sheeppox, goatpox), the genus Suipoxvirus (Swinepox), the genus Parapoxvirus (contagious postular dermatitis virus, pseudocowpox, bovine papular stomatitis virus); the family Iridoviridae (African swine fever virus, Frog viruses 2 and 3, Lymphocystis virus of fish); the family Herpesviridae, including the alpha-Herpesviruses (Herpes Simplex Types 1 and 2, Varicella-Zoster, Equine abortion virus, Equine herpes virus 2 and 3, pseudorabies virus, infectious bovine keratoconjunctivitis virus, infectious bovine rhinotracheitis virus, feline rhinotracheitis virus, infectious laryngotracheitis virus) the Beta-herpesviruses (Human cytomegalovirus and cytomegaloviruses of swine, monkeys and rodents); the gamma-herpesviruses (Epstein-Barr virus (EBV), Marek's disease virus, Herpes saimiri, Herpesvirus ateles, Herpesvirus sylvilagus, guinea pig herpes virus, Lucke tumor virus); the family Adenoviridae, including the genus Mastadenovirus (Human subgroups A, B,

- 35 -

C, D and E and ungrouped; simian adenoviruses (at least 23 serotypes), infectious canine hepatitis, and adenoviruses of cattle, pigs, sheep, frogs and many other species, the genus Aviadenovirus (Avian adenoviruses); and non-cultivable adenoviruses; the family Papoviridae, including the genus Papillomavirus (Human papilloma viruses, bovine papilloma viruses, Shope rabbit papilloma virus, and various pathogenic papilloma viruses of other species), the genus Polyomavirus (polyomavirus, Simian vacuolating agent (SV-40), Rabbit vacuolating agent (RKV), K virus, BK virus, JC virus, and other primate polyoma viruses such as Lymphotropic papilloma virus); the family Parvoviridae including the genus Adeno-associated viruses, the genus Parvovirus (Feline panleukopenia virus, bovine parvovirus, canine parvovirus, Aleutian mink disease virus, etc.). Finally, DNA viruses may include viruses such as chronic infectious neuropathic agents (CHINA virus).

[0098] In other embodiments, a bacterial pilin, a subportion of a bacterial pilin, or a fusion protein which contains either a bacterial pilin or subportion thereof is used to prepare compositions and vaccine compositions, respectively, of the invention. Examples of pilin proteins include pilins produced by *Escherichia coli*, *Haemophilus influenzae*, *Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Caulobacter crescentus*, *Pseudomonas stutzeri*, and *Pseudomonas aeruginosa*. The amino acid sequences of pilin proteins suitable for use with the present invention include those set out in GenBank reports AJ000636 (SEQ ID NO:1), AJ132364 (SEQ ID NO:2), AF229646 (SEQ ID NO: 3), AF051814 (SEQ ID NO:4), AF051815 (SEQ ID NO:5), and X00981 (SEQ ID NO:6), the entire disclosures of which are incorporated herein by reference.

[0099] Bacterial pilin proteins are generally processed to remove N-terminal leader sequences prior to export of the proteins into the bacterial periplasm. Further, as one skilled in the art would recognize, bacterial pilin proteins used to prepare compositions and vaccine compositions, respectively, of the invention will generally not have the naturally present leader sequence.

- 36 -

[00100] One specific example of a pilin protein suitable for use in the present invention is the P-pilin of *E. coli* (GenBank report AF237482 (SEQ ID NO:7)). An example of a Type-1 *E. coli* pilin suitable for use with the invention is a pilin having the amino acid sequence set out in GenBank report P04128 (SEQ ID NO:8), which is encoded by nucleic acid having the nucleotide sequence set out in GenBank report M27603 (SEQ ID NO:9). The entire disclosures of these GenBank reports are incorporated herein by reference. Again, the mature form of the above referenced protein would generally be used to prepare compositions and vaccine compositions, respectively, of the invention.

[00101] Bacterial pilins or pilin subportions suitable for use in the practice of the present invention will generally be able to associate to form ordered and repetitive antigen arrays.

[00102] Methods for preparing pili and pilus-like structures *in vitro* are known in the art. Bullitt *et al.*, *Proc. Natl. Acad. Sci. USA* 93:12890-12895 (1996), for example, describe the *in vitro* reconstitution of *E. coli* P-pili subunits. Furthermore, Eshdat *et al.*, *J. Bacteriol.* 148:308-314 (1981) describe methods suitable for dissociating Type-1 pili of *E. coli* and the reconstitution of pili. In brief, these methods are as follows: pili are dissociated by incubation at 37°C in saturated guanidine hydrochloride. Pilin proteins are then purified by chromatography, after which pilin dimers are formed by dialysis against 5 mM tris(hydroxymethyl) aminomethane hydrochloride (pH 8.0). Eshdat *et al.* also found that pilin dimers reassemble to form pili upon dialysis against the 5 mM tris(hydroxymethyl) aminomethane (pH 8.0) containing 5 mM MgCl₂.

[00103] Further, using, for example, conventional genetic engineering and protein modification methods, pilin proteins may be modified to contain a first attachment site to which an antigen or antigenic determinant is linked through a second attachment site. Alternatively, antigens or antigenic determinants can be directly linked through a second attachment site to amino acid residues which are naturally resident in these proteins. These modified pilin proteins may then be used in vaccine compositions of the invention.

- 37 -

[00104] Bacterial pilin proteins used to prepare compositions and vaccine compositions, respectively, of the invention may be modified in a manner similar to that described herein for HBcAg. For example, cysteine and lysine residues may be either deleted or substituted with other amino acid residues and first attachment sites may be added to these proteins. Further, pilin proteins may either be expressed in modified form or may be chemically modified after expression. Similarly, intact pili may be harvested from bacteria and then modified chemically.

[00105] In another embodiment, pili or pilus-like structures are harvested from bacteria (*e.g.*, *E. coli*) and used to form compositions and vaccine compositions of the invention. One example of pili suitable for preparing compositions and vaccine compositions is the Type-1 pilus of *E. coli*, which is formed from pilin monomers having the amino acid sequence set out in SEQ ID NO:8.

[00106] A number of methods for harvesting bacterial pili are known in the art. Bullitt and Makowski (*Biophys. J.* 74:623-632 (1998)), for example, describe a pilus purification method for harvesting P-pili from *E. coli*. According to this method, pili are sheared from hyperpiliated *E. coli* containing a P-pilus plasmid and purified by cycles of solubilization and $MgCl_2$ (1.0 M) precipitation.

[00107] Once harvested, pili or pilus-like structures may be modified in a variety of ways. For example, a first attachment site can be added to the pili to which antigens or antigen determinants may be attached through a second attachment site. In other words, bacterial pili or pilus-like structures can be harvested and modified to lead to ordered and repetitive antigen arrays.

[00108] Antigens or antigenic determinants could be linked to naturally occurring cysteine residues or lysine residues present in Pili or pilus-like structures. In such instances, the high order and repetitiveness of a naturally occurring amino acid residue would guide the coupling of the antigens or antigenic determinants to the pili or pilus-like structures. For example, the pili or pilus-like structures could be linked to the second attachment sites of the

- 38 -

antigens or antigenic determinants using a heterobifunctional cross-linking agent.

[00109] When structures which are naturally synthesized by organisms (*e.g.*, pili) are used to prepare compositions and vaccine compositions of the invention, it will often be advantageous to genetically engineer these organisms so that they produce structures having desirable characteristics. For example, when Type-1 pili of *E. coli* are used, the *E. coli* from which these pili are harvested may be modified so as to produce structures with specific characteristics. Examples of possible modifications of pilin proteins include the insertion of one or more lysine residues, the deletion or substitution of one or more of the naturally resident lysine residues, and the deletion or substitution of one or more naturally resident cysteine residues (*e.g.*, the cysteine residues at positions 44 and 84 in SEQ ID NO:8).

[00110] Further, additional modifications can be made to pilin genes which result in the expression products containing a first attachment site other than a lysine residue (*e.g.*, a *FOS* or *JUN* domain). Of course, suitable first attachment sites will generally be limited to those which do not prevent pilin proteins from forming pili or pilus-like structures suitable for use in vaccine compositions of the invention.

[00111] Pilin genes which naturally reside in bacterial cells can be modified *in vivo* (*e.g.*, by homologous recombination) or pilin genes with particular characteristics can be inserted into these cells. For examples, pilin genes could be introduced into bacterial cells as a component of either a replicable cloning vector or a vector which inserts into the bacterial chromosome. The inserted pilin genes may also be linked to expression regulatory control sequences (*e.g.*, a *lac* operator).

[00112] In most instances, the pili or pilus-like structures used in compositions and vaccine compositions, respectively, of the invention will be composed of single type of a pilin subunit. Pili or pilus-like structures composed of identical subunits will generally be used because they are expected to form structures which present highly ordered and repetitive antigen arrays.

- 39 -

[00113] However, the compositions of the invention also include compositions and vaccines comprising pili or pilus-like structures formed from heterogenous pilin subunits. The pilin subunits which form these pili or pilus-like structures can be expressed from genes naturally resident in the bacterial cell or may be introduced into the cells. When a naturally resident pilin gene and an introduced gene are both expressed in a cell which forms pili or pilus-like structures, the result will generally be structures formed from a mixture of these pilin proteins. Further, when two or more pilin genes are expressed in a bacterial cell, the relative expression of each pilin gene will typically be the factor which determines the ratio of the different pilin subunits in the pili or pilus-like structures.

[00114] When pili or pilus-like structures having a particular composition of mixed pilin subunits is desired, the expression of at least one of the pilin genes can be regulated by a heterologous, inducible promoter. Such promoters, as well as other genetic elements, can be used to regulate the relative amounts of different pilin subunits produced in the bacterial cell and, hence, the composition of the pili or pilus-like structures.

[00115] In additional, the antigen or antigenic determinant can be linked to bacterial pili or pilus-like structures by a bond which is not a peptide bond, bacterial cells which produce pili or pilus-like structures used in the compositions of the invention can be genetically engineered to generate pilin proteins which are fused to an antigen or antigenic determinant. Such fusion proteins which form pili or pilus-like structures are suitable for use in vaccine compositions of the invention.

[00116] Virus-like particles in the context of the present application refer to structures resembling a virus particle but which are not pathogenic. In general, virus-like particles lack the viral genome and, therefore, are noninfectious. Also, virus-like particles can be produced in large quantities by heterologous expression and can be easily purified.

[00117] In a preferred embodiment, the virus-like particle is a recombinant virus-like particle. The skilled artisan can produce VLPs using recombinant

- 40 -

DNA technology and virus coding sequences which are readily available to the public. For example, the coding sequence of a virus envelope or core protein can be engineered for expression in a baculovirus expression vector using a commercially available baculovirus vector, under the regulatory control of a virus promoter, with appropriate modifications of the sequence to allow functional linkage of the coding sequence to the regulatory sequence. The coding sequence of a virus envelope or core protein can also be engineered for expression in a bacterial expression vector, for example.

[00118] Examples of VLPs include, but are not limited to, the capsid proteins of Hepatitis B virus (Ulrich, *et al.*, *Virus Res.* 50:141-182 (1998)), measles virus (Warnes, *et al.*, *Gene* 160:173-178 (1995)), Sindbis virus, rotavirus (US 5,071,651 and US 5,374,426), foot-and-mouth-disease virus (Twomey, *et al.*, *Vaccine* 13:1603-1610, (1995)), Norwalk virus (Jiang, X., *et al.*, *Science* 250:1580-1583 (1990); Matsui, S.M., *et al.*, *J. Clin. Invest.* 87:1456-1461 (1991)), the retroviral GAG protein (WO 96/30523), the retrotransposon Ty protein p1, the surface protein of Hepatitis B virus (WO 92/11291), human papilloma virus (WO 98/15631), RNA phages, Ty, fr-phage, GA-phage and Q β -phage.

[00119] As will be readily apparent to those skilled in the art, the VLP of the invention is not limited to any specific form. The particle can be synthesized chemically or through a biological process, which can be natural or non-natural. By way of example, this type of embodiment includes a virus-like particle or a recombinant form thereof.

[00120] In a more specific embodiment, the VLP can comprise, or alternatively essentially consist of, or alternatively consist of recombinant polypeptides, or fragments thereof, being selected from recombinant polypeptides of Rotavirus, recombinant polypeptides of Norwalk virus, recombinant polypeptides of Alphavirus, recombinant polypeptides of Foot and Mouth Disease virus, recombinant polypeptides of measles virus, recombinant polypeptides of Sindbis virus, recombinant polypeptides of Polyoma virus, recombinant polypeptides of Retrovirus, recombinant polypeptides of Hepatitis B virus

(e.g., a HBcAg), recombinant polypeptides of Tobacco mosaic virus, recombinant polypeptides of Flock House Virus, recombinant polypeptides of human Papillomavirus, recombinant polypeptides of bacteriophages, recombinant polypeptides of RNA phages, recombinant polypeptides of Ty, recombinant polypeptides of fr-phage, recombinant polypeptides of GA-phage and recombinant polypeptides of Q β -phage. The virus-like particle can further comprise, or alternatively essentially consist of, or alternatively consist of, one or more fragments of such polypeptides, as well as variants of such polypeptides. Variants of polypeptides can share, for example, at least 80%, 85%, 90%, 95%, 97%, or 99% identity at the amino acid level with their wild-type counterparts.

[00121] In a preferred embodiment, the virus-like particle comprises, consists essentially of, or alternatively consists of recombinant proteins, or fragments thereof, of a RNA-phage. Preferably, the RNA-phage is selected from the group consisting of a) bacteriophage Q β ; b) bacteriophage R17; c) bacteriophage fr; d) bacteriophage GA; e) bacteriophage SP; f) bacteriophage MS2; g) bacteriophage M11; h) bacteriophage MX1; i) bacteriophage NL95; k) bacteriophage f2; and l) bacteriophage PP7.

[00122] In another preferred embodiment of the present invention, the virus-like particle comprises, or alternatively consists essentially of, or alternatively consists of recombinant proteins, or fragments thereof, of the RNA-bacteriophage Q β or of the RNA-bacteriophage fr.

[00123] In a further preferred embodiment of the present invention, the recombinant proteins comprise, or alternatively consist essentially of, or alternatively consist of coat proteins of RNA phages.

[00124] RNA-phage coat proteins forming capsids or VLPs, or fragments of the bacteriophage coat proteins compatible with self-assembly into a capsid or a VLP, are, therefore, further preferred embodiments of the present invention. Bacteriophage Q β coat proteins, for example, can be expressed recombinantly in *E. coli*. Further, upon such expression these proteins spontaneously form

- 42 -

capsids. Additionally, these capsids form a structure with an inherent repetitive organization.

[00125] Specific preferred examples of bacteriophage coat proteins which can be used to prepare compositions of the invention include the coat proteins of RNA bacteriophages such as bacteriophage Q β (SEQ ID NO:10; PIR Database, Accession No. VCBPQ β referring to Q β CP and SEQ ID NO: 11; Accession No. AAA16663 referring to Q β A1 protein), bacteriophage R17 (SEQ ID NO:12; PIR Accession No. VCBPR7), bacteriophage fr (SEQ ID NO:13; PIR Accession No. VCBPFR), bacteriophage GA (SEQ ID NO:14; GenBank Accession No. NP-040754), bacteriophage SP (SEQ ID NO:15; GenBank Accession No. CAA30374 referring to SP CP and SEQ ID NO: 16; Accession No. referring to SP A1 protein), bacteriophage MS2 (SEQ ID NO:17; PIR Accession No. VCBPM2), bacteriophage M11 (SEQ ID NO:18; GenBank Accession No. AAC06250), bacteriophage MX1 (SEQ ID NO:19; GenBank Accession No. AAC14699), bacteriophage NL95 (SEQ ID NO:20; GenBank Accession No. AAC14704), bacteriophage f2 (SEQ ID NO: 21; GenBank Accession No. P03611), bacteriophage PP7 (SEQ ID NO: 22). Furthermore, the A1 protein of bacteriophage Q β or C-terminal truncated forms missing as much as 100, 150 or 180 amino acids from its C-terminus may be incorporated in a capsid assembly of Q β coat proteins. Generally, the percentage of Q β A1 protein relative to Q β CP in the capsid assembly will be limited, in order to ensure capsid formation.

[00126] Q β coat protein has also been found to self-assemble into capsids when expressed in *E. coli* (Kozlovskaya TM. *et al.*, *GENE* 137: 133-137 (1993)). The obtained capsids or virus-like particles showed an icosahedral phage-like capsid structure with a diameter of 25 nm and T=3 quasi symmetry. Further, the crystal structure of phage Q β has been solved. The capsid contains 180 copies of the coat protein, which are linked in covalent pentamers and hexamers by disulfide bridges (Golmohammadi, R. *et al.*, *Structure* 4: 543-5554 (1996)) leading to a remarkable stability of the capsid of Q β coat protein.

- 43 -

Capsids or VLPs made from recombinant Q β coat protein may contain, however, subunits not linked via disulfide links to other subunits within the capsid, or incompletely linked. Thus, upon loading recombinant Q β capsid on non-reducing SDS-PAGE, bands corresponding to monomeric Q β coat protein as well as bands corresponding to the hexamer or pentamer of Q β coat protein are visible. Incompletely disulfide-linked subunits could appear as dimer, trimer or even tetramer band in non-reducing SDS-PAGE. Q β capsid protein also shows unusual resistance to organic solvents and denaturing agents. Surprisingly, we have observed that DMSO and acetonitrile concentrations as high as 30%, and Guanidinium concentrations as high as 1 M do not affect the stability of the capsid. The high stability of the capsid of Q β coat protein is an advantageous feature, in particular, for its use in immunization and vaccination of mammals and humans in accordance of the present invention.

[00127] Upon expression in *E. coli*, the N-terminal methionine of Q β coat protein is usually removed, as we observed by N-terminal Edman sequencing as described in Stoll, E. et al. *J. Biol. Chem.* 252:990-993 (1977). VLP composed from Q β coat proteins where the N-terminal methionine has not been removed, or VLPs comprising a mixture of Q β coat proteins where the N-terminal methionine is either cleaved or present are also within the scope of the present invention.

[00128] Further RNA phage coat proteins have also been shown to self-assemble upon expression in a bacterial host (Kastelein, RA. *et al.*, *Gene* 23: 245-254 (1983), Kozlovskaya, TM. *et al.*, *Dokl. Akad. Nauk SSSR* 287: 452-455 (1986), Adhin, MR. *et al.*, *Virology* 170: 238-242 (1989), Ni, CZ., *et al.*, *Protein Sci.* 5: 2485-2493 (1996), Priano, C. *et al.*, *J. Mol. Biol.* 249: 283-297 (1995)). The Q β phage capsid contains, in addition to the coat protein, the so called read-through protein A1 and the maturation protein A2. A1 is generated by suppression at the UGA stop codon and has a length of 329 aa. The capsid of phage Q β recombinant coat protein used in the invention is devoid of the A2 lysis protein, and contains RNA from the host. The coat protein of RNA

- 44 -

phages is an RNA binding protein, and interacts with the stem loop of the ribosomal binding site of the replicase gene acting as a translational repressor during the life cycle of the virus. The sequence and structural elements of the interaction are known (Witherell, GW. & Uhlenbeck, OC. *Biochemistry* 28: 71-76 (1989); Lim F. et al., *J. Biol. Chem.* 271: 31839-31845 (1996)). The stem loop and RNA in general are known to be involved in the virus assembly (Golmohammadi, R. et al., *Structure* 4: 543-5554 (1996)).

[00129] In a further preferred embodiment of the present invention, the virus-like particle comprises, or alternatively consists essentially of, or alternatively consists of recombinant proteins, or fragments thereof, of a RNA-phage, wherein the recombinant proteins comprise, consist essentially of or alternatively consist of mutant coat proteins of a RNA phage, preferably of mutant coat proteins of the RNA phages mentioned above. In another preferred embodiment, the mutant coat proteins of the RNA phage have been modified by removal of at least one lysine residue by way of substitution, or by addition of at least one lysine residue by way of substitution; alternatively, the mutant coat proteins of the RNA phage have been modified by deletion of at least one lysine residue, or by addition of at least one lysine residue by way of insertion.

[00130] In another preferred embodiment, the virus-like particle comprises, or alternatively consists essentially of, or alternatively consists of recombinant proteins, or fragments thereof, of the RNA-bacteriophage Q β , wherein the recombinant proteins comprise, or alternatively consist essentially of, or alternatively consist of coat proteins having an amino acid sequence of SEQ ID NO:10, or a mixture of coat proteins having amino acid sequences of SEQ ID NO:10 and of SEQ ID NO: 11 or mutants of SEQ ID NO: 11 and wherein the N-terminal methionine is preferably cleaved.

[00131] In a further preferred embodiment of the present invention, the virus-like particle comprises, consists essentially of or alternatively consists of recombinant proteins of Q β , or fragments thereof, wherein the recombinant proteins comprise, or alternatively consist essentially of, or alternatively

- 45 -

consist of mutant Q β coat proteins. In another preferred embodiment, these mutant coat proteins have been modified by removal of at least one lysine residue by way of substitution, or by addition of at least one lysine residue by way of substitution. Alternatively, these mutant coat proteins have been modified by deletion of at least one lysine residue, or by addition of at least one lysine residue by way of insertion.

[00132] Four lysine residues are exposed on the surface of the capsid of Q β coat protein. Q β mutants, for which exposed lysine residues are replaced by arginines can also be used for the present invention. The following Q β coat protein mutants and mutant Q β VLPs can, thus, be used in the practice of the invention: "Q β -240" (Lys13-Arg; SEQ ID NO:23), "Q β -243" (Asn 10-Lys; SEQ ID NO:24), "Q β -250" (Lys 2-Arg, Lys13-Arg; SEQ ID NO:25), "Q β -251" (SEQ ID NO:26) and "Q β -259" (Lys 2-Arg, Lys16-Arg; SEQ ID NO:27). Thus, in further preferred embodiment of the present invention, the virus-like particle comprises, consists essentially of or alternatively consists of recombinant proteins of mutant Q β coat proteins, which comprise proteins having an amino acid sequence selected from the group of a) the amino acid sequence of SEQ ID NO:23; b) the amino acid sequence of SEQ ID NO:24; c) the amino acid sequence of SEQ ID NO:25; d) the amino acid sequence of SEQ ID NO:26; and e) the amino acid sequence of SEQ ID NO:27. The construction, expression and purification of the above indicated Q β coat proteins, mutant Q β coat protein VLPs and capsids, respectively, are disclosed in pending U.S. Application No. 10/050,902 filed by the present assignee on January 18, 2002. In particular is hereby referred to Example 18 of above mentioned application.

[00133] In a further preferred embodiment of the present invention, the virus-like particle comprises, or alternatively consists essentially of, or alternatively consists of recombinant proteins of Q β , or fragments thereof, wherein the recombinant proteins comprise, consist essentially of or alternatively consist

- 46 -

of a mixture of either one of the foregoing Q β mutants and the corresponding A1 protein.

[00134] In a further preferred embodiment of the present invention, the virus-like particle comprises, or alternatively essentially consists of, or alternatively consists of recombinant proteins, or fragments thereof, of RNA-phage AP205.

[00135] The AP205 genome consists of a maturation protein, a coat protein, a replicase and two open reading frames not present in related phages; a lysis gene and an open reading frame playing a role in the translation of the maturation gene (Klovins, J., *et al.*, *J. Gen. Virol.* 83: 1523-33 (2002)). AP205 coat protein can be expressed from plasmid pAP283-58 (SEQ ID NO: 111), which is a derivative of pQb10 (Kozlovskaya, T. M., *et al.*, *Gene* 137:133-37 (1993)), and which contains an AP205 ribosomal binding site. Alternatively, AP205 coat protein may be cloned into pQb185, downstream of the ribosomal binding site present in the vector. Both approaches lead to expression of the protein and formation of capsids as described in the co-pending US provisional patent application with the title "Molecular Antigen Arrays" and having filed by the present assignee on July 16, 2002, which is incorporated by reference in its entirety. Vectors pQb10 and pQb185 are vectors derived from pGEM vector, and expression of the cloned genes in these vectors is controlled by the *trp* promoter (Kozlovskaya, T. M., *et al.*, *Gene* 137:133-37 (1993)). Plasmid pAP283-58 (SEQ ID NO:111) comprises a putative AP205 ribosomal binding site in the following sequence, which is downstream of the XbaI site, and immediately upstream of the ATG start codon of the AP205 coat protein: *tctaga*ATTTTCTGCGCACCCATCCCGGGTGGCGC-CCAAAGTGAGGAAAATCACatg. The vector pQb185 comprises a Shine Delagarno sequence downstream from the XbaI site and upstream of the start codon (*tctaga*TTAACCCAACGCGTAGGAG TCAGGCCatg, Shine Delagarno sequence underlined).

[00136] In a further preferred embodiment of the present invention, the virus-like particle comprises, or alternatively essentially consists of, or alternatively

consists of recombinant coat proteins, or fragments thereof, of the RNA-phage AP205..

[00137] This preferred embodiment of the present invention, thus, comprises AP205 coat proteins that form capsids. Such proteins are recombinantly expressed, or prepared from natural sources. AP205 coat proteins produced in bacteria spontaneously form capsids, as evidenced by Electron Microscopy (EM) and immunodiffusion. The structural properties of the capsid formed by the AP205 coat protein (SEQ ID NO: 112) and those formed by the coat protein of the AP205 RNA phage are nearly indistinguishable when seen in EM. AP205 VLPs are highly immunogenic, and can be linked with antigens and/or antigenic determinants to generate vaccine constructs displaying the antigens and/or antigenic determinants oriented in a repetitive manner. High titers are elicited against the so displayed antigens showing that bound antigens and/or antigenic determinants are accessible for interacting with antibody molecules and are immunogenic.

[00138] In a further preferred embodiment of the present invention, the virus-like particle comprises, or alternatively essentially consists of, or alternatively consists of recombinant mutant coat proteins, or fragments thereof, of the RNA-phage AP205.

[00139] Assembly-competent mutant forms of AP205 VLPs, including AP205 coat protein with the substitution of proline at amino acid 5 to threonine (SEQ ID NO: 113), may also be used in the practice of the invention and leads to a further preferred embodiment of the invention. These VLPs, AP205 VLPs derived from natural sources, or AP205 viral particles, may be bound to antigens to produce ordered repetitive arrays of the antigens in accordance with the present invention.

[00140] AP205 P5-T mutant coat protein can be expressed from plasmid pAP281-32 (SEQ ID No. 114), which is derived directly from pQb185, and which contains the mutant AP205 coat protein gene instead of the Q β coat protein gene. Vectors for expression of the AP205 coat protein are transfected into *E. coli* for expression of the AP205 coat protein.

- 48 -

[00141] Methods for expression of the coat protein and the mutant coat protein, respectively, leading to self-assembly into VLPs are described in co-pending US provisional patent application with the title "Molecular Antigen Arrays" and having filed by the present assignee on July 17, 2002, which is incorporated by reference in its entirety. Suitable *E. coli* strains include, but are not limited to, *E. coli* K802, JM 109, RR1. Suitable vectors and strains and combinations thereof can be identified by testing expression of the coat protein and mutant coat protein, respectively, by SDS-PAGE and capsid formation and assembly by optionally first purifying the capsids by gel filtration and subsequently testing them in an immunodiffusion assay (Ouchterlony test) or Electron Microscopy (Kozlovska, T. M.. *et al.*, *Gene* 137:133-37 (1993)).

[00142] AP205 coat proteins expressed from the vectors pAP283-58 and pAP281-32 may be devoid of the initial Methionine amino-acid, due to processing in the cytoplasm of *E. coli*. Cleaved, uncleaved forms of AP205 VLP, or mixtures thereof are further preferred embodiments of the invention.

[00143] In a further preferred embodiment of the present invention, the virus-like particle comprises, or alternatively essentially consists of, or alternatively consists of a mixture of recombinant coat proteins, or fragments thereof, of the RNA-phage AP205 and of recombinant mutant coat proteins, or fragments thereof, of the RNA-phage AP205.

[00144] In a further preferred embodiment of the present invention, the virus-like particle comprises, or alternatively essentially consists of, or alternatively consists of fragments of recombinant coat proteins or recombinant mutant coat proteins of the RNA-phage AP205.

[00145] Recombinant AP205 coat protein fragments capable of assembling into a VLP and a capsid, respectively are also useful in the practice of the invention. These fragments may be generated by deletion, either internally or at the termini of the coat protein and mutant coat protein, respectively. Insertions in the coat protein and mutant coat protein sequence or fusions of antigen sequences to the coat protein and mutant coat protein sequence, and

- 49 -

compatible with assembly into a VLP, are further embodiments of the invention and lead to chimeric AP205 coat proteins, and particles, respectively. The outcome of insertions, deletions and fusions to the coat protein sequence and whether it is compatible with assembly into a VLP can be determined by electron microscopy.

[00146] The particles formed by the AP205 coat protein, coat protein fragments and chimeric coat proteins described above, can be isolated in pure form by a combination of fractionation steps by precipitation and of purification steps by gel filtration using *e.g.* Sepharose CL-4B, Sepharose CL-2B, Sepharose CL-6B columns and combinations thereof as described in the co-pending US provisional patent application with the title "Molecular Antigen Arrays" and having filed by the present assignee on July 17, 2002, which is incorporated by reference in its entirety. Other methods of isolating virus-like particles are known in the art, and may be used to isolate the virus-like particles (VLPs) of bacteriophage AP205. For example, the use of ultracentrifugation to isolate VLPs of the yeast retrotransposon Ty is described in U.S. Patent No. 4,918,166, which is incorporated by reference herein in its entirety.

[00147] The crystal structure of several RNA bacteriophages has been determined (Golmohammadi, R. *et al.*, *Structure* 4:543-554 (1996)). Using such information, surface exposed residues can be identified and, thus, RNA-phage coat proteins can be modified such that one or more reactive amino acid residues can be inserted by way of insertion or substitution. As a consequence, those modified forms of bacteriophage coat proteins can also be used for the present invention. Thus, variants of proteins which form capsids or capsid-like structures (*e.g.*, coat proteins of bacteriophage Q β , bacteriophage R17, bacteriophage fr, bacteriophage GA, bacteriophage SP, and bacteriophage MS2) can also be used to prepare compositions of the present invention.

[00148] Although the sequence of the variants proteins discussed above will differ from their wild-type counterparts, these variant proteins will generally retain the ability to form capsids or capsid-like structures. Thus, the invention

- 50 -

further includes compositions and vaccine compositions, respectively, which further includes variants of proteins which form capsids or capsid-like structures, as well as methods for preparing such compositions and vaccine compositions, respectively, individual protein subunits used to prepare such compositions, and nucleic acid molecules which encode these protein subunits. Thus, included within the scope of the invention are variant forms of wild-type proteins which form capsids or capsid-like structures and retain the ability to associate and form capsids or capsid-like structures.

[00149] As a result, the invention further includes compositions and vaccine compositions, respectively, comprising proteins, which comprise, or alternatively consist essentially of, or alternatively consist of amino acid sequences which are at least 80%, 85%, 90%, 95%, 97%, or 99% identical to wild-type proteins which form ordered arrays and having an inherent repetitive structure, respectively.

[00150] Further included within the scope of the invention are nucleic acid molecules which encode proteins used to prepare compositions of the present invention.

[00151] In other embodiments, the invention further includes compositions comprising proteins, which comprise, or alternatively consist essentially of, or alternatively consist of amino acid sequences which are at least 80%, 85%, 90%, 95%, 97%, or 99% identical to any of the amino acid sequences shown in SEQ ID NOs:10-27.

[00152] Proteins suitable for use in the present invention also include C-terminal truncation mutants of proteins which form capsids or capsid-like structures, or VLPs. Specific examples of such truncation mutants include proteins having an amino acid sequence shown in any of SEQ ID NOs:10-27 where 1, 2, 5, 7, 9, 10, 12, 14, 15, or 17 amino acids have been removed from the C-terminus. Typically, these C-terminal truncation mutants will retain the ability to form capsids or capsid-like structures.

[00153] Further proteins suitable for use in the present invention also include N-terminal truncation mutants of proteins which form capsids or capsid-like

- 51 -

structures. Specific examples of such truncation mutants include proteins having an amino acid sequence shown in any of SEQ ID NOs:10-27 where 1, 2, 5, 7, 9, 10, 12, 14, 15, or 17 amino acids have been removed from the N-terminus. Typically, these N-terminal truncation mutants will retain the ability to form capsids or capsid-like structures.

[00154] Additional proteins suitable for use in the present invention include N- and C-terminal truncation mutants which form capsids or capsid-like structures. Suitable truncation mutants include proteins having an amino acid sequence shown in any of SEQ ID NOs:10-27 where 1, 2, 5, 7, 9, 10, 12, 14, 15, or 17 amino acids have been removed from the N-terminus and 1, 2, 5, 7, 9, 10, 12, 14, 15, or 17 amino acids have been removed from the C-terminus. Typically, these N-terminal and C-terminal truncation mutants will retain the ability to form capsids or capsid-like structures.

[00155] The invention further includes compositions comprising proteins which comprise, or alternatively consist essentially of, or alternatively consist of, amino acid sequences which are at least 80%, 85%, 90%, 95%, 97%, or 99% identical to the above described truncation mutants.

[00156] The invention thus includes compositions and vaccine compositions prepared from proteins which form capsids or VLPs, methods for preparing these compositions from individual protein subunits and VLPs or capsids, methods for preparing these individual protein subunits, nucleic acid molecules which encode these subunits, and methods for vaccinating and/or eliciting immunological responses in individuals using these compositions of the present invention.

[00157] As previously stated, the invention includes virus-like particles or recombinant forms thereof. In one further preferred embodiment, the particles used in compositions of the invention are composed of a Hepatitis B core protein (HBcAg) or a fragment of a HBcAg. In a further embodiment, the particles used in compositions of the invention are composed of a Hepatitis B core protein (HBcAg) or a fragment of a HBcAg protein, which has been modified to either eliminate or reduce the number of free cysteine residues.

- 52 -

Zhou *et al.* (*J. Virol.* 66:5393-5398 (1992)) demonstrated that HBcAgs which have been modified to remove the naturally resident cysteine residues retain the ability to associate and form capsids. Thus, VLPs suitable for use in compositions of the invention include those comprising modified HBcAgs, or fragments thereof, in which one or more of the naturally resident cysteine residues have been either deleted or substituted with another amino acid residue (*e.g.*, a serine residue).

[00158] The HBcAg is a protein generated by the processing of a Hepatitis B core antigen precursor protein. A number of isotypes of the HBcAg have been identified and their amino acids sequences are readily available to those skilled in the art. In most instances, compositions and vaccine compositions, respectively, of the invention will be prepared using the processed form of a HBcAg (*i.e.*, a HBcAg from which the N-terminal leader sequence of the Hepatitis B core antigen precursor protein have been removed).

[00159] Further, when HBcAgs are produced under conditions where processing will not occur, the HBcAgs will generally be expressed in "processed" form. For example, when an *E. coli* expression system directing expression of the protein to the cytoplasm is used to produce HBcAgs of the invention, these proteins will generally be expressed such that the N-terminal leader sequence of the Hepatitis B core antigen precursor protein is not present.

[00160] The preparation of Hepatitis B virus-like particles, which can be used for the present invention, is disclosed, for example, in WO 00/32227, and hereby in particular in Examples 17 to 19 and 21 to 24, as well as in WO 01/85208, and hereby in particular in Examples 17 to 19, 21 to 24, 31 and 41, and in pending U.S. Application No. 10/050,902 filed by the present assignee on January 18, 2002. For the latter application, it is in particular referred to Example 23, 24, 31 and 51. All three documents are explicitly incorporated herein by reference.

[00161] The present invention also includes HBcAg variants which have been modified to delete or substitute one or more additional cysteine residues. It is

known in the art that free cysteine residues can be involved in a number of chemical side reactions. These side reactions include disulfide exchanges, reaction with chemical substances or metabolites that are, for example, injected or formed in a combination therapy with other substances, or direct oxidation and reaction with nucleotides upon exposure to UV light. Toxic adducts could thus be generated, especially considering the fact that HBcAgs have a strong tendency to bind nucleic acids. The toxic adducts would thus be distributed between a multiplicity of species, which individually may each be present at low concentration, but reach toxic levels when together.

[00162] In view of the above, one advantage to the use of HBcAgs in vaccine compositions which have been modified to remove naturally resident cysteine residues is that sites to which toxic species can bind when antigens or antigenic determinants are attached would be reduced in number or eliminated altogether.

[00163] A number of naturally occurring HBcAg variants suitable for use in the practice of the present invention have been identified. Yuan *et al.*, (*J. Virol.* 73:10122-10128 (1999)), for example, describe variants in which the isoleucine residue at position corresponding to position 97 in SEQ ID NO:28 is replaced with either a leucine residue or a phenylalanine residue. The amino acid sequences of a number of HBcAg variants, as well as several Hepatitis B core antigen precursor variants, are disclosed in GenBank reports AAF121240 (SEQ ID NO:29), AF121239 (SEQ ID NO:30), X85297 (SEQ ID NO:31), X02496 (SEQ ID NO:32), X85305 (SEQ ID NO:33), X85303 (SEQ ID NO:34), AF151735 (SEQ ID NO:35), X85259 (SEQ ID NO:36), X85286 (SEQ ID NO:37), X85260 (SEQ ID NO:38), X85317 (SEQ ID NO:39), X85298 (SEQ ID NO:40), AF043593 (SEQ ID NO:41), M20706 (SEQ ID NO:42), X85295 (SEQ ID NO:43), X80925 (SEQ ID NO:44), X85284 (SEQ ID NO:45), X85275 (SEQ ID NO:46), X72702 (SEQ ID NO:47), X85291 (SEQ ID NO:48), X65258 (SEQ ID NO:49), X85302 (SEQ ID NO:50), M32138 (SEQ ID NO:51), X85293 (SEQ ID NO:52), X85315 (SEQ ID NO:53), U95551 (SEQ ID NO:54), X85256 (SEQ ID NO:55), X85316 (SEQ

- 54 -

ID NO:56), X85296 (SEQ ID NO:57), AB033559 (SEQ ID NO:58), X59795 (SEQ ID NO:59), X85299 (SEQ ID NO:60), X85307 (SEQ ID NO:61), X65257 (SEQ ID NO:62), X85311 (SEQ ID NO:63), X85301 (SEQ ID NO:64), X85314 (SEQ ID NO:65), X85287 (SEQ ID NO:66), X85272 (SEQ ID NO:67), X85319 (SEQ ID NO:68), AB010289 (SEQ ID NO:69), X85285 (SEQ ID NO:70), AB010289 (SEQ ID NO:71), AF121242 (SEQ ID NO:72), M90520 (SEQ ID NO:73), P03153 (SEQ ID NO:74), AF110999 (SEQ ID NO:75), and M95589 (SEQ ID NO:76), the disclosures of each of which are incorporated herein by reference. These HBcAg variants differ in amino acid sequence at a number of positions, including amino acid residues which corresponds to the amino acid residues located at positions 12, 13, 21, 22, 24, 29, 32, 33, 35, 38, 40, 42, 44, 45, 49, 51, 57, 58, 59, 64, 66, 67, 69, 74, 77, 80, 81, 87, 92, 93, 97, 98, 100, 103, 105, 106, 109, 113, 116, 121, 126, 130, 133, 135, 141, 147, 149, 157, 176, 178, 182 and 183 in SEQ ID NO:77. Further HBcAg variants suitable for use in the compositions of the invention, and which may be further modified according to the disclosure of this specification are described in WO 00/198333, WO 00/177158 and WO 00/214478.

[00164] As noted above, generally processed HBcAgs (*i.e.*, those which lack leader sequences) will be used in the compositions and vaccine compositions, respectively, of the invention. The present invention includes vaccine compositions, as well as methods for using these compositions, which employ the above described variant HBcAgs.

[00165] Whether the amino acid sequence of a polypeptide has an amino acid sequence that is at least 80%, 85%, 90%, 95%, 97% or 99% identical to one of the above wild-type amino acid sequences, or a subportion thereof, can be determined conventionally using known computer programs such the Bestfit program. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference amino acid sequence, the parameters are set such that the percentage of identity is calculated over the full length of the reference amino acid

- 55 -

sequence and that gaps in homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

[00166] The HBcAg variants and precursors having the amino acid sequences set out in SEQ ID NOs: 29-72 and 73-76 are relatively similar to each other. Thus, reference to an amino acid residue of a HBcAg variant located at a position which corresponds to a particular position in SEQ ID NO:77, refers to the amino acid residue which is present at that position in the amino acid sequence shown in SEQ ID NO:77. The homology between these HBcAg variants is for the most part high enough among Hepatitis B viruses that infect mammals so that one skilled in the art would have little difficulty reviewing both the amino acid sequence shown in SEQ ID NO:77 and that of a particular HBcAg variant and identifying "corresponding" amino acid residues. Furthermore, the HBcAg amino acid sequence shown in SEQ ID NO:73, which shows the amino acid sequence of a HBcAg derived from a virus which infect woodchucks, has enough homology to the HBcAg having the amino acid sequence shown in SEQ ID NO:77 that it is readily apparent that a three amino acid residue insert is present in SEQ ID NO:64 between amino acid residues 155 and 156 of SEQ ID NO:77.

[00167] The invention also includes vaccine compositions which comprise HBcAg variants of Hepatitis B viruses which infect birds, as well as vaccine compositions which comprise fragments of these HBcAg variants. For these HBcAg variants one, two, three or more of the cysteine residues naturally present in these polypeptides could be either substituted with another amino acid residue or deleted prior to their inclusion in vaccine compositions of the invention.

[00168] As discussed above, the elimination of free cysteine residues reduces the number of sites where toxic components can bind to the HBcAg, and also eliminates sites where cross-linking of lysine and cysteine residues of the same or of neighboring HBcAg molecules can occur. Therefore, in another embodiment of the present invention, one or more cysteine residues of the

- 56 -

Hepatitis B virus capsid protein have been either deleted or substituted with another amino acid residue.

[00169] In other embodiments, compositions and vaccine compositions, respectively, of the invention will contain HBcAgs from which the C-terminal region (*e.g.*, amino acid residues 145-185 or 150-185 of SEQ ID NO:77) has been removed. Thus, additional modified HBcAgs suitable for use in the practice of the present invention include C-terminal truncation mutants. Suitable truncation mutants include HBcAgs where 1, 5, 10, 15, 20, 25, 30, 34, 35, amino acids have been removed from the C-terminus.

[00170] HBcAgs suitable for use in the practice of the present invention also include N-terminal truncation mutants. Suitable truncation mutants include modified HBcAgs where 1, 2, 5, 7, 9, 10, 12, 14, 15, or 17 amino acids have been removed from the N-terminus.

[00171] Further HBcAgs suitable for use in the practice of the present invention include N- and C-terminal truncation mutants. Suitable truncation mutants include HBcAgs where 1, 2, 5, 7, 9, 10, 12, 14, 15, and 17 amino acids have been removed from the N-terminus and 1, 5, 10, 15, 20, 25, 30, 34, 35 amino acids have been removed from the C-terminus.

[00172] The invention further includes compositions and vaccine compositions, respectively, comprising HBcAg polypeptides comprising, or alternatively essentially consisting of, or alternatively consisting of, amino acid sequences which are at least 80%, 85%, 90%, 95%, 97%, or 99% identical to the above described truncation mutants.

[00173] In certain embodiments of the invention, a lysine residue is introduced into a HBcAg polypeptide, to mediate the binding of the RANKL protein, RANKL fragment or RANKL peptide to the VLP of HBcAg. In preferred embodiments, compositions of the invention are prepared using a HBcAg comprising, or alternatively consisting of, amino acids 1-144, or 1-149, 1-185 of SEQ ID NO:77, which is modified so that the amino acids corresponding to positions 79 and 80 are replaced with a peptide having the amino acid sequence of Gly-Gly-Lys-Gly-Gly (SEQ ID NO:78). In further preferred

- 57 -

embodiments, the cysteine residues at positions 48 and 107 of SEQ ID NO:77 are mutated to serine. The invention further includes compositions comprising the corresponding polypeptides having amino acid sequences shown in any of SEQ ID NOs:29-74, which also have above noted amino acid alterations. Further included within the scope of the invention are additional HBcAg variants which are capable of associating to form a capsid or VLP and have the above noted amino acid alterations. Thus, the invention further includes compositions and vaccine compositions, respectively, comprising HBcAg polypeptides which comprise, or alternatively consist of, amino acid sequences which are at least 80%, 85%, 90%, 95%, 97% or 99% identical to any of the wild-type amino acid sequences, and forms of these proteins which have been processed, where appropriate, to remove the N-terminal leader sequence and modified with above noted alterations.

[00174] Compositions or vaccine compositions of the invention may comprise mixtures of different HBcAgs. Thus, these vaccine compositions may be composed of HBcAgs which differ in amino acid sequence. For example, vaccine compositions could be prepared comprising a "wild-type" HBcAg and a modified HBcAg in which one or more amino acid residues have been altered (*e.g.*, deleted, inserted or substituted). Further, preferred vaccine compositions of the invention are those which present highly ordered and repetitive antigen array, wherein the antigen is a RANKL protein, RANKL fragment or RANKL peptide.

[00175] In a further preferred embodiment of the present invention, the at least one RANKL protein, RANKL fragment or RANKL peptide is bound to said core particle and virus-like particle, respectively, by at least one covalent bond. Preferably, the least one RANKL protein, RANKL fragment or RANKL peptide is bound to the core particle and virus-like particle, respectively, by at least one covalent bond, said covalent bond being a non-peptide bond leading to a core particle-RANKL ordered and repetitive array and a RANKL-VLP-array or -conjugate, respectively. This RANKL-VLP array and conjugate, respectively, has typically and preferably a repetitive and ordered structure

- 58 -

since the at least one, but usually more than one, RANKL protein, RANKL fragment or RANKL peptide is bound to the VLP in an oriented manner. Preferably, more than 10, 20, 40, 80, 120 RANKL proteins, RANKL fragments or RANKL peptides are bound to the VLP. The formation of a repetitive and ordered RANKL-VLP array and conjugate, respectively, is ensured by an oriented and directed as well as defined binding and attachment, respectively, of the at least one RANKL protein, RANKL fragment or RANKL peptide to the VLP as will become apparent in the following. Furthermore, the typical inherent highly repetitive and organized structure of the VLPs advantageously contributes to the display of the RANKL protein, RANKL fragment or RANKL peptide in a highly ordered and repetitive fashion leading to a highly organized and repetitive RANKL-VLP array and conjugate, respectively.

[00176] Therefore, the preferred inventive conjugates and arrays, respectively, differ from prior art conjugates in their highly organized structure, dimensions, and in the repetitiveness of the antigen on the surface of the array. The preferred embodiment of this invention, furthermore, allows expression of both the particle and the antigen in an expression host guaranteeing proper folding of the antigen, i.e. the at least one RANKL protein, RANKL fragment or RANKL peptide, and proper folding and assembly of the VLP.

[00177] The present invention discloses methods of binding of RANKL protein, RANKL fragment or RANKL peptide to core particles and VLPs, respectively. As indicated, in one aspect of the invention, the RANKL protein, RANKL fragment or RANKL peptide is bound to the core particle and VLP, respectively, by way of chemical cross-linking, typically and preferably by using a heterobifunctional cross-linker. Several hetero-bifunctional cross-linkers are known to the art. In preferred embodiments, the hetero-bifunctional cross-linker contains a functional group which can react with preferred first attachment sites, i.e. with the side-chain amino group of lysine residues of the core particle and the VLP or at least one VLP subunit, respectively, and a further functional group which can react with a preferred second attachment

- 59 -

site, i.e. a cysteine residue naturally present, made available for reaction by reduction, or engineered on the RANKL protein, RANKL fragment or RANKL peptide, and optionally also made available for reaction by reduction. The first step of the procedure, typically called the derivatization, is the reaction of the core particle or the VLP with the cross-linker. The product of this reaction is an activated core particle or activated VLP, also called activated carrier. In the second step, unreacted cross-linker is removed using usual methods such as gel filtration or dialysis. In the third step, the RANKL protein, RANKL fragment or RANKL peptide is reacted with the activated carrier, and this step is typically called the coupling step. Unreacted RANKL protein, RANKL fragment or RANKL peptide may be optionally removed in a fourth step, for example by dialysis. Several hetero-bifunctional cross-linkers are known to the art. These include the preferred cross-linkers SMPH (Pierce), Sulfo-MBS, Sulfo-EMCS, Sulfo-GMBS, Sulfo-SIAB, Sulfo-SMPB, Sulfo-SMCC, SVSB, SIA and other cross-linkers available for example from the Pierce Chemical Company (Rockford, IL, USA), and having one functional group reactive towards amino groups and one functional group reactive towards cysteine residues. The above mentioned cross-linkers all lead to formation of a thioether linkage. Another class of cross-linkers suitable in the practice of the invention is characterized by the introduction of a disulfide linkage between the RANKL protein, RANKL fragment or RANKL peptide and the core particle or VLP upon coupling. Preferred cross-linkers belonging to this class include for example SPDP and Sulfo-LC-SPDP (Pierce). The extent of derivatization of the core particle and VLP, respectively, with cross-linker can be influenced by varying experimental conditions such as the concentration of each of the reaction partners, the excess of one reagent over the other, the pH, the temperature and the ionic strength. The degree of coupling, i.e. the amount of RANKL protein, RANKL fragment or RANKL peptides per subunits of the core particle and VLP, respectively, can be adjusted by varying the experimental conditions described above to match the requirements of the vaccine. Solubility of the RANKL protein, RANKL

- 60 -

fragment or RANKL peptide may impose a limitation on the amount of RANKL protein, RANKL fragment or RANKL peptide that can be coupled on each subunit, and in those cases where the obtained vaccine would be insoluble, reducing the amount of RANKL protein, RANKL fragment or RANKL peptides per subunit is beneficial.

[00178] A particularly favored method of binding of RANKL protein, RANKL fragment or RANKL peptides to the core particle and the VLP, respectively, is the linking of a lysine residue on the surface of the core particle and the VLP, respectively, with a cysteine residue on the RANKL protein, RANKL fragment or RANKL peptide. Thus, in a preferred embodiment of the present invention, the first attachment site is a lysine residue and the second attachment site is a cysteine residue. In some embodiments, engineering of an amino acid linker containing a cysteine residue, as a second attachment site or as a part thereof, to the RANKL protein, RANKL fragment or RANKL peptide for coupling to the core particle and VLP, respectively, may be required. Alternatively, a cysteine may be introduced either by insertion or mutation within the RANKL protein, RANKL fragment or RANKL peptide. Alternatively, the cysteine residue or a thiol group may be introduced by chemical coupling.

[00179] The selection of the amino acid linker will be dependent on the nature of the antigen and self-antigen, respectively, i.e. on the nature of the RANKL protein, RANKL fragment or RANKL peptide, on its biochemical properties, such as pI, charge distribution and glycosylation. In general, flexible amino acid linkers are favored. Preferred embodiments of the amino acid linker are selected from the group consisting of: (a) CGG; (b) N-terminal gamma 1-linker; (c) N-terminal gamma 3-linker; (d) Ig hinge regions; (e) N-terminal glycine linkers; (f) $(G)_kC(G)_n$ with $n=0-12$ and $k=0-5$; (g) N-terminal glycine-serine linkers; (h) $(G)_kC(G)_m(S)_l(GGGGS)_n$ with $n=0-3$, $k=0-5$, $m=0-10$, $l=0-2$; (i) GGC; (k) GGC-NH₂; (l) C-terminal gamma 1-linker; (m) C-terminal gamma 3-linker; (n) C-terminal glycine linkers; (o) $(G)_nC(G)_k$ with $n=0-12$

- 61 -

and $k=0-5$; (p) C-terminal glycine-serine linkers; (q) $(G)_m(S)_l(GGGGS)_n(G)_oC(G)_k$ with $n=0-3$, $k=0-5$, $m=0-10$, $l=0-2$, and $o=0-8$.

[00180] Further preferred examples of amino acid linkers are the hinge region of Immunoglobulins, glycine serine linkers $(GGGGS)_n$, and glycine linkers $(G)_n$ all further containing a cysteine residue as second attachment site and optionally further glycine residues. Typically preferred examples of said amino acid linkers are N-terminal gamma1: CGDKTHTSPP; C-terminal gamma 1: DKTHTSPPCG; N-terminal gamma 3: CGGPKPSTPPGSSGGAP; C-terminal gamma 3: PKPSTPPGSSGGAPGGCG; N-terminal glycine linker: GCGGGG; C-terminal glycine linker: GGGGCG; C-terminal glycine-lysine linker: GGKKGC; N-terminal glycine-lysine linker: CGKKGG.

[00181] In a further preferred embodiment of the present invention, and in particular if the antigen is a RANKL peptide, GGCG, GGC or GGC-NH₂ ("NH₂" stands for amidation) linkers at the C-terminus of the peptide or CGG at its N-terminus are preferred as amino acid linkers. In general, glycine residues will be inserted between bulky amino acids and the cysteine to be used as second attachment site, to avoid potential steric hindrance of the bulkier amino acid in the coupling reaction.

[00182] The cysteine residue present on the RANKL protein, RANKL fragment or RANKL peptide has to be in its reduced state to react with the hetero-bifunctional cross-linker on the activated VLP, that is a free cysteine or a cysteine residue with a free sulfhydryl group has to be available. In the instance where the cysteine residue to function as binding site is in an oxidized form, for example if it is forming a disulfide bridge, reduction of this disulfide bridge with *e.g.* DTT, TCEP or β -mercaptoethanol is required.

[00183] Binding of the RANKL protein, RANKL fragment or RANKL peptide to the core particle and VLP, respectively, by using a hetero-bifunctional cross-linker according to the preferred methods described above, allows coupling of the RANKL protein, RANKL fragment or RANKL peptide to the core particle and the VLP, respectively, in an oriented fashion. Other methods of binding the RANKL protein, RANKL fragment or RANKL peptide to the

- 62 -

core particle and the VLP, respectively, include methods wherein the RANKL protein, RANKL fragment or RANKL peptide is cross-linked to the core particle and the VLP, respectively, using the carbodiimide EDC, and NHS. The RANKL protein, RANKL fragment or RANKL peptide may also be first thiolated through reaction, for example with SATA, SATP or iminothiolane. The RANKL protein, RANKL fragment or RANKL peptide, after deprotection if required, may then be coupled to the core particle and the VLP, respectively, as follows. After separation of the excess thiolation reagent, the RANKL protein, RANKL fragment or RANKL peptide is reacted with the core particle and the VLP, respectively, previously activated with a hetero-bifunctional cross-linker comprising a cysteine reactive moiety, and therefore displaying at least one or several functional groups reactive towards cysteine residues, to which the thiolated RANKL protein, RANKL fragment or RANKL peptide can react, such as described above. Optionally, low amounts of a reducing agent are included in the reaction mixture. In further methods, the RANKL protein, RANKL fragment or RANKL peptide is attached to the core particle and the VLP, respectively, using a homo-bifunctional cross-linker such as glutaraldehyde, DSG, BM[PEO]₄, BS³, (Pierce Chemical Company, Rockford, IL, USA) or other known homo-bifunctional cross-linkers which functional groups reactive towards amine groups or carboxyl groups of the core particle and the VLP, respectively,.

[00184] In a further embodiment, the RANKL protein, RANKL fragment or RANKL peptide is bound to the core particle and the VLP, respectively, through modification of the carbohydrate moieties present on glycosylated RANKL protein, RANKL fragment or RANKL peptide and subsequent reaction with the core particle and the VLP, respectively. In one embodiment, the glycosylated RANKL protein, RANKL fragment or RANKL peptide is reacted with sodium periodate in a mild oxidation reaction of the carbohydrate moiety, to yield an activated RANKL protein, RANKL fragment or RANKL peptide with one or more aldehyde functional groups. The so activated RANKL protein, RANKL fragment or RANKL peptide is separated from

- 63 -

excess sodium periodate, and further reacted with the core particle and the VLP, respectively, wherein lysine residues of the core particle and the VLP, respectively, or of at least one VLP subunit are reacting with the previously formed aldehyde functional group on the RANKL protein, RANKL fragment or RANKL peptide, for example as described by Hermanson, G.T. in *Bioconjugate Techniques*, Academic Press Inc., San Diego, CA, USA. Self polymerization of the activated RANKL protein, RANKL fragment or RANKL peptide may be controlled by adjusting the pH as described in the aforementioned publication. The formed Schiff base is preferably further reduced with sodium cyanoborohydride, which is subsequently removed by gel filtration or dialysis. Alternatively, the core particle and the VLP, respectively, may be reacted with EDC at carboxyl groups of the core particle and the VLP, respectively, or at least one VLP subunit and a dihydrazide, such as adipic acid dihydrazide, to yield a hydrazide moiety available for reaction with the one or more aldehyde functional groups present on the activated RANKL protein, RANKL fragment or RANKL peptide. The so formed hydrazone may be further reduced with sodium cyanoborohydride. Alternatively, the activated RANKL protein, RANKL fragment or RANKL peptide with one or more aldehyde functional groups is reacted with cysteamine, resulting in the introduction of a cysteine group in the RANKL protein, RANKL fragment or RANKL peptide. Additional cross-linking methods and cross-linkers, suitable for binding a RANKL protein, RANKL fragment or RANKL peptide to a core particle and a VLP, respectively, as well as guidance on performing the coupling reactions and on the use of chemical cross-linkers and chemical cross-linking procedures can be found in Hermanson, G.T. in *Bioconjugate Techniques*, Academic Press Inc., San Diego, CA, USA.

[00185] Other methods of binding the VLP to a RANKL protein, RANKL fragment or RANKL peptide include methods where the core particle and the VLP, respectively, is biotinylated, and the RANKL protein, RANKL fragment or RANKL peptide expressed as a streptavidin-fusion protein, or methods

- 64 -

wherein both the RANKL protein, RANKL fragment or RANKL peptides and the core particle and the VLP, respectively, are biotinylated, for example as described in WO 00/23955. In this case, the RANKL protein, RANKL fragment or RANKL peptide may be first bound to streptavidin or avidin by adjusting the ratio of RANKL protein, RANKL fragment or RANKL peptide to streptavidin such that free binding sites are still available for binding of the core particle and the VLP, respectively,, which is added in the next step. Alternatively, all components may be mixed in a "one pot" reaction. Other ligand-receptor pairs, where a soluble form of the receptor and of the ligand is available, and are capable of being cross-linked to the core particle and the VLP, respectively, or the RANKL protein, RANKL fragment or RANKL peptide, may be used as binding agents for binding the RANKL protein, RANKL fragment or RANKL peptide to the core particle and the VLP, respectively,. Alternatively, either the ligand or the receptor may be fused to the RANKL protein, RANKL fragment or RANKL peptide, and so mediate binding to the core particle and the VLP, respectively, chemically bound or fused either to the receptor, or the ligand respectively. Fusion may also be effected by insertion or substitution.

[00186] As already indicated, in a favored embodiment of the present invention, the VLP is the VLP of a RNA phage, and in a more preferred embodiment, the VLP is the VLP of RNA phage Q β coat protein.

[00187] One or several antigen molecules, i.e. a RANKL protein, RANKL fragment or RANKL peptide, can be attached to one subunit of the capsid or VLP of RNA phages coat proteins, preferably through the exposed lysine residues of the VLP of RNA phages, if sterically allowable. A specific feature of the VLP of the coat protein of RNA phages and in particular of the Q β coat protein VLP is thus the possibility to couple several antigens per subunit. This allows for the generation of a dense antigen array.

[00188] In a preferred embodiment of the invention, the binding and attachment, respectively, of the at least RANKL protein, RANKL fragment or RANKL peptide to the core particle and the virus-like particle, respectively, is

- 65 -

by way of interaction and association, respectively, between at least one first attachment site of the virus-like particle and at least one second attachment of the antigen or antigenic determinant.

[00189] VLPs or capsids of Q β coat protein display a defined number of lysine residues on their surface, with a defined topology with three lysine residues pointing towards the interior of the capsid and interacting with the RNA, and four other lysine residues exposed to the exterior of the capsid. These defined properties favor the attachment of antigens to the exterior of the particle, rather than to the interior of the particle where the lysine residues interact with RNA. VLPs of other RNA phage coat proteins also have a defined number of lysine residues on their surface and a defined topology of these lysine residues.

[00190] In further preferred embodiments of the present invention, the first attachment site is a lysine residue and/or the second attachment comprises sulfhydryl group or a cysteine residue. In a very preferred embodiment of the present invention, the first attachment site is a lysine residue and the second attachment is a cysteine residue.

[00191] In very preferred embodiments of the invention, the RANKL protein, RANKL fragment or RANKL peptide is bound via a cysteine residue, either naturally present on the RANKL protein, RANKL fragment or RANKL peptide or engineered, to lysine residues of the VLP of RNA phage coat protein, and in particular to the VLP of Q β coat protein.

[00192] Another advantage of the VLPs derived from RNA phages is their high expression yield in bacteria that allows production of large quantities of material at affordable cost.

[00193] As indicated, the inventive conjugates and arrays, respectively, differ from prior art conjugates in their highly organized structure, dimensions, and in the repetitiveness of the antigen on the surface of the array. Moreover, the use of the VLPs as carriers allow the formation of robust antigen arrays and conjugates, respectively, with variable antigen density. In particular, the use of VLPs of RNA phages, and hereby in particular the use of the VLP of RNA

- 66 -

phage Q β coat protein allows to achieve very high epitope density. The preparation of compositions of VLPs of RNA phage coat proteins with a high epitope density can be effected by using the teaching of this application.

[00194] The second attachment site, as defined herein, may be either naturally or non-naturally present with the antigen or the antigenic determinant. In the case of the absence of a suitable natural occurring second attachment site on the antigen or antigenic determinant, such a, then non-natural second attachment has to be engineered to the antigen.

[00195] As described above, four lysine residues are exposed on the surface of the VLP of Q β coat protein. Typically these residues are derivatized upon reaction with a cross-linker molecule. In the instance where not all of the exposed lysine residues can be coupled to an antigen, the lysine residues which have reacted with the cross-linker are left with a cross-linker molecule attached to the ϵ -amino group after the derivatization step. This leads to disappearance of one or several positive charges, which may be detrimental to the solubility and stability of the VLP. By replacing some of the lysine residues with arginines, as in the disclosed Q β coat protein mutants described below, we prevent the excessive disappearance of positive charges since the arginine residues do not react with the cross-linker. Moreover, replacement of lysine residues by arginines may lead to more defined antigen arrays, as fewer sites are available for reaction to the antigen.

[00196] Accordingly, exposed lysine residues were replaced by arginines in the following Q β coat protein mutants and mutant Q β VLPs disclosed in this application: Q β -240 (Lys13-Arg; SEQ ID NO:23), Q β -250 (Lys 2-Arg, Lys13-Arg; SEQ ID NO:25) and Q β -259 (Lys 2-Arg, Lys16-Arg; SEQ ID NO:27). The constructs were cloned, the proteins expressed, the VLPs purified and used for coupling to peptide and protein antigens. Q β -251 ; (SEQ ID NO:26) was also constructed, and guidance on how to express, purify and couple the VLP of Q β -251 coat protein can be found throughout the application.

- 67 -

[00197] In a further embodiment, we disclose a Q β mutant coat protein with one additional lysine residue, suitable for obtaining even higher density arrays of antigens. This mutant Q β coat protein, Q β -243 (Asn 10-Lys; SEQ ID NO:24), was cloned, the protein expressed, and the capsid or VLP isolated and purified, showing that introduction of the additional lysine residue is compatible with self-assembly of the subunits to a capsid or VLP. Thus, RANKL protein, RANKL fragment or RANKL peptide arrays and conjugates, respectively, may be prepared using VLP of Q β coat protein mutants. A particularly favored method of attachment of antigens to VLPs, and in particular to VLPs of RNA phage coat proteins is the linking of a lysine residue present on the surface of the VLP of RNA phage coat proteins with a cysteine residue naturally present or engineered on the antigen, i.e. the RANKL protein, RANKL fragment or RANKL peptide. In order for a cysteine residue to be effective as second attachment site, a sulfhydryl group must be available for coupling. Thus, a cysteine residue has to be in its reduced state, that is, a free cysteine or a cysteine residue with a free sulfhydryl group has to be available. In the instant where the cysteine residue to function as second attachment site is in an oxidized form, for example if it is forming a disulfide bridge, reduction of this disulfide bridge with *e.g.* DTT, TCEP or β -mercaptoethanol is required. The concentration of reductant, and the molar excess of reductant over antigen has to be adjusted for each antigen. A titration range, starting from concentrations as low as 10 μ M or lower, up to 10 to 20 mM or higher reductant if required is tested, and coupling of the antigen to the carrier assessed. Although low concentrations of reductant are compatible with the coupling reaction as described in pending U.S. Application No. 10/050,902 filed by the present assignee on January 18, 2002, higher concentrations inhibit the coupling reaction, as a skilled artisan would know, in which case the reductant has to be removed by dialysis or gel filtration. Advantageously, the pH of the dialysis or equilibration buffer is lower than 7, preferably 6. The compatibility of the low pH buffer with antigen activity or stability has to be tested.

- 68 -

[00198] Epitope density on the VLP of RNA phage coat proteins can be modulated by the choice of cross-linker and other reaction conditions. For example, the cross-linkers Sulfo-GMBS and SMPH typically allow reaching high epitope density. Derivatization is positively influenced by high concentration of reactands, and manipulation of the reaction conditions can be used to control the number of antigens coupled to VLPs of RNA phage coat proteins, and in particular to VLPs of Q β coat protein.

[00199] Prior to the design of a non-natural second attachment site the position at which it should be fused, inserted or generally engineered has to be chosen. The selection of the position of the second attachment site may, by way of example, be based on a crystal structure of the antigen. Such a crystal structure of the antigen may provide information on the availability of the C- or N-termini of the molecule (determined for example from their accessibility to solvent), or on the exposure to solvent of residues suitable for use as second attachment sites, such as cysteine residues. Exposed disulfide bridges, as is the case for Fab fragments, may also be a source of a second attachment site, since they can be generally converted to single cysteine residues through mild reduction. Mild reduction conditions not affecting the immunogenicity of RANKL protein, RANKL fragment or RANKL peptide will be chosen. In general, in the case where immunization with a self-antigen is aiming at inhibiting the interaction of this self-antigen with its natural ligands, the second attachment site will be added such that it allows generation of antibodies against the site of interaction with the natural ligands. Thus, the location of the second attachment site will be selected such that steric hindrance from the second attachment site or any amino acid linker containing the same is avoided. In further embodiments, an antibody response directed at a site distinct from the interaction site of the self-antigen with its natural ligand is desired. In such embodiments, the second attachment site may be selected such that it prevents generation of antibodies against the interaction site of the self-antigen with its natural ligands.

- 69 -

[00200] Other criteria in selecting the position of the second attachment site include the oligomerization state of the antigen, the site of oligomerization, the presence of a cofactor, and the availability of experimental evidence disclosing sites in the antigen structure and sequence where modification of the antigen is compatible with the function of the self-antigen, or with the generation of antibodies recognizing the self-antigen.

[00201] In the most preferred embodiments, the RANKL protein, RANKL fragment or RANKL peptide comprises a single second attachment site or a single reactive attachment site capable of association with the first attachment sites on the core particle and the VLPs or VLP subunits, respectively. This ensures a defined and uniform binding and association, respectively, of the at least one, but typically more than one, preferably more than 10, 20, 40, 80, 120 antigens to the core particle and VLP, respectively. The provision of a single second attachment site or a single reactive attachment site on the antigen, thus, ensures a single and uniform type of binding and association, respectively leading to a very highly ordered and repetitive array. For example, if the binding and association, respectively, is effected by way of a lysine- (as the first attachment site) and cysteine- (as a second attachment site) interaction, it is ensured, in accordance with this preferred embodiment of the invention, that only one cysteine residue per antigen, independent whether this cysteine residue is naturally or non-naturally present on the antigen, is capable of binding and associating, respectively, with the VLP and the first attachment site of the core particle, respectively.

[00202] In some embodiments, engineering of a second attachment site onto the antigen require the fusion of an amino acid linker containing an amino acid suitable as second attachment site according to the disclosures of this invention. Therefore, in a preferred embodiment of the present invention, an amino acid linker is bound to the antigen or the antigenic determinant by way of at least one covalent bond. Preferably, the amino acid linker comprises, or alternatively consists of, the second attachment site. In a further preferred embodiment, the amino acid linker comprises a sulfhydryl group or a cysteine

- 70 -

residue. In another preferred embodiment, the amino acid linker is cysteine. Some criteria of selection of the amino acid linker as well as further preferred embodiments of the amino acid linker according to the invention have already mentioned above.

[00203] In a further preferred embodiment of the invention, the at least one antigen or antigenic determinant, i.e. the RANKL protein, RANKL fragment or RANKL peptide arrays is fused to the core particle and the virus-like particle, respectively. As outlined above, a VLP is typically composed of at least one subunit assembling into a VLP. Thus, in again a further preferred embodiment of the invention, the antigen or antigenic determinant, preferably the at least one RANKL protein, RANKL fragment or RANKL peptide, is fused to at least one subunit of the virus-like particle or of a protein capable of being incorporated into a VLP generating a chimeric VLP-subunit-RANKL protein, RANKL fragment or RANKL peptide fusion.

[00204] Fusion of the RANKL protein, RANKL fragment or RANKL peptide can be effected by insertion into the VLP subunit sequence, or by fusion to either the N- or C-terminus of the VLP-subunit or protein capable of being incorporated into a VLP. Hereinafter, when referring to fusion proteins of a peptide to a VLP subunit, the fusion to either ends of the subunit sequence or internal insertion of the peptide within the subunit sequence are encompassed.

[00205] Fusion may also be effected by inserting the RANKL protein, RANKL fragment or RANKL peptide sequences into a variant of a VLP subunit where part of the subunit sequence has been deleted, that are further referred to as truncation mutants. Truncation mutants may have N- or C-terminal, or internal deletions of part of the sequence of the VLP subunit. For example, the specific VLP HBcAg with, for example, deletion of amino acid residues 79 to 81 is a truncation mutant with an internal deletion. Fusion of RANKL protein, RANKL fragment or RANKL peptide to either the N- or C-terminus of the truncation mutants VLP-subunits also lead to embodiments of the invention. Likewise, fusion of an epitope into the sequence of the VLP subunit may also be effected by substitution, where for example for the

- 71 -

specific VLP HBcAg, amino acids 79-81 are replaced with a foreign epitope. Thus, fusion, as referred to hereinafter, may be effected by insertion of the RANKL protein, RANKL fragment or RANKL peptide sequence in the sequence of a VLP subunit, by substitution of part of the sequence of the VLP subunit with the RANKL protein, RANKL fragment or RANKL peptide sequence, or by a combination of deletion, substitution or insertions.

[00206] The chimeric RANKL protein, RANKL fragment or RANKL peptide-VLP subunit will be in general capable of self-assembly into a VLP. VLP displaying epitopes fused to their subunits are also herein referred to as chimeric VLPs. As indicated, the virus-like particle comprises or alternatively is composed of at least one VLP subunit. In a further embodiment of the invention, the virus-like particle comprises or alternatively is composed of a mixture of chimeric VLP subunits and non-chimeric VLP subunits, i.e. VLP subunits not having an antigen fused thereto, leading to so called mosaic particles. This may be advantageous to ensure formation of and assembly to a VLP. In those embodiments, the proportion of chimeric VLP-subunits may be 1, 2, 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, 95% or higher.

[00207] Flanking amino acid residues may be added to either end of the sequence of the peptide or epitope to be fused to either end of the sequence of the subunit of a VLP, or for internal insertion of such peptidic sequence into the sequence of the subunit of a VLP. Glycine and serine residues are particularly favored amino acids to be used in the flanking sequences added to the RANKL protein, RANKL fragment or RANKL peptide to be fused. Glycine residues confer additional flexibility, which may diminish the potentially destabilizing effect of fusing a foreign sequence into the the sequence of a VLP subunit.

[00208] In a specific embodiment of the invention, the VLP is a Hepatitis B core antigen VLP. Fusion proteins to either the N-terminus of a HBcAg (Neyrinck, S. *et al.*, *Nature Med.* 5:1157-1163 (1999)) or insertions in the so called major immunodominant region (MIR) have been described (Pumpens, P. and Grens, E., *Intervirology* 44:98-114 (2001)), WO 01/98333, and are

- 72 -

preferred embodiments of the invention. Naturally occurring variants of HBcAg with deletions in the MIR have also been described (Pumpens, P. and Grens, E., *Intervirology* 44:98-114 (2001), which is expressly incorporated by reference in their entirety), and fusions to the N- or C-terminus, as well as insertions at the position of the MIR corresponding to the site of deletion as compared to a wt HBcAg are further embodiments of the invention. Fusions to the C-terminus have also been described (Pumpens, P. and Grens, E., *Intervirology* 44:98-114 (2001)). One skilled in the art will easily find guidance on how to construct fusion proteins using classical molecular biology techniques (Sambrook, J. et al., eds., *Molecular Cloning, A Laboratory Manual*, 2nd. edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), Ho et al., *Gene* 77:51 (1989)). Vectors and plasmids encoding HBcAg and HBcAg fusion proteins and useful for the expression of a HBcAg and HBcAg fusion proteins have been described (Pumpens, P. & Grens, E. *Intervirology* 44: 98-114 (2001), Neyrinck, S. et al., *Nature Med.* 5:1157-1163 (1999)) and can be used in the practice of the invention. We also describe by way of example (Example 6) the insertion of an epitope into the MIR of HBcAg, resulting in a chimeric self-assembling HBcAg. An important factor for the optimization of the efficiency of self-assembly and of the display of the epitope to be inserted in the MIR of HBcAg is the choice of the insertion site, as well as the number of amino acids to be deleted from the HBcAg sequence within the MIR (Pumpens, P. and Grens, E., *Intervirology* 44:98-114 (2001); EP 421'635; US 6'231'864) upon insertion, or in other words, which amino acids form HBcAg are to be substituted with the new epitope. For example, substitution of HBcAg amino acids 76-80, 79-81, 79-80, 75-85 or 80-81 with foreign epitopes has been described (Pumpens, P. and Grens, E., *Intervirology* 44:98-114 (2001); EP0421635; US 6'231'864). HBcAg contains a long arginine tail (Pumpens, P. and Grens, E., *Intervirology* 44:98-114 (2001)) which is dispensable for capsid assembly and capable of binding nucleic acids (Pumpens, P. and Grens, E., *Intervirology* 44:98-114

- 73 -

(2001)). HBcAg either comprising or lacking this arginine tail are both embodiments of the invention.

[00209] In a further preferred embodiment of the invention, the VLP is a VLP of a RNA phage. The major coat proteins of RNA phages spontaneously assemble into VLPs upon expression in bacteria, and in particular in *E. coli*. Specific examples of bacteriophage coat proteins which can be used to prepare compositions of the invention include the coat proteins of RNA bacteriophages such as bacteriophage Q β (SEQ ID NO:10; PIR Database, Accession No. VCBPQ β referring to Q β CP and SEQ ID NO: 11; Accession No. AAA16663 referring to Q β A1 protein) and bacteriophage fr (SEQ ID NO:4; PIR Accession No. VCBPFR).

[00210] In a more preferred embodiment, the at least one RANKL protein, RANKL fragment or RANKL peptide is fused to a Q β coat protein. Fusion protein constructs wherein epitopes have been fused to the C-terminus of a truncated form of the A1 protein of Q β , or inserted within the A1 protein have been described (Kozlovskaja, T. M., *et al.*, *Intervirology*, 39:9-15 (1996)). The A1 protein is generated by suppression at the UGA stop codon and has a length of 329 aa, or 328 aa, if the cleavage of the N-terminal methionine is taken into account. Cleavage of the N-terminal methionine before an alanine (the second amino acid encoded by the Q β CP gene) usually takes place in *E. coli*, and such is the case for N-termini of the Q β coat proteins CP. The part of the A1 gene, 3' of the UGA amber codon encodes the CP extension, which has a length of 195 amino acids. Insertion of the at least one RANKL protein, RANKL fragment or RANKL peptide between position 72 and 73 of the CP extension leads to further embodiments of the invention (Kozlovskaja, T. M., *et al.*, *Intervirology* 39:9-15 (1996)). Fusion of a RANKL protein, RANKL fragment or RANKL peptide at the C-terminus of a C-terminally truncated Q β A1 protein leads to further preferred embodiments of the invention. For example, Kozlovskaja *et al.*, (*Intervirology*, 39: 9-15 (1996)) describe Q β A1

- 74 -

protein fusions where the epitope is fused at the C-terminus of the Q β CP extension truncated at position 19.

[00211] As described by Kozlovskaja *et al.* (*Intervirology*, 39: 9-15 (1996)), assembly of the particles displaying the fused epitopes typically requires the presence of both the A1 protein-RANKL protein, RANKL fragment or RANKL peptide fusion and the wt CP to form a mosaic particle. However, embodiments comprising virus-like particles, and hereby in particular the VLPs of the RNA phage Q β coat protein, which are exclusively composed of VLP subunits having at least one RANKL protein, RANKL fragment or RANKL peptide fused thereto, are also within the scope of the present invention.

[00212] The production of mosaic particles may be effected in a number of ways. Kozlovskaja *et al.*, *Intervirology*, 39:9-15 (1996), describe two methods, which both can be used in the practice of the invention. In the first approach, efficient display of the fused epitope on the VLPs is mediated by the expression of the plasmid encoding the Q β A1 protein fusion having a UGA stop codon between CP and CP extension in a *E. coli* strain harboring a plasmid encoding a cloned UGA suppressor tRNA which leads to translation of the UGA codon into Trp (pISM3001 plasmid (Smiley B.K., *et al.*, *Gene* 134:33-40 (1993))). In another approach, the CP gene stop codon is modified into UAA, and a second plasmid expressing the A1 protein-RANKL protein, RANKL fragment or RANKL peptide fusion is cotransformed. The second plasmid encodes a different antibiotic resistance and the origin of replication is compatible with the first plasmid (Kozlovskaja, T. M., *et al.*, *Intervirology* 39:9-15 (1996)). In a third approach, CP and the A1 protein-RANKL protein, RANKL fragment or RANKL peptide fusion are encoded in a bicistronic manner, operatively linked to a promoter such as the Trp promoter, as described in FIG. 1 of Kozlovskaja *et al.*, *Intervirology*, 39:9-15 (1996).

[00213] In a further embodiment, the RANKL protein, RANKL fragment or RANKL peptide is inserted between amino acid 2 and 3 (numbering of the cleaved CP, that is wherein the N-terminal methionine is cleaved) of the fr CP,

thus leading to a RANKL protein, RANKL fragment or RANKL peptide -fr CP fusion protein. Vectors and expression systems for construction and expression of fr CP fusion proteins self-assembling to VLP and useful in the practice of the invention have been described (Pushko P. *et al.*, *Prot. Eng.* 6:883-891 (1993)). In a specific embodiment, the RANKL protein, RANKL fragment or RANKL peptide sequence is inserted into a deletion variant of the fr CP after amino acid 2, wherein residues 3 and 4 of the fr CP have been deleted (Pushko P. *et al.*, *Prot. Eng.* 6:883-891 (1993)).

[00214] Fusion of epitopes in the N-terminal protuberant β -hairpin of the coat protein of RNA phage MS-2 and subsequent presentation of the fused epitope on the self-assembled VLP of RNA phage MS-2 has also been described (WO 92/13081), and fusion of RANKL protein, RANKL fragment or RANKL peptide by insertion or substitution into the coat protein of MS-2 RNA phage is also falling under the scope of the invention.

[00215] In another embodiment of the invention, the RANKL protein, RANKL fragment or RANKL peptide are fused to a capsid protein of papillomavirus. In a more specific embodiment, the RANKL protein, RANKL fragment or RANKL peptide are fused to the major capsid protein L1 of bovine papillomavirus type 1 (BPV-1). Vectors and expression systems for construction and expression of BPV-1 fusion proteins in a baculovirus/insect cells systems have been described (Chackerian, B. *et al.*, *Proc. Natl. Acad. Sci. USA* 96:2373-2378 (1999), WO 00/23955). Substitution of amino acids 130-136 of BPV-1 L1 with a RANKL protein, RANKL fragment or RANKL peptide leads to a BPV-1 L1- RANKL protein, RANKL fragment or RANKL peptide fusion protein, which is a preferred embodiment of the invention. Cloning in a baculovirus vector and expression in baculovirus infected Sf9 cells has been described, and can be used in the practice of the invention (Chackerian, B. *et al.*, *Proc. Natl. Acad. Sci. USA* 96:2373-2378 (1999), WO 00/23955). Purification of the assembled particles displaying the fused RANKL protein, RANKL fragment or RANKL peptide can be performed in a number of ways, such as for example gel filtration or sucrose gradient

- 76 -

ultracentrifugation (Chackerian, B. *et al.*, *Proc. Natl. Acad. Sci. USA* 96:2373-2378 (1999), WO 00/23955).

[00216] In a further embodiment of the invention, the RANKL protein, RANKL fragment or RANKL peptide are fused to a Ty protein capable of being incorporated into a Ty VLP. In a more specific embodiment, the RANKL protein, RANKL fragment or RANKL peptide are fused to the p1 or capsid protein encoded by the TYA gene (Roth, J.F., *Yeast* 16:785-795 (2000)). The yeast retrotransposons Ty1, 2, 3 and 4 have been isolated from *Saccharomyces Serevisiae*, while the retrotransposon Tfl has been isolated from *Schizosaccharomyces Pombae* (Boeke, J.D. and Sandmeyer, S.B., "Yeast Transposable elements," in *The molecular and Cellular Biology of the Yeast Saccharomyces: Genome dynamics, Protein Synthesis, and Energetics.*, p. 193, Cold Spring Harbor Laboratory Press (1991)). The retrotransposons Ty1 and 2 are related to the *copia* class of plant and animal elements, while Ty3 belongs to the *gypsy* family of retrotransposons, which is related to plants and animal retroviruses. In the Ty1 retrotransposon, the p1 protein, also referred to as Gag or capsid protein, has a length of 440 amino acids. P1 is cleaved during maturation of the VLP at position 408, leading to the p2 protein, the essential component of the VLP.

[00217] Fusion proteins to p1 and vectors for the expression of said fusion proteins in Yeast have been described (Adams, S.E., *et al.*, *Nature* 329:68-70 (1987)). So, for example, a RANKL protein, RANKL fragment or RANKL peptide may be fused to p1 by inserting a sequence coding for the RANKL protein, RANKL fragment or RANKL peptide into the BamH1 site of the pMA5620 plasmid (Adams, S.E., *et al.*, *Nature* 329:68-70 (1987)). The cloning of sequences coding for foreign epitopes into the pMA5620 vector leads to expression of fusion proteins comprising amino acids 1-381 of p1 of Ty1-15, fused C-terminally to the N-terminus of the foreign epitope. Likewise, N-terminal fusion of RANKL protein, RANKL fragment or RANKL peptide s, or internal insertion into the p1 sequence, or substitution of part of the p1 sequence is also meant to fall within the scope of the invention. In particular,

- 77 -

insertion of RANKL protein, RANKL fragment or RANKL peptide into the Ty sequence between amino acids 30-31, 67-68, 113-114 and 132-133 of the Ty protein p1 (EP0677111) leads to preferred embodiments of the invention.

[00218] Further VLPs suitable for fusion of RANKL protein, RANKL fragment or RANKL peptide are, for example, Retrovirus-like-particles (WO9630523), HIV2 Gag (Kang, Y.C., *et al*, *Biol. Chem.* 380:353-364 (1999)), Cowpea Mosaic Virus (Taylor, K.M.*et al.*, *Biol. Chem.* 380:387-392 (1999)), parvovirus VP2 VLP (Rueda, P. *et al.*, *Virology* 263:89-99 (1999)), HBsAg (US 4,722,840, EP0020416B1).

[00219] Examples of chimeric VLPs suitable for the practice of the invention are also those described in *Intervirology* 39:1 (1996). Further examples of VLPs contemplated for use in the invention are: HPV-1, HPV-6, HPV-11, HPV-16, HPV-18, HPV-33, HPV-45, CRPV, COPV, HIV GAG, Tobacco Mosaic Virus. Virus-like particles of SV-40, Polyomavirus, Adenovirus, Herpes Simplex Virus, Rotavirus and Norwalk virus have also been made, and chimeric VLPs of those VLPs are also within the scope of the present invention.

[00220] In a further preferred embodiment of the present invention, the antigen or antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide.

[00221] In a further very preferred embodiment of the invention, the antigen or antigenic determinant is a human RANKL protein, RANKL fragment or RANKL peptide.

[00222] In a further very preferred embodiment of the invention, the antigen or antigenic determinant comprises, alternatively essentially consists of, or alternatively consists of an amino acid sequence selected from the group consisting of a) the amino acid sequence of SEQ ID NO: 79; b) the amino acid sequence of any fragment of SEQ ID NO: 79.

[00223] In a further preferred embodiment of the invention, the antigen or antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide variant, e.g. in particular containing amino acid substitutions or

- 78 -

peptide insertions or polymorphisms. As already indicated, compositions and vaccine compositions, respectively, comprising RANKL protein, RANKL fragment or RANKL peptide variants are included within the scope of the present invention.

[00224] In a further preferred embodiment of the invention, the antigen or antigenic determinant is a RANKL fragment. Preferably, the antigen or antigenic determinant is human RANKL fragment selected from the group of a) the extracellular region of human RANKL, b) the splice isoform 1 of human RANKL, c) the splice isoform 2 resulting in a secreted human RANKL, d) the proteolytically produced soluble region of human RANKL, and e) the TNF- α homolog region. .

[00225] In a further very preferred embodiment of the invention, the antigen or antigenic determinant comprises, alternatively essentially consists of, or alternatively consists of an amino acid sequence selected from the group consisting a) the amino acid sequence of SEQ ID NO: 79; b) the amino acid sequence of SEQ ID NO: 80; c) the amino acid sequence of SEQ ID NO: 81; d) the amino acid sequence of SEQ ID NO: 82; e) the amino acid sequence of SEQ ID NO: 83; f) the amino acid sequence of SEQ ID NO: 84; g) the amino acid sequence of SEQ ID NO: 100; h) the amino acid sequence of SEQ ID NO: 101; i) the amino acid sequence of any fragment of any of SEQ ID NO:79-84,100,101.

[00226] RANKL protein and RANKL fragments can be produced by expression of the RANKL cDNA in procaryotic or eucaryotic expression systems. Various examples hereto have been described in the literature and can be used, possibly after modifications, to express any RANKL protein, RANKL fragment or RANKL peptide of any desired species. RANKL protein and RANKL fragments have been expressed in mammalian cells (Anderson, D. M., *et al.*, *Nature* 390: 175-179 (1997), Lacey, D.L., *et al.*, *Cell* 93: 165-176 (1998), Wong B.R., *et al.*, *J. Biol. Chem.* 272: 25190-25194 (1997), Lum, L., *et al.*, *J. Biol. Chem.* 274: 13613-13618 (2000)), in insect cells (Willard, D., *et al.*, *Prot. Express. Purif.* 20: 48-57 (2000)), and procaryotic cells (Xu,

- 79 -

J., et al., *J. Bone Mineral Res.* 15: 2178-86 (2000), Yasuda *et al.*, *Proc. Natl. Acad. Sci USA* 95: 3597-3602 (1998)). Disclosures how to produce RANKL proteins and fragments are also given in WO 9846751, US 5,843,678, WO 98259958, US 6,242,586, WO 9828426, US 6,242,213, WO 9929865, JP 2000102390 and WO 0015807.

[00227] In a further preferred embodiment of the invention, the antigen or antigenic determinant is a RANKL peptide. Such RANKL peptides or fragments thereof can be produced using standard molecular biological technologies where the nucleotide sequence coding for the fragment of interest is amplified by PCR and is cloned as a fusion to a polypeptide tag, such as the histidine tag, the Flag tag, myc tag or the constant region of an antibody (Fc region). By introducing a protease cleavage site between the RANKL fragment and the tag, the RANKL fragment can be separated from the tag after purification by digestion with corresponding protease. In another approach the RANKL fragment can be synthesized in vitro using standard peptide synthesis reactions known to a person skilled in the art. In a further approach RANKL peptides or RANKL fragments can be produced by protease digestion or chemical cleavage of the full length RANKL protein or RANKL fragments, both methods which are well known to people trained in the art.

[00228] In a still further preferred embodiment of the present invention, the antigen or antigenic determinant further comprise at least one second attachment site being selected from the group consisting of: (i) an attachment site not naturally occurring with said antigen or antigenic determinant; and (ii) an attachment site naturally occurring with said antigen or antigenic determinant. Guidance on how to modify RANKL protein, RANKL fragment or RANKL peptide for binding to the virus-like particle is given throughout the application. Preferred second attachment sites contain a cysteine residue for binding to the derivatized VLP and examples are given in the above description and in Example 12 and 13.

[00229] We have built a model for the 3-dimensional structure of the region of human RANKL that is homologous to TNF- α . We found that the naturally

- 80 -

occurring cysteine may not be accessible in the folded structure for interaction with a first attachment site on the VLP in accordance with the present invention. Our model was confirmed by the X-ray structure of mouse RANKL that was recently solved (Lam, J., *et al.*, *J. Clin. Invest.*, 108: 971-979 (2002)). The N-terminus is preferred for attaching a second attachment site comprising an amino acid linker with an additional cysteine residue as shown in Example 12. However, an amino-acid linker containing a cysteine residue as second attachment site and being fused at the C-terminus of the RANKL construct leads to a further preferred embodiment of the invention as shown in EXAMPLE 13. A human RANKL construct with an N-terminal amino acid linker containing a cysteine residue fused to the extracellular region of RANKL is a very preferred embodiment of the invention.

[00230] Mouse RANKL fragment constructs (SEQ ID NO:96 and SEQ ID NO:99) are disclosed, and preferred human RANKL fragment constructs can also be generated and have, for example, the sequence of SEQ ID NO:100-104. Further preferred constructs comprise the whole human RANKL protein, a human RANKL fragment selected from the group of a) the extracellular region of RANKL (SEQ ID NO:82), b) the splice isoform 1 of RANKL (SEQ ID NO: 80), c) the splice isoform 2 resulting in a secreted RANKL (SEQ ID NO:81), d) the proteolytically produced soluble region of RANKL (SEQ ID NO:83), and e) the TNF- α homolog region (SEQ ID NO:84) or human RANKL peptide sequences. Immunization against RANKL protein, RANKL fragment or RANKL peptide using the inventive compositions comprising, preferably a human RANKL protein, RANKL fragment or RANKL peptide bound to a VLP may provide a way of treatment or prevention of bone diseases.

[00231] In a further preferred embodiment of the present invention, the RANKL protein, RANKL fragment or RANKL peptide comprises at least one antigenic site of a RANKL protein. The skilled person in the art knows how to identify the corresponding peptides and amino acid sequences, respectively.

[00232] In a further preferred embodiment of the present invention, the antigen or antigenic determinant is a RANKL peptide that is crucial for interaction with the receptor RANK. Our modeling of the human RANKL structure and the published crystal structure of the mouse RANKL showed that the RANKL monomer consists of a β -sandwich, composed of two flat antiparallel β -sheets. The first sheet is formed by β -strands A'', A, H, C and F while the second sheet is formed by β -strands B', B, G, D, and E. The inner A''AHCF β -sheet is involved in intersubunit association, whereas the B'BGDE β -sheet contributes largely to the outer surface. β -strands are connected via the AA'' loop, the CD loop, the DE loop, the EF loop. The homotrimer is assembled such that one edge of the β -sandwich in each RANKL monomer packs against the inner hydrophobic face of the AHCF β -sheet of the neighbouring monomer. The RANK binding site is thought to encompass the cleft formed by neighbouring monomers of the homotrimer. Based on the homology between the mouse and the human sequence peptides are selected which encompass the RANK binding site. In a preferred embodiment peptides from the interaction site of RANKL with RANK are selected from the group consisting of a) the AA'' loop encompassing amino acids 171-194 (SEQ ID NO: 87), b) the DE loop encompassing amino acids 246-253 (SEQ ID NO:88), c) the β -strand D encompassing amino acids 235-245 (SEQ ID NO: 89), d) the CD loop encompassing amino acids 223-234 (SEQ ID NO:90), e) the EF loop encompassing 262-272 (SEQ ID NO:91), f) β -strand B'-loop B'B encompassing amino acids 200-207 (SEQ ID NO:92), g) loop GH encompassing amino acids 300-305 (SEQ ID NO:93), h) any fragment of said peptides a-g, i) any N- and/or C-terminal extensions of said peptides a-g of at least one amino acid and up to 25 amino acids, j) any fusion of peptides a-i). Further RANKL peptides suitable for use in the present invention can be experimentally determined by their intrinsic property to induce a T cell or an antibody response. This is generally achieved by immunizing an experimental animal separately with selected peptides in an immunologically suitable

- 82 -

formulation and by measuring T cell and B cell, i.e. antibody responses, using methods known to a person trained in the art. In the case where the antigen is a protein, a polypeptide or a peptide, this region can be formed by a continuous amino acid sequence. Alternatively, the antibody epitope can be formed by a discontinuous amino acid sequence in which, after three dimensional folding of the protein, polypeptide or peptide, the aminoacids are arranged in such a manner that they spatially come close together and form the epitope. Continuous peptide fragments of interest can identified by immunization experiments as described above.

[00233] Further preferred RANKL peptides suitable for use for the present invention can be identified by using existing or future monoclonal or polyclonal antibodies, the procedures hereto are know to those skilled in the art.

[00234] Further RANKL peptides suitable for use for the present invention may be identified by screening phage display peptide libraries with antibodies specific for RANKL, a method well known to a person trained in the art.

[00235] In a further preferred embodiment of the invention, the antigen or antigenic determinant is isolated RANKL of any animal as well as any antigenic fragments of RANKL of any animal. Those skilled in the art know how to produce fragments and peptides from those isolated RANKL protein or fragments.

[00236] It will be understood by one of ordinary skill in the relevant arts that other suitable modifications and adaptations to the methods and applications described herein are readily apparent and may be made without departing from the scope of the invention or any embodiment thereof. Having now described the present invention in detail, the same will be more clearly understood by reference to the following examples, which are included herewith for purposes of illustration only and are not intended to be limiting of the invention.

- 83 -

EXAMPLE 1

Construction and expression of mutant Q β coat proteins, and purification of mutant Q β coat protein VLPs or Capsids.

Plasmid construction and cloning of mutant coat proteins

Construction of pQ β -240:

[00237] The plasmid pQ β 10 (Kozlovskaya, TM, *et al.*, *Gene* 137:133-137) was used as an initial plasmid for the construction of pQ β -240. The mutation Lys13 \rightarrow Arg was created by inverse PCR. The inverse primers were designed in inverted tail-to-tail directions:

5'-GGTAACATCGGTTCGAGATGGAAAACAACTCTGGTCC-3'

and

5'-GGACCAGAGTTTGTTCCTTCGACCGATGTTACC-3'.

The products of the first PCR were used as templates for the second PCR reaction, in which an upstream primer

5'-AGCTCGCCCGGGGATCCTCTAG-3'

and a downstream primer

5'-CGATGCATTTTCATCCTTAGTTATCAATACGCTGGGTTTCAG-

3'

were used. The product of the second PCR was digested with *Xba*I and *Mph*1103I and cloned into the pQ β 10 expression vector, which was cleaved by the same restriction enzymes. The PCR reactions were performed with PCR kit reagents and according to producer protocol (MBI Fermentas, Vilnius, Lithuania).

[00238] Sequencing using the direct label incorporation method verified the desired mutations. *E.coli* cells harbouring pQ β -240 supported efficient synthesis of 14-kD protein co migrating upon SDS-PAGE with control Q β coat protein isolated from Q β phage particles.

- 84 -

Resulting amino acid sequence: (SEQ ID NO: 23)

AKLETVTLGNI GRD GKQTLVLNPRGVNPTNGVASLSQAGAVP
ALEKRVTVSVSQPSRNRKNYKVQVKIQNPTACTANGSCDPSVTRQ
KYADVTFSTQYSTDEERAFVRTELAALLASPLLIDAIDQLNPAY

Construction of pQ β -243:

[00239] The plasmid pQ β 10 was used as an initial plasmid for the construction of pQ β -243. The mutation Asn10 \rightarrow Lys was created by inverse PCR. The inverse primers were designed in inverted tail-to-tail directions:

5'-GGCAAATTAGAGACTGTTACTTTAGGTAAGATCGG-3'

and

5'-CCGATCTTACCTAAAGTAACAGTCTCTAATTTTGCC-3'.

The products of the first PCR were used as templates for the second PCR reaction, in which an upstream primer

5'-AGCTCGCCCGGGGATCCTCTAG-3'

and a downstream primer

5'-CGATGCATTTTCATCCTTAGTTATCAATACGCTGGGTTCAG-3'

were used. The product of the second PCR was digested with *Xba*I and *Mph*1103I and cloned into the pQ β 10 expression vector, which was cleaved by the same restriction enzymes. The PCR reactions were performed with PCR kit reagents and according to producer protocol (MBI Fermentas, Vilnius, Lithuania).

[00240] Sequencing using the direct label incorporation method verified the desired mutations. *E.coli* cells harbouring pQ β -243 supported efficient synthesis of 14-kD protein co migrating upon SDS-PAGE with control Q β coat protein isolated from Q β phage particles.

Resulting amino acid sequence: (SEQ ID NO: 24)

AKLETVTLGKIGKD GKQTLVLNPRGVNPTNGVASLSQAGAVP
ALEKRVTVSVSQPSRNRKNYKVQVKIQNPTACTANGSCDPSVTRQ

- 85 -

KYADVTFSTQYSTDEERAFVRTELAALLASPLLIDAIDQLNPAY

Construction of pQ β -250:

[00241] The plasmid pQ β -240 was used as an initial plasmid for the construction of pQ β -250. The mutation Lys2→Arg was created by site-directed mutagenesis. An upstream primer

5'-GGCCATGGCACGACTCGAGACTGTTACTTTAGG-3'

and a downstream primer

5'-GATTTAGGTGACACTATAG-3'

were used for the synthesis of the mutant PCR-fragment, which was introduced into the pQ β -185 expression vector at the unique restriction sites *NcoI* and *HindIII*. The PCR reactions were performed with PCR kit reagents and according to producer protocol (MBI Fermentas, Vilnius, Lithuania).

[00242] Sequencing using the direct label incorporation method verified the desired mutations. *E.coli* cells harbouring pQ β -250 supported efficient synthesis of 14-kD protein co migrating upon PAGE with control Q β coat protein isolated from Q β phage particles.

Resulting amino acid sequence: (SEQ ID NO: 25)

ARLETVTGLGNIGRDGKQTLVLNPRGVNPTNGVASLSQAGAVP

ALEKRVTVSVSQPSRNRKNYKVQVKIQNPACTANGSCDPSVTRQ

KYADVTFSTQYSTDEERAFVRTELAALLASPLLIDAIDQLNPAY

Construction of pQ β -251:

[00243] The plasmid pQ β 10 was used as an initial plasmid for the construction of pQ β -251. The mutation Lys16→Arg was created by inverse PCR. The inverse primers were designed in inverted tail-to-tail directions:

5'-GATGGACGTCAAACCTCTGGTCCTCAATCCGCGTGGGG -3'

and

5'-CCCCACGCGGATTGAGGACCAGAGTTTGACGTCCATC -3'.

- 86 -

The products of the first PCR were used as templates for the second PCR reaction, in which an upstream primer

5'-AGCTCGCCCGGGGATCCTCTAG-3'

and a downstream primer

5'-CGATGCATTTTCATCCTTAGTTATCAATACGCTGGGTTCAG-

3'

were used. The product of the second PCR was digested with *XbaI* and *MphI103I* and cloned into the pQ β 10 expression vector, which was cleaved by the same restriction enzymes. The PCR reactions were performed with PCR kit reagents and according to producer protocol (MBI Fermentas, Vilnius, Lithuania).

[00244] Sequencing using the direct label incorporation method verified the desired mutations. *E.coli* cells harbouring pQ β -251 supported efficient synthesis of 14-kD protein co migrating upon SDS-PAGE with control Q β coat protein isolated from Q β phage particles. The resulting amino acid sequence encoded by this construct is shown in SEQ. ID NO: 26.

Construction of pQ β -259:

[00245] The plasmid pQ β -251 was used as an initial plasmid for the construction of pQ β -259. The mutation Lys2 \rightarrow Arg was created by site-directed mutagenesis. An upstream primer

5'-GGCCATGGCACGACTCGAGACTGTTACTTTAGG-3'

and a downstream primer

5'-GATTTAGGTGACACTATAG-3'

were used for the synthesis of the mutant PCR-fragment, which was introduced into the pQ β -185 expression vector at the unique restriction sites *NcoI* and *HindIII*. The PCR reactions were performed with PCR kit reagents and according to producer protocol (MBI Fermentas, Vilnius, Lithuania).

- 87 -

[00246] Sequencing using the direct label incorporation method verified the desired mutations. *E.coli* cells harbouring pQ β -259 supported efficient synthesis of 14-kD protein co migrating upon SDS-PAGE with control Q β coat protein isolated from Q β phage particles.

Resulting amino acid sequence: (SEQ ID NO: 27)

AKLETVTTLGNIGKDGKQTLVLNPRGVNPTNGVASLSQAGAVP

ALEKRVTVSVSQPSRNRKNYKVQVKIQNPTACTANGSCDPSVTRQ

KYADVTFSTQYSTDEERAFVRTELAALLASPLLIDAIDQLNPAY

General procedures for Expression and purification of Q β and Q β mutants

Expression

[00247] *E.coli* JM109 was transformed with Q β coat protein expression plasmids. 5 ml of LB liquid medium containing 20 μ g/ml ampicillin were inoculated with clones transformed with with Q β coat protein expression plasmids. The inoculated culture was incubated at 37 °C for 16-24 h without shaking. The prepared inoculum was subsequently diluted 1:100 in 100-300 ml of fresh LB medium, containing 20 μ g/ml ampicillin. and incubated at 37°C overnight without shaking. The resulting second inoculum was diluted 1:50 in M9 medium containing 1 % Casamino acids and 0.2 % glucose in flasks, and incubated at 37 °C overnight under shaking.

Purification

[00248] Solutions and buffers for the purification procedure:

1. Lysis buffer LB

50mM Tris-HCl pH8,0 with 5mM EDTA , 0,1%

tritonX100 and freshly prepared PMSF at a concentration of 5micrograms per

ml. Without lysozyme and DNase.

- 88 -

2. SAS
Saturated ammonium sulphate in water
3. Buffer NET.
20 mM Tris-HCl, pH 7.8 with 5mM EDTA and
150 mM NaCl.
4. PEG
40% (w/v) polyethylenglycol 6000 in NET

Disruption and lysis

[00249] Frozen cells were resuspended in LB at 2 ml/g cells. The mixture was sonicated with 22 kHz five times for 15 seconds, with intervals of 1 min to cool the solution on ice. The lysate was then centrifuged at 14 000 rpm, for 1 h using a Janekki K 60 rotor. The centrifugation steps described below were all performed using the same rotor, except otherwise stated. The supernatant was stored at 4° C, while cell debris were washed twice with LB. After centrifugation, the supernatants of the lysate and wash fractions were pooled.

Fractionation

[00250] A saturated ammonium sulphate solution was added dropwise under stirring to the above pooled lysate. The volume of the SAS was adjusted to be one fifth of total volume, to obtain 20% of saturation. The solution was left standing overnight, and was centrifuged the next day at 14 000 rpm, for 20 min. The pellet was washed with a small amount of 20% ammonium sulphate, and centrifuged again. The obtained supernatants were pooled, and SAS was added dropwise to obtain 40% of saturation. The solution was left standing overnight, and was centrifuged the next day at 14 000 rpm, for 20 min. The obtained pellet was solubilised in NET buffer.

Chromatography

[00251] The capsid or VLP protein resolubilized in NET buffer was loaded on a Sepharose CL- 4B column. Three peaks eluted during chromatography. The first one mainly contained membranes and membrane fragments, and was not collected. Capsids were contained in the second peak, while the third one contained other *E.coli* proteins.

[00252] The peak fractions were pooled, and the NaCl concentration was adjusted to a final concentration of 0.65 M. A volume of PEG solution corresponding to one half of the pooled peak fraction was added dropwise under stirring. The solution was left to stand overnight without stirring. The capsid protein was sedimented by centrifugation at 14 000 rpm for 20 min. It was then solubilized in a minimal volume of NET and loaded again on the Sepharose CL- 4B column. The peak fractions were pooled, and precipitated with ammonium sulphate at 60% of saturation (w/v). After centrifugation and resolubilization in NET buffer, capsid protein was loaded on a Sepharose CL- 6B column for rechromatography.

Dialysis and drying

[00253] The peak fractions obtained above were pooled and extensively dialysed against sterile water, and lyophilized for storage.

Expression and purification Q β -240

[00254] Cells (*E. coli* JM 109, transformed with the plasmid pQ β -240) were resuspended in LB, sonicated five times for 15 seconds (water ice jacket) and centrifuged at 13000 rpm for one hour. The supernatant was stored at 4°C until further processing, while the debris were washed 2 times with 9 ml of LB, and finally with 9 ml of 0,7 M urea in LB. All supernatants were pooled, and loaded on the Sepharose CL-4B column. The pooled peak fractions were precipitated with ammonium sulphate and centrifuged. The resolubilized

- 90 -

protein was then purified further on a Sepharose 2B column and finally on a Sepharose 6B column. The capsid peak was finally extensively dialyzed against water and lyophilized as described above. The assembly of the coat protein into a capsid was confirmed by electron microscopy.

Expression and purification Q β -243

[00255] Cells (*E. coli* RR1) were resuspended in LB and processed as described in the general procedure. The protein was purified by two successive gel filtration steps on the sepharose CL-4B column and finally on a sepharose CL-2B column. Peak fractions were pooled and lyophilized as described above. The assembly of the coat protein into a capsid was confirmed by electron microscopy.

Expression and purification of Q β -250

[00256] Cells (*E. coli* JM 109, transformed with pQ β -250) were resuspended in LB and processed as described above. The protein was purified by gel filtration on a Sepharose CL-4B and finally on a Sepharose CL-2B column, and lyophilized as described above. The assembly of the coat protein into a capsid was confirmed by electron microscopy.

Expression and purification of Q β -259

[00257] Cells (*E. coli* JM 109, transformed with pQ β -259) were resuspended in LB and sonicated. The debris were washed once with 10 ml of LB and a second time with 10 ml of 0,7 M urea in LB. The protein was purified by two gel-filtration chromatography steps, on a Sepharose CL-4 B column. The protein was dialyzed and lyophilized, as described above. The assembly of the coat protein into a capsid was confirmed by electron microscopy.

EXAMPLE 2

Insertion of a peptide containing a Lysine residue into the c/e1 epitope of HBcAg(1-149).

[00258] The c/e1 epitope (residues 72 to 88) of HBcAg is located in the tip region on the surface of the Hepatitis B virus capsid (HBcAg). A part of this region (Proline 79 and Alanine 80) was genetically replaced by the peptide Gly-Gly-Lys-Gly-Gly (HBcAg-Lys construct). The introduced Lysine residue contains a reactive amino group in its side chain that can be used for intermolecular chemical crosslinking of HBcAg particles with any antigen containing a free cysteine group.

[00259] HBcAg-Lys DNA, having the amino acid sequence shown in SEQ ID NO:78, was generated by PCRs: The two fragments encoding HBcAg fragments (amino acid residues 1 to 78 and 81 to 149) were amplified separately by PCR. The primers used for these PCRs also introduced a DNA sequence encoding the Gly-Gly-Lys-Gly-Gly peptide. The HBcAg (1 to 78) fragment was amplified from pEco63 using primers EcoRIHBcAg(s) and Lys-HBcAg(as). The HBcAg (81 to 149) fragment was amplified from pEco63 using primers Lys-HBcAg(s) and HBcAg(1-149)Hind(as). Primers Lys-HBcAg(as) and Lys-HBcAg(s) introduced complementary DNA sequences at the ends of the two PCR products allowing fusion of the two PCR products in a subsequent assembly PCR. The assembled fragments were amplified by PCR using primers EcoRIHBcAg(s) and HbcAg(1-149)Hind(as).

[00260] For the PCRs, 100 pmol of each oligo and 50 ng of the template DNAs were used in the 50 ml reaction mixtures with 2 units of Pwo polymerase, 0.1 mM dNTPs and 2 mM MgSO₄. For both reactions, temperature cycling was carried out as follows: 94°C for 2 minutes; 30 cycles of 94°C (1 minute), 50°C (1 minute), 72°C (2 minutes).

- 92 -

Primer sequences:

EcoRIHBcAg(s):

(5'-CCGGAATTCATGGACATTGACCCTTATAAAG-3');

Lys-HBcAg(as):

(5'-

CCTAGAGCCACCTTTGCCACCATCTTCTAAATTAGTACCCACCCAG
GTAGC-3');

Lys-HBcAg(s):

(5'-

GAAGATGGTGGCAAAGGTGGCTCTAGGGACCTAGTAGTCAGTTAT
GTC -3');

HBcAg(1-149)Hind(as):

(5'-CGCGTCCCAAGCTTCTAAACAACAGTAGTCTCCGGAAG-3').

[00261] For fusion of the two PCR fragments by PCR 100 pmol of primers EcoRIHBcAg(s) and HBcAg(1-149)Hind(as) were used with 100 ng of the two purified PCR fragments in a 50 ml reaction mixture containing 2 units of Pwo polymerase, 0.1 mM dNTPs and 2 mM MgSO₄. PCR cycling conditions were: 94°C for 2 minutes; 30 cycles of 94°C (1 minute), 50°C (1 minute), 72°C (2 minutes). The assembled PCR product was analyzed by agarose gel electrophoresis, purified and digested for 19 hours in an appropriate buffer with EcoRI and HindIII restriction enzymes. The digested DNA fragment was ligated into EcoRI/HindIII-digested pKK vector to generate pKK-HBcAg-Lys expression vector. Insertion of the PCR product into the vector was analyzed by EcoRI/HindIII restriction analysis and DNA sequencing of the insert.

EXAMPLE 3

Expression and purification of HBcAg-Lys.

- [00262] *E. coli* strains K802 or JM109 were transformed with pKK-HBcAg-Lys. 1 ml of an overnight culture of bacteria was used to inoculate 100 ml of LB medium containing 100 µg/ml ampicillin. This culture was grown for 4 hours at 37°C until an OD at 600 nm of approximately 0.8 was reached. Induction of the synthesis of HBcAg-Lys was performed by addition of IPTG to a final concentration of 1 mM. After induction, bacteria were further shaken at 37°C for 4 hours. Bacteria were harvested by centrifugation at 5000 x g for 15 minutes. The pellet was frozen at -80°C. The pellet was thawed and resuspended in bacteria lysis buffer (10 mM Na₂HPO₄, pH 7.0, 30 mM NaCl, 0.25% Tween-20, 10 mM EDTA) supplemented with 200 µg/ml lysozyme and 10 µl of Benzonase (Merck). Cells were incubated for 30 minutes at room temperature and disrupted by sonication. *E. coli* cells harboring pKK-HBcAg-Lys expression plasmid or a control plasmid were used for induction of HBcAg-Lys expression with IPTG. Prior to the addition of IPTG, a sample was removed from the bacteria culture carrying the pKK-HBcAg-Lys plasmid and from a culture carrying the control plasmid. Four hours after addition of IPTG, samples were again removed from the culture containing pKK-HBcAg-Lys and from the control culture. Protein expression was monitored by SDS-PAGE followed by Coomassie staining.
- [00263] The lysate was then centrifuged for 30 minutes at 12,000 x g in order to remove insoluble cell debris. The supernatant and the pellet were analyzed by Western blotting using a monoclonal antibody against HBcAg (YVS1841, purchased from Accurate Chemical and Scientific Corp., Westbury, NY, USA), indicating that a significant amount of HBcAg-Lys protein was soluble. Briefly, lysates from *E. coli* cells expressing HBcAg-Lys and from control cells were centrifuged at 14,000 x g for 30 minutes. Supernatant (= soluble fraction) and pellet (= insoluble fraction) were separated and diluted with SDS sample buffer to equal volumes. Samples were analyzed by SDS-PAGE

followed by Western blotting with anti-HBcAg monoclonal antibody YVS 1841.

[00264] The cleared cell lysate was used for step-gradient centrifugation using a sucrose step gradient consisting of a 4 ml 65% sucrose solution overlaid with 3 ml 15% sucrose solution followed by 4 ml of bacterial lysate. The sample was centrifuged for 3 hrs with 100,000 x g at 4°C. After centrifugation, 1 ml fractions from the top of the gradient were collected and analyzed by SDS-PAGE followed by Coomassie staining. The HBcAg-Lys protein was detected by Coomassie staining.

[00265] The HBcAg-Lys protein was enriched at the interface between 15 and 65% sucrose indicating that it had formed a capsid particle. Most of the bacterial proteins remained in the sucrose-free upper layer of the gradient, therefore step-gradient centrifugation of the HBcAg-Lys particles led both to enrichment and to a partial purification of the particles.

[00266] Expression and purification of HBcAg-Lys in large scale was performed as follows. An overnight culture was prepared by inoculating a single colony in 100 ml LB, 100 µg/ml Ampicillin and growing the culture overnight at 37°C. 25 ml of the preculture were diluted in 800 ml LB Ampicillin medium the next day, and the culture grown to an optical density OD⁶⁰⁰ of 0.6-0.8. The culture was then induced with 1 mM IPTG, and left to grow for another 4 hours. The cells were harvested and lysed essentially as described above.

[00267] HBcAg-Lys was then purified by first precipitating the protein with ammonium sulphate (30% saturation) from the cleared cell lysate, then loading the resolubilized pellet on a gel filtration column (Sephacryl S-400, Pharmacia). The pooled fractions were precipitated again with ammonium sulphate, the pellet resolubilized and loaded a second time on the same gel filtration column. The fractions were finally pooled and concentrated, and the concentration assessed using a Bradford test (BioRad).

- 95 -

EXAMPLE 4

Construction of a HBcAg devoid of free cysteine residues and containing an inserted lysine residue.

[00268] A Hepatitis core Antigen (HBcAg), referred to herein as HBcAg-lys-2cys-Mut, devoid of cysteine residues at positions corresponding to 48 and 107 in SEQ ID NO:77 and containing an inserted lysine residue was constructed using the following methods.

[00269] The two mutations were introduced by first separately amplifying three fragments of the HBcAg-Lys gene prepared as described above in Example 2 with the following PCR primer combinations. PCR methods and conventional cloning techniques were used to prepare the HBcAg-lys-2cys-Mut gene.

[00270] In brief, the following primers were used to prepare fragment 1:

Primer 1: EcoRIHBcAg(s)

CCGGAATTCATGGACATTGACCCTTATAAAG

Primer 2: 48as

GTGCAGTATGGTGAGGTGAGGAATGCTCAGGAGACTC

[00271] The following primers were used to prepare fragment 2:

Primer 3: 48s

GSGTCTCCTGAGCATTCTCACCTCACCATACTGCAC

Primer 4: 107as

CTTCCAAAAGTGAGGGAAGAAATGTGAAACCAC

[00272] The following primers were used to prepare fragment 3:

Primer 5: HBcAg149hind-as

CGCGTCCCAAGCTTCTAAACAACAGTAGTCTCCGGAAGCGTTGATA

G

- 96 -

Primer 6: 107s

GTGGTTTCACATTCTTCCCTCACTTTTGGAAG

[00273] Fragments 1 and 2 were then combined with PCR primers EcoRIHBcAg(s) and 107as to give fragment 4. Fragment 4 and fragment 3 were then combined with primers EcoRIHBcAg(s) and HBcAg149hind-as to produce the full length gene. The full length gene was then digested with the EcoRI (GAATTC) and HindIII (AAGCTT) enzymes and cloned into the pKK vector (Pharmacia) cut at the same restriction sites. Expression and purification of HBcAg-lys-2cys-Mut were performed as set out in Example 3.

EXAMPLE 5

Construction of HBcAg1-185-Lys.

[00274] Hepatitis core Antigen (HBcAg) 1-185 was modified as described in Example 2. A part of the c/e1 epitope (residues 72 to 88) region (Proline 79 and Alanine 80) was genetically replaced by the peptide Gly-Gly-Lys-Gly-Gly (HBcAg1-185-Lys construct). The introduced Lysine residue contains a reactive amino group in its side chain that can be used for intermolecular chemical crosslinking of HBcAg particles with any antigen containing a free cysteine group. PCR methods and conventional cloning techniques were used to prepare the HBcAg1-185-Lys gene.

[00275] The Gly-Gly-Lys-Gly-Gly sequence was inserted by amplifying two separate fragments of the HBcAg gene from pEco63, as described above in Example 2 and subsequently fusing the two fragments by PCR to assemble the full length gene. The following PCR primer combinations were used:

[00276] fragment 1:

Primer 1: EcoRIHBcAg(s) (see Example 2)

Primer 2: Lys-HBcAg(as) (see Example 2)

fragment 2:

Primer 3: Lys-HBcAg(s) (see Example 2)

- 97 -

Primer 4: HBcAgwtHindIII

CGCGTCCCAAGCTTCTAACATTGAGATTCCCGAGATTG

Assembly:

Primer 1: EcoRIHBcAg(s) (see example 2)

Primer 2: HBcAgwtHindIII

[00277] The assembled full length gene was then digested with the EcoRI (GAATTC) and HindIII (AAGCTT) enzymes and cloned into the pKK vector (Pharmacia) cut at the same restriction sites.

EXAMPLE 6

Fusion of a peptide epitope in the MIR region of HbcAg.

[00278] The residues 79 and 80 of HBcAg1-185 were substituted with the epitope CεH3 of sequence VNLTWSRASG . The CεH3 sequence stems from the sequence of the third constant domain of the heavy chain of human IgE. The epitope was inserted in the HBcAg1-185 sequence using an assembly PCR method. In the first PCR step, the HBcAg1-185 gene originating from ATCC clone pEco63 and amplified with primers HBcAg-wt EcoRI fwd and HBcAg-wt Hind III rev was used as template in two separate reactions to amplify two fragments containing sequence elements coding for the CεH3 sequence. These two fragments were then assembled in a second PCR step, in an assembly PCR reaction.

[00279] Primer combinations in the first PCR step: CεH3fwd with HBcAg-wt Hind III rev, and HBcAg-wt EcoRI fwd with CεH3rev. In the assembly PCR reaction, the two fragments isolated in the first PCR step were first assembled during 3 PCR cycles without outer primers, which were added afterwards to the reaction mixture for the next 25 cycles. Outer primers: HBcAg-wt EcoRI fwd and HBcAg-wt Hind III rev.

- 98 -

[00280] The PCR product was cloned in the pKK223.3 using the EcoRI and HindIII sites, for expression in *E. coli* (see Example 2). The chimeric VLP was expressed in *E. coli* and purified as described in Example 2. The elution volume at which the HBcAg1-185- CεH3 eluted from the gel filtration showed assembly of the fusion proteins to a chimeric VLP.

Primer sequences:

CεH3fwd:

5' GTT AAC TTG ACC TGG TCT CGT GCT TCT GGT GCA TCC AGG GAT CTA GTA GTC 3'
V N L T W S R A S G A80 S R D L V V86

CεH3rev:

5' ACC AGA AGC ACG AGA CCA GGT CAA GTT AAC ATC TTC CAA ATT ATT ACC CAC 3'
D78 E L N N G V72

HBcAg-wt EcoRI fwd:

5' CCGgaattcATGGACATTGACCCTTATAAAG

HBcAg-wt Hind III rev:

5' CGCGTCCCcaagcttCTAACATTGAGATTCCCGAGATTG

EXAMPLE 7

Fusion of a RANKL peptide epitope in the MIR region of HbcAg.

[00281] The residues 79 and 80 of HBcAg1-185 are substituted with the RANKL peptide epitope of sequence: SIKIPSSH. Two overlapping primers are designed using the same strategy described in Example 6, and the fusion protein constructed by assembly PCR. The PCR product is cloned in the pKK223.3 vector, and expressed in *E. coli* K802. The chimeric VLPs are expressed and purified as described in Example 3.

- 99 -

EXAMPLE 8

Fusion of a RANKL peptide epitope to the C-terminus of the Q β A1 protein truncated at position 19 of the CP extension.

[00282] A primer annealing to the 5' end of the Q β A1 gene and a primer annealing to the 3' end of the A1 gene and comprising additionally a sequence element coding for the RANKL peptide epitope of sequence: SIKIPSSH, are used in a PCR reaction with pQ β 10 as template. The PCR product is cloned in pQ β 10 (Kozlovska T.M. *et al.*, *Gene* 137: 133-37 (1993)), and the chimeric VLP expressed and purified as described in Example 1.

EXAMPLE 9

Insertion of a RANKL peptide epitope between positions 2 and 3 of fr coat protein.

[00283] Complementary primers coding for the sequence of the RANKL peptide epitope of sequence: SIKIPSSH, and containing *Bsp*119I compatible ends and additional nucleotides enabling in frame insertion, are inserted in the *Bsp*119I site of the pFrd8 vector (Pushko, P. *et al.*, *Prot. Eng.* 6: 883-91 (1993)) by standard molecular biology techniques. Alternatively, the overhangs of the pFrd8 vector are filled in with Klenow after digestion with *Bsp*119I, and oligonucleotides coding for the sequence of the RANKL protein, RANKL fragment or RANKL peptide and additional nucleotides for in frame cloning are ligated in pFrd8 after the Klenow treatment. Clones with the insert in the right orientation are analysed by sequencing. Expression and purification of the chimeric fusion protein in *E. coli* JM109 or *E. coli* K802 is performed as described in Pushko, P. *et al.*, *Prot. Eng.* 6:883-91 (1993), but for the chromatography steps which are performed using a Sepharose CL-4B or Sephacryl S-400 (Pharmacia). The cell lysate is precipitated with ammonium sulphate, and purified by two successive gel filtration purification steps, similarly to the procedure described for Q β in Example 1.

- 100 -

EXAMPLE 10

Insertion of a RANKL peptide epitope between positions 67 and 68 of Ty1 protein p1 in the vector pOGS8111.

[00284] Two complementary oligonucleotides coding for the RANKL peptide epitope of sequence: SIKIPSSH, with ends compatible with the NheI site of pOGS8111 are synthesized. Additional nucleotides are added to allow for in frame insertion of a sequence coding for the RANKL epitope according to the description of EP0677111. The amino acids AS and SS flanking the inserted epitope are encoded by the altered NheI sites resulting from the insertion of the oligonucleotide in the TyA(d) gene of pOGS8111.

[00285] pOGS8111 is transformed into *S. cerevisiae* strain MC2, for expression of the chimeric Ty VLP as described in EP0677111 and references therein. The chimeric Ty VLP is purified by sucrose gradient ultracentrifugation as described in EP0677111.

EXAMPLE 11

Insertion of a RANKL peptide epitope in to the major capsid protein L1 of papillomavirus type 1 (BPV-1).

[00286] A sequence coding for the RANKL peptide epitope of sequence SIKIPSSH is substituted to the sequence coding for amino acids 130-136 of the BPV-1 L1 gene cloned in the pFastBac1 (GIBCO/BRL) vector as described (Chackerian, B. *et al.*, *Proc. Natl. Acad. USA* 96: 2373-2378 (1999)). The sequence of the construct is verified by nucleotide sequence analysis. Recombinant baculovirus is generated using the GIBCO/BRL baculovirus system as described by the manufacturer. The chimeric VLPs are purified from baculovirus infected Sf9 cells as described by Kimbauer, R. *et al.*, *Proc. Natl. Acad. Sci.* 89:12180-84 (1992) and Greenstone, H.L., *et al.*, *Proc. Natl. Acad. Sci.* 95:1800-05 (1998).

- 101 -

EXAMPLE 12

Introduction of an N-terminal cys-containing linker, expression and purification of RANKL

A fragment of RANKL was recombinantly expressed with an N-terminal linker containing one cysteine for coupling to VLP.

Construction of expression plasmid

The C-terminal coding region of the RANKL gene was amplified by PCR with oligos RANKL-UP and RANKL-DOWN (Oligos: RANKL-UP: 5'-CTGCCAGGGGCCCGGGTGCGGCGGTGGCCATCATCACCACCAT-CACCAGCGCTTCTCAGGAG-3'; RANKL-DOWN : 5'-CCGCTCGAGT-TAGTCTATGTCCTGAACTTTGAAAG-3'). RANKL-UP had an internal *Apal* site and RANKL-DOWN had an internal *XhoI* site. The PCR product was digested with *Apal* and *XhoI* and ligated into pGEX-6p1 (Amersham Pharmacia). The resulting plasmid was named pGEX-RANKL. All steps were performed by standard molecular biology protocols and the sequence was verified. The plasmid pGEX-RANKL codes for a fusion protein of a glutathione S-transferase-PreScission cleavage site-cysteine-containing amino acid linker-RANKL (GST-PS-C-RANKL). The cysteine-containing amino acid linker had the sequence GPGCGGG. The construct also contains a hexahistidine tag between the cysteine containing amino acid linker and the RANKL sequence. Sequences of the resulting cDNA and protein constructs are given as SEQ-ID NO: 94 and SEQ-ID NO: 95.

Expression and Purification of C-RANKL

Competent *Escherichia coli* BL21 (DE3) Gold pLys cells were transformed with the plasmid pGEX-RANKL. Single colonies from ampicillin-containing agar plates were expanded in liquid culture (LB

- 102 -

medium, 100 µg/ml ampicillin) and incubated at 30°C with 220 rpm shaking overnight. One litre of LB (with 100 µg/ml ampicillin) was then inoculated 1:100 v/v with the overnight culture and grown at 24°C to OD₆₀₀=1. Expression was induced with 0.4 mM IPTG. Cells were harvested after 16 h and centrifuged at 5000 rpm. Cell pellet was suspended in lysis buffer (50 mM Tris-HCl, pH 8.0; 25 % sucrose; 1 mM EDTA, 1% NaN₃; 10 mM DTT; 5 mM MgCl₂; 1 mg/ml Lysozyme; 0.4 U/ml DNase) for 30 min. Then 2.5 volumes of buffer A (50 mM Tris-HCl, pH 8.0; 1% Triton X100; 100 mM NaCl; 0.1% NaN₃; 10 mM DTT; 1 mM PMSF) were added and incubated at 37°C for 15 min. The cells were sonicated and pelleted at 9000 rpm for 15 min. The supernatant was immediately used for GST-affinity chromatography.

A GST-Trap FF column of 5 ml (Amersham Pharmacia) was equilibrated in PBS, pH 7.3 (140 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, 1.8 mM KH₂PO₄). The supernatant was loaded on the 5 ml GST-Trap FF column and subsequently the column was rinsed with 5 column volumes of PBS. The protein GST-PS-C-RANKL was eluted with 50 mM Tris-HCl, pH 8.0 containing 10 mM reduced glutathione.

The purified GST-PS-C-RANKL protein was digested using the protease PreScission (Amersham Pharmacia). The digestion was performed at 37°C for 1 hour using a molar ratio of 500/1 of GST-PS-C-RANKL to PreScission.

Furthermore, the reaction of protease digestion was buffer exchanged using a HiPrep 26/10 desalting column (Amersham Pharmacia), the fractions containing the proteins were pooled and immediately used for another step of GST affinity chromatography using the same conditions reported before. Purification of C-RANKL was analysed on a SDS-PAGE gel under reducing conditions, shown in Fig.1. The cleaved C-RANKL is present in the flow-through (unbound fraction) while the uncleaved GST-PS-C-RANKL, the cleaved GST-PS and the PreScission remain bound to the column. C-RANKL protein (SEQ ID NO:96) of the expected size of 22 kDa was obtained in high purity.

EXAMPLE 13

Introduction of a C-terminal cys-containing linker, expression and purification of RANKL

A fragment of the RANKL was recombinantly expressed with a C-terminal linker containing one cysteine for coupling to VLP.

Construction of expression plasmid

The MCS of pET22b(+) (Novagen, Inc.) was changed to GTTTAACTTTAAGAAGGAGATATACATATGGATCCGGCTAGCGCT-CGAGGGTTTAAACGGCGGCCGCATGCACC by replacing the original sequence from the NdeI site to XhoI site with annealed oligos primerMCS-1F and primerMCS-1R (annealing in 15 mM TrisHCl pH 8 buffer). The resulting plasmid was termed pMod00, which had NdeI, BamHI, NheI, XhoI, PmeI and NotI restriction sites in its MCS. The annealed pair of oligos Bamhis6-EK-Nhe-F and Bamhis6-EKNhe-R and the annealed pair of oligo1F-C-glycine-linker and oligo1R-C-glycine-linker were together ligated into BamHI-NotI digested pMod00 plasmid to get pModEC1, which had an N terminal hexahistidine tag, an enterokinase cleavage site and a C-terminal amino acid glycine linker containing one cysteine residue.

A DNA fragment comprising the glutathione S transferase gene with a C-terminal enterokinase cleavage site was amplified by PCR with oligonucleotides GST-UP and GST-EK from plasmid SP-GST-EK-pCEP-Pu (Wuttke, M., *et al.*, *J. Biol. Chem.*, 276: 36839-36848), digested with *NheI* and *BamHI* and cloned into the pModEC1 vector.

The resulting plasmid pMod-GST-EK-C1 comprises the gene coding for glutathione S transferase fused to an enterokinase cleavage site and a C-terminal cys-containing linker. The C-terminal coding sequence of RANKL

- 104 -

was then amplified by PCR with oligonucleotides mRANKL-1 and mRANKL-2, digested with *NheI* and *XhoI*, and cloned into plasmid pMod-GST-EK-C1. The resulting plasmid pMod-GST-EK-mRANKL-C1 encodes a fusion protein consisting of glutathione S-transferase, an enterokinase cleavage site, the RANKL fragment, and a cysteine-containing amino acid linker (GST-EK-RANKL-C). Sequences of the resulting cDNA and protein constructs are given as SEQ ID NO:97 and SEQ ID NO: 98.

Sequence of oligonucleotides:

GST-UP: 5'-ATATATGGATCCTATACTAGGTTATTGGAAAAT-3';

GST-EK: 5'-ATATATGCTAGCTTATCGTCATCGTCG-3';

mRANKL-1: 5'-ATATATGCTAGCAAAGCCTGAGGCCAGCC-
ATTG3';

mRANKL-2: 5'-ATATATCTCGAGGTCTATGTCCTGAACTTT-
GAAAG3';

Expression and Purification of RANKL-C

Competent *Escherichia coli* BL21 (DE3) Gold pLys cells were transformed with the plasmid pMod-GST-EK-mRANKL-C1. Single colonies from ampicillin-containing agar plates were expanded in 100 ml liquid culture (LB medium, 200µg/ml ampicillin) and incubated at 30°C with 220 rpm shaking overnight. One litre of medium (SB with 150mM MOPS, pH 7.0, 200µg/ml Amp) was then inoculated 1:100 v/v with the overnight culture and grown at 30°C with 125 rpm shaking to OD₆₀₀=2.5. Cultures were then shifted to 18°C and protein expression was induced after 30 min by addition of 0.1 mM IPTG. Bacteria were harvested after overnight culture at 18°C by centrifugation (SLA-3000, 15 min, 4°C, 6000 rpm), resuspended in 40 ml lysis buffer (10 mM Na₂HPO₄, 30 mM NaCl, 10 mM EDTA and 0.25% Tween-20) and incubated for 30 min on ice with 0.8 mg/ml lysozyme. Bacteria were then

- 105 -

lysed by sonication and incubated for 30 min at RT with 0.2 M MgCl_2 and 8 μl Benzonase. The lysate was cleared from insoluble material by centrifugation (SS-34, 30 min, 4°C, 20000 rpm) and used immediately for glutathione sepharose affinity chromatography.

A GST-Trap FF column of 5 ml (Amersham Pharmacia) was therefore equilibrated with lysis buffer (10 mM Na_2HPO_4 , 30 mM NaCl, 10 mM EDTA and 0.25% Tween-20) and loaded with the cleared lysate at a constant flow rate of 0.5 ml/min. The column was then washed three times with 5 column volumes of lysis buffer and the protein GST-EK-RANKL-C was eluted in 9 fractions of 1ml elution buffer (50 mM Tris-HCl, pH 8.0, 10 mM reduced glutathione) each. The purification, shown in Fig. 2A, resulted in GST-EK-RANKL-C fusion protein of about 45 kDa with small proportion of GST-EK.

Elution fractions were pooled and the purified GST-EK-RANKL-C protein was digested using EnterokinaseMaxTM (Invitrogen). The digestion was performed at 4°C for 16 hours using 10 units EnterokinaseMax per mg of purified GST-EK-RANKL-C. Fig. 2B shows that the cleavage reaction lead to RANKL-C with an apparent MW of about 16 kDa.

After cleavage the protein solution was dialysed against PBS pH 7.2 (140 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4 , 1.8 mM KH_2PO_4) and the glutathione S transferase was removed by a second glutathione sepharose affinity chromatography. Therefore a GST-Trap FF column of 5 ml was equilibrated with PBS pH 7.2, and the protein solution was loaded on the column at a constant flow rate of 0.5 ml/min. Protein fractions were analysed before and after glutathione sepharose chromatography on SDS gels. As shown in Fig. 2C the cleaved RANKL-C protein (SEQ ID NO:99) was contained in the flow through in high purity, while the GST-EK protein remained bound to the column.

- 106 -

EXAMPLE 14

Coupling of C-RANKL to Q β capsid protein

A solution of 1.48 ml of 6 mg/ml Q β capsid protein in 20 mM Hepes, 150 mM NaCl pH 7.2 was reacted for 30 minutes with 14.8 μ l of a SMPH (Pierce) (from a 100 mM stock solution dissolved in DMSO) at 25°C. The reaction solution was subsequently dialyzed twice for 3 hours against 2 l of 20 mM Hepes, 150 mM NaCl, pH 7.0 at 4°C. A solution of 230 μ l of 9.8 mg/ml C-RANKL protein in 20 mM Hepes, 150 mM NaCl pH 7.2 was reacted for 30 min with 1.7 μ l of a 100 mM solution of TCEP (Pierce) at 25°C. For coupling 80 μ l of the derivatized and dialyzed Q β was then mixed with 24 μ l of reduced C-RANKL and 96 μ l of 20 mM Hepes, 150 mM NaCl, pH 7.0 and incubated over night at 25°C.

Coupled products were analysed on 16% SDS-PAGE gels under reducing conditions. Gels were either stained with Coomassie Brilliant Blue or blotted onto nitrocellulose membranes. In the latter case membranes were blocked and incubated either with a polyclonal rabbit anti-Q β antiserum (dilution 1:2000) followed by a horse radish peroxidase-conjugated goat anti-rabbit IgG (dilutions 1:5000), or a monoclonal mouse anti-RANKL antibody (dilution 1:2000) followed by a horse raddish peroxidase-conjugated goat anti mouse antibody (dilution 1:5000). Blots were then developed with the ECLTM Western Blotting Detection Reagents (Amersham Pharmacia). The results are shown in Fig. 3A and Fig. 3B. Coupled products could be detected in the Coomassie-stained gels (Fig. 3A) and by both anti-Q β antiserum and the anti-RANKL antibody (Fig. 3B), clearly demonstrating the covalent coupling of C-RANKL to Q β capsid protein.

- 107 -

EXAMPLE 15

Immunization of mice with C-RANKL coupled to Q β capsid protein

A total of 8 female Balb/c mice were vaccinated with C-RANKL coupled to Q β capsid protein. 25 μ g of total protein of each sample was diluted in PBS to 200 μ l and injected subcutaneously (100 μ l on two ventral sides) on day 0, day 16 and day 64. Four mice received the vaccine without addition of adjuvants while the other four received the vaccine with the addition of alum. Mice were bled retroorbitally on day 0, 16, 23, 64, and 78 and their serum was analyzed using a RANKL-specific ELISA.

EXAMPLE 16

Detection of RANKL-specific antibodies in an ELISA

ELISA plates were coated with C-RANKL at a concentration of 10 μ g/ml. The plates were blocked and then incubated with serially diluted mouse sera from day 16, 23, 64, and 78. Bound antibodies were detected with enzymatically labeled anti-mouse IgG antibody. As a control, preimmune serum of the same mice was also tested. Fig. 4 shows the average titers of RANKL-specific antibodies that could be detected in sera mice which had been immunized with Q β -C-RANKL with or without alum. ELISA titers are expressed as serum dilutions which lead to half maximal OD in the ELISA assay. In mice immunized without alum an average titer of 28000 was reached, while the average titer in mice immunized with the addition of alum was 160000. Preimmune sera did not show any reactivity with C-RANKL. This clearly demonstrates that a RANKL-VLP conjugate is able to induce a high antibody titer against a selfprotein.

- 108 -

EXAMPLE 17

Inhibition of RANKL-RANK interaction by sera of mice immunized with Q β -C-RANKL

To test whether the antibodies generated in mice vaccinated with Q β -C-RANKL have neutralizing activity, an in vitro binding assay for RANKL and its cognate receptor RANK was established. ELISA plates were therefore coated with C-RANKL protein at a concentration of 10 μ g/ml and incubated with serial dilutions of a purified RANK-Fc fusion protein or an unrelated Fc-fusion protein as negative control. Binding proteins were detected with a horse raddish peroxidase conjugated anti-Fc antibody. Fig. 5A shows the result of this analysis. The purified RANK-Fc fusion protein was found to bind with a high affinity (half maximal binding at 1-3 nM) to its ligand RANKL, while virtually no binding was observed when the unrelated Fc-fused protein was used.

Sera of mice vaccinated with C-RANKL coupled to Q β were then tested for their ability to inhibit the binding of RANKL to RANK-Fc. ELISA plates were therefore coated with C-RANKL protein at a concentration of 10 μ g/ml, and co-incubated with serial dilutions of mouse sera from day 78 mixed with 1 nM RANK-Fc fusion protein. Binding of the RANK-Fc fusion protein to C-RANKL was detected with horse raddish peroxidase conjugated anti-Fc antibody. Fig. 5B shows that all 8 mice, that received the vaccine, produced antibodies that specifically inhibited binding of RANK-Fc to C-RANKL. On average, half maximal inhibition was achieved when sera dilutions of 1:90 were used. This clearly demonstrates that immunization with an RANKL-VLP conjugate does induce antibodies with high titers that are able to inhibit the interaction of RANKL with its receptor RANK. Thus, inhibition of the RANK-RANKL interaction by way of injecting the specific embodiment of this invention may reverse or prevent bone diseases characterized by increased bone resorption.

EXAMPLE 18

Expression and Purification of Recombinant AP205 VLP

A. Expression of recombinant AP205 VLP

E.coli JM109 was transformed with plasmid pAP283-58. 5 ml of LB liquid medium with 20 µg/ml ampicillin were inoculated with a single colony, and incubated at 37 °C for 16-24 h without shaking.

The prepared inoculum was diluted 1:100 in 100-300 ml of LB medium, containing 20 µg/ml ampicillin and incubated at 37 °C overnight without shaking. The resulting second inoculum was diluted 1:50 in 2TY medium, containing 0.2 % glucose and phosphate for buffering, and incubated at 37 °C overnight on a shaker. Cells were harvested by centrifugation and frozen at -80°C.

B. Purification of recombinant AP205 VLP

Solutions and buffers:

1. Lysis buffer
50mM Tris-HCl pH 8.0 with 5mM EDTA , 0.1% tritonX100 and PMSF at 5 micrograms per ml.
2. SAS
Saturated ammonium sulphate in water
3. Buffer NET.
20 mM Tris-HCl, pH 7.8 with 5mM EDTA and 150 mM NaCl.
4. PEG
40% (w/v) polyethylenglycol 6000 in NET

- 110 -

Lysis:

Frozen cells were resuspended in lysis buffer at 2 ml/g cells. The mixture was sonicated with 22 kHz five times for 15 seconds, with intervals of 1 min to cool the solution on ice. The lysate was then centrifuged for 20 minutes at 12 000 rpm, using a F34-6-38 rotor (Ependorf). The centrifugation steps described below were all performed using the same rotor, except otherwise stated. The supernatant was stored at 4° C, while cell debris were washed twice with lysis buffer. After centrifugation, the supernatants of the lysate and wash fractions were pooled.

Fractionation:

Ammonium-sulphate precipitation can be further used to purify AP205 VLP. In a first step, a concentration of ammonium-sulphate at which AP205 VLP does not precipitate is chosen. The resulting pellet is discarded. In the next step, an ammonium sulphate concentration at which AP205 VLP quantitatively precipitates is selected, and AP205 VLP is isolated from the pellet of this precipitation step by centrifugation (14 000 rpm, for 20 min). The obtained pellet is solubilised in NET buffer.

Chromatography:

The capsid protein from the pooled supernatants was loaded on a Sepharose 4B column (2.8 X 70 cm), and eluted with NET buffer, at 4 ml/hour/fraction. Fractions 28-40 were collected, and precipitated with ammonium sulphate at 60% saturation. The fractions were analyzed by SDS-PAGE and Western Blot with an antiserum specific for AP205 prior to precipitation (FIG. 6A and FIG. 6B). The pellet isolated by centrifugation was resolubilized in NET buffer, and loaded on a Sepharose 2B column (2.3 X 65 cm), eluted at 3 ml/h/fraction. Fractions were analysed by SDS-PAGE, and fractions 44-50 were collected, pooled and precipitated with ammonium sulphate at 60% saturation. The pellet isolated by centrifugation was

- 111 -

resolubilized in NET buffer, and purified on a Sepharose 6B column (2.5 X 47 cm), eluted at 3 ml/hour/fraction. The fractions were analysed by SDS-PAGE. Fractions 23-27 were collected, the salt concentration adjusted to 0.5 M, and precipitated with PEG 6000, added from a 40% stock in water and to a final concentration of 13.3%. The pellet isolated by centrifugation was resolubilized in NET buffer, and loaded on the same Sepharose 2B column as above, eluted in the same manner. Analysis of the fractions by SDS-PAGE is shown in FIG. 6C. Fractions 43-53 were collected, and precipitated with ammonium sulphate at a saturation of 60%. The pellet isolated by centrifugation was resolubilized in water, and the obtained protein solution was extensively dialyzed against water. About 10 mg of purified protein per gram of cells could be isolated.

Examination of the virus-like particles in Electron microscopy showed that they were identical to the phage particles (FIG. 7A and 7B).

FIG. 6A shows in the top panel, the silver-stained SDS-PAGE run under reducing conditions of the fractions of the first Sepharose 4B chromatography step. Lane 1- 13 were loaded with every second fraction from fraction 20 to 44. Fraction 50 was loaded in lane 14. A second gel was loaded with the same fractions and analysed by Western blotting with an anti-serum specific for AP205, and is shown in the lower panel (FIG. 6B).

FIG. 6C shows the silver-stained SDS-PAGE run under reducing conditions of the fractions of the last Sepharose 2B chromatography step. Fractions 38-54 are loaded in Lane 1- 16.

FIG. 7A shows an EM picture of AP205 phage particles, while an EM picture of self assembled particles of recombinant AP205 VLP is shown in FIG. 7B.

EXAMPLE 19

Inhibition of RANKL-induced osteoclast formation by sera of mice
immunized with Q β -C-RANKL

- 112 -

To test whether the antibodies generated in mice immunized with Q β -C-RANKL are able to inhibit the biological activity of RANKL, an in vitro osteoclast differentiation assay was established. Bone marrow cells were therefore isolated from Balb/c mice (4 weeks of age) and incubated at a density of 10^6 /ml with recombinant mouse M-CSF (5 ng/ml) in α -MEM/10% FCS for 16 hours. Floating cells were then collected and further cultivated with M-CSF (30 ng/ml), PGE₂ (1 μ M) and different concentrations of C-RANKL. On day 4 osteoclast formation was assessed by the number of multinucleated cells staining positive for tartrate resistant acid phosphatase (TRAP). C-RANKL was found to induce a significant number of TRAP positive multinucleated cells at a concentration of 100-1000 ng/ml.

Sera of mice vaccinated with Q β -C-RANKL are then tested for their ability to inhibit the formation of osteoclasts from C-RANKL treated bone marrow cells. Osteoclast precursor cells are therefore isolated from Balb/c mice and incubated with M-CSF (30 ng/ml), PGE₂ (1 μ M), C-RANKL (1000 ng/ml) and serial dilutions of sera derived from immunized mice in α -MEM/10% FCS for 4 days. Osteoclast formation is then assayed by counting the number of TRAP positive multinucleated cells and comparison to the controls incubated with C-RANKL alone and with C-RANKL and preimmune sera.

EXAMPLE 20

Inhibition of bone loss by vaccination with Q β -C-RANKL in a murine ovariectomy model

To test whether vaccination with Q β -C-RANKL protects from bone loss induced by estrogen deficiency a mouse ovariectomy (ovx) model is established. A total of 40 C57/BL6 mice at the age of 12 weeks is therefore randomized into 4 groups and treated as follows: Group 1 is not immunized

- 113 -

and subjected to sham operation, group 2 is not immunized and subjected to ovx, group 3 is immunized with 25 µg of Q β -C-RANKL without Alum and subjected to ovx, and group 4 is immunized with 25 µg of Q β -C-RANKL with Alum and also subjected to ovx. Animals from group 3 and 4 receive immunizations at days 0, 14, 21, and 42 and all animals are operated (either sham or ovx) on day 28 and sacrificed on day 63. Body weight is measured at days 0, 28, and 63 and blood samples are collected on days 0, 14, 21, 28, 42 and 63. Antibody titres as well as bone formation and bone degradation markers are monitored continuously and bone mineral density is assessed after scarification of the animals by DXA scan of the excised vertebrae columns and pQCT scanning of excised femurs at one distal and one midshaft site.

Having now fully described the present invention in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious to one of ordinary skill in the art that the same can be performed by modifying or changing the invention within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any specific embodiment thereof, and that such modifications or changes are intended to be encompassed within the scope of the appended claims.

All publications, patents and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains, and are herein incorporated by reference to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference.

- 114 -

WHAT IS CLAIMED IS:

1. A composition comprising:
 - (a) a virus-like particle; and
 - (b) at least one antigen or antigenic determinant, wherein said antigen or said antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide, and
wherein said at least one antigen or antigenic determinant is bound to said virus-like particle.
2. The composition of claim 1, wherein said virus-like particle (a) is a recombinant virus-like particle.
3. The composition of claim 1, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, selected from the group consisting of:
 - (a) recombinant proteins of Hepatitis B virus;
 - (b) recombinant proteins of measles virus;
 - (c) recombinant proteins of Sindbis virus;
 - (d) recombinant proteins of Rotavirus;
 - (e) recombinant proteins of Foot-and-Mouth-Disease virus;
 - (f) recombinant proteins of Retrovirus;
 - (g) recombinant proteins of Norwalk virus;
 - (h) recombinant proteins of Alphavirus;
 - (i) recombinant proteins of human Papilloma virus;
 - (j) recombinant proteins of Polyoma virus;
 - (k) recombinant proteins of bacteriophages;
 - (l) recombinant proteins of RNA-phages;
 - (m) recombinant proteins of Ty;
 - (n) recombinant proteins of Q β -phage;
 - (o) recombinant proteins of GA-phage;

- 115 -

- (p) recombinant proteins of fr-phage; and
- (q) fragments of any of the recombinant proteins from (a) to (p).

4. The composition of claim 1, wherein said virus-like particle is Hepatitis B virus core antigen.

5. The composition of claim 1, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, of a RNA-phage.

6. The composition of claim 5, wherein said RNA-phage is selected from the group consisting of:

- (a) bacteriophage Q β ;
- (b) bacteriophage R17;
- (c) bacteriophage fr;
- (d) bacteriophage GA;
- (e) bacteriophage SP;
- (f) bacteriophage MS2;
- (g) bacteriophage M11;
- (h) bacteriophage MX1;
- (i) bacteriophage NL95;
- (k) bacteriophage f2;
- (l) bacteriophage PP7; and
- (m) bacteriophage AP205.

7. The composition of claim 1, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, of RNA-phage Q β .

8. The composition of claim 1, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, of RNA-phage fr.

- 116 -

9. The composition of claim 1, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, of RNA-phage AP205.

10. The composition of claim 1, wherein said at least one antigen or antigenic determinant is bound to said virus-like particle by at least one covalent bond.

11. The composition of claim 1, wherein said at least one antigen or antigenic determinant is bound to said virus-like particle by at least one covalent bond, and wherein said covalent bond is a non-peptide bond.

12. The composition of claim 1, wherein said at least one antigen or antigenic determinant is fused to said virus-like particle.

13. The composition of claim 1, wherein said antigen or antigenic determinant is a RANKL protein or RANKL fragment.

14. The composition of claim 1, wherein said antigen or antigenic determinant is a human RANKL protein or a human RANKL fragment

15. The composition of claim 1, wherein said antigen or antigenic determinant has an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:79;
- (b) the amino acid sequence of SEQ ID NO:80;
- (c) the amino acid sequence of SEQ ID NO:81;
- (d) the amino acid sequence of SEQ ID NO:82;
- (e) the amino acid sequence of SEQ ID NO:83;
- (f) the amino acid sequence of SEQ ID NO:84
- (g) the amino acid sequence of SEQ ID NO:100
- (h) the amino acid sequence of SEQ ID NO:101;

- 117 -

- (i) the amino acid sequence of a fragment of any of SEQ ID NO:79-84, 100, 101.

16. The composition of claim 1, wherein said antigen or antigenic determinant is a RANKL peptide.

17. The composition of claim 16, wherein said RANKL peptide is a human RANKL peptide.

18. The composition of claim 1, wherein said antigen or antigenic determinant is a RANKL peptide comprising at least one antigenic site of a RANKL protein.

19. The composition of claim 1, wherein said antigen or antigenic determinant is a RANKL peptide comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:87;
- (b) the amino acid sequence of SEQ ID NO:88;
- (c) the amino acid sequence of SEQ ID NO:89;
- (d) the amino acid sequence of SEQ ID NO:90;
- (e) the amino acid sequence of SEQ ID NO:91;
- (f) the amino acid sequence of SEQ ID NO:92;
- (f) the amino acid sequence of SEQ ID NO:93;
- (k) the amino acid sequence of a fragment of any of SEQ ID NO:87-93.

20. The composition of claim 1, wherein said antigen or antigenic determinant further comprises at least one second attachment site selected from the group consisting of:

- (i) an attachment site not naturally occurring with said antigen or antigenic determinant; and

- 118 -

(ii) an attachment site naturally occurring with said antigen or antigenic determinant.

21. The composition of claim 20, wherein said antigen or antigenic determinant with said at least second attachment site comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:104;
- (b) the amino acid sequence of SEQ ID NO:105;
- (c) the amino acid sequence of SEQ ID NO:106;
- (d) the amino acid sequence of SEQ ID NO:107;
- (e) the amino acid sequence of SEQ ID NO:108;
- (f) the amino acid sequence of SEQ ID NO:109;
- (g) the amino acid sequence of SEQ ID NO:110;
- (h) the amino acid sequence of a fragment of any of SEQ ID NO:104-110.

22. A composition comprising:

- (a) a core particle with at least one first attachment site; and
- (b) at least one antigen or antigenic determinant with at least one second attachment site,

wherein said antigen or antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide, and wherein said second attachment site being selected from the group consisting of:

- (i) an attachment site not naturally occurring with said antigen or antigenic determinant; and
- (ii) an attachment site naturally occurring with said antigen or antigenic determinant,

wherein said second attachment site is capable of association to said first attachment site; and wherein said antigen or antigenic determinant

- 119 -

and said core particle interact through said association to form an ordered and repetitive antigen array.

23. The composition of claim 22, wherein said second attachment site is capable of association to said first attachment site through at least one non-peptide bond.

24. The composition of claim 22, wherein said core particle is selected from the group consisting of:

- i) a virus;
- ii) a virus-like particle;
- iii) a bacteriophage;
- iv) a bacterial pilus;
- v) a viral capsid particle; and
- vi) a recombinant form of (i), (ii), (iii), (iv) or (v).

25. The composition of claim 22, wherein said core particle is selected from the group consisting of:

- i) a virus-like particle;
- ii) a bacterial pilus; and
- iii) a virus-like particle of a RNA-phage.

26. The composition of claim 22, wherein said virus-like particle (a) is a recombinant virus-like particle.

27. The composition of claim 22, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, selected from the group consisting of:

- (a) recombinant proteins of Hepatitis B virus;
- (b) recombinant proteins of measles virus;
- (c) recombinant proteins of Sindbis virus;

- 120 -

- (d) recombinant proteins of Rotavirus;
- (e) recombinant proteins of Foot-and-Mouth-Disease virus;
- (f) recombinant proteins of Retrovirus;
- (g) recombinant proteins of Norwalk virus;
- (h) recombinant proteins of Alphavirus;
- (i) recombinant proteins of human Papilloma virus;
- (j) recombinant proteins of Polyoma virus;
- (k) recombinant proteins of bacteriophages;
- (l) recombinant proteins of RNA-phages;
- (m) recombinant proteins of Ty;
- (n) recombinant proteins of Q β -phage;
- (o) recombinant proteins of GA-phage;
- (p) recombinant proteins of fr-phage; and
- (q) fragments of any of the recombinant proteins from (a) to (p).

28. The composition of claim 22, wherein said virus-like particle is Hepatitis B virus core antigen.

29. The composition of claim 22, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, of a RNA-phage.

30. The composition of claim 29, wherein said RNA-phage is selected from the group consisting of:

- (a) bacteriophage Q β ;
- (b) bacteriophage R17;
- (c) bacteriophage fr;
- (d) bacteriophage GA;
- (e) bacteriophage SP;
- (f) bacteriophage MS2;
- (g) bacteriophage M11;

- 121 -

- (h) bacteriophage MX1;
- (i) bacteriophage NL95;
- (k) bacteriophage f2;
- (l) bacteriophage PP7; and
- (m) bacteriophage AP205.

31. The composition of claim 22, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, of RNA-phage Q β .

32. The composition of claim 22, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, of RNA-phage fr.

33. The composition of claim 22, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, of RNA-phage AP205

34. The composition of claim 22, wherein said second attachment site is capable of association to said first attachment site through at least one covalent bond.

35. The composition of claim 22, wherein said second attachment site is capable of association to said first attachment site through at least one covalent bond, and wherein said covalent bond is a non-peptide bond.

36. The composition of claim 22, wherein said antigen or antigenic determinant is a RANKL protein or RANKL fragment.

37. The composition of claim 22, wherein said antigen or antigenic determinant is a human RANKL protein or a human RANKL fragment

38. The composition of claim 22, wherein said antigen or antigenic determinant has an amino acid sequence selected from the group consisting of:

- 122 -

- (a) the amino acid sequence of SEQ ID NO:79;
- (b) the amino acid sequence of SEQ ID NO:80;
- (c) the amino acid sequence of SEQ ID NO:81;
- (d) the amino acid sequence of SEQ ID NO:82;
- (e) the amino acid sequence of SEQ ID NO:83;
- (f) the amino acid sequence of SEQ ID NO:84
- (g) the amino acid sequence of SEQ ID NO:100
- (h) the amino acid sequence of SEQ ID NO:101;
- (i) the amino acid sequence of a fragment of any of SEQ ID NO:79-84, 100, 101.

39. The composition of claim 22, wherein said antigen or antigenic determinant is a RANKL peptide.

40. The composition of claim 39, wherein said RANKL peptide is a human RANKL peptide.

41. The composition of claim 22, wherein said antigen or antigenic determinant is a RANKL peptide comprising at least one antigenic site of a RANKL protein.

42. The composition of claim 22, wherein said antigen or antigenic determinant is a RANKL peptide comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:87;
- (b) the amino acid sequence of SEQ ID NO:88;
- (c) the amino acid sequence of SEQ ID NO:89;
- (d) the amino acid sequence of SEQ ID NO:90;
- (e) the amino acid sequence of SEQ ID NO:91;
- (f) the amino acid sequence of SEQ ID NO:92;
- (f) the amino acid sequence of SEQ ID NO:93;

- 123 -

- (k) the amino acid sequence of a fragment of any of SEQ ID NO:87-93.

43. The composition of claim 22, wherein said antigen or antigenic determinant with said at least second attachment site comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:104;
- (b) the amino acid sequence of SEQ ID NO:105;
- (c) the amino acid sequence of SEQ ID NO:106;
- (d) the amino acid sequence of SEQ ID NO:107;
- (e) the amino acid sequence of SEQ ID NO:108;
- (f) the amino acid sequence of SEQ ID NO:109;
- (g) the amino acid sequence of SEQ ID NO:110;
- (h) the amino acid sequence of a fragment of any of SEQ ID NO:104-110.

44. A pharmaceutical composition comprising:

- (a) the composition of claim 1; and
- (b) an acceptable pharmaceutical carrier.

45. A pharmaceutical composition comprising:

- (a) the composition of claim 22; and
- (b) an acceptable pharmaceutical carrier.

46. A vaccine composition comprising a composition, wherein said composition comprises:

- (a) a virus-like particle; and
- (b) at least one antigen or antigenic determinant, wherein said antigen or said antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide, and

- 124 -

wherein said at least one antigen or antigenic determinant is bound to said virus-like particle.

47. The vaccine composition of claim 46, further comprising an adjuvant.

48. The vaccine composition of claim 46, wherein said virus-like particle (a) is a recombinant virus-like particle.

49. The vaccine composition of claim 46, wherein said virus-like particle is Hepatitis B virus core antigen.

50. The vaccine composition of claim 46, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, of a RNA-phage.

51. The vaccine composition of claim 46, wherein said virus-like particle comprises recombinant proteins or fragments thereof, of RNA-phage Q β .

52. The vaccine composition of claim 46, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, of RNA-phage fr.

53. The vaccine composition of claim 46, wherein said virus-like particle comprises recombinant proteins, or fragments thereof, of RNA-phage AP205.

54. The vaccine composition of claim 46, wherein said at least one antigen or antigenic determinant is bound to said virus-like particle by at least one covalent bond, and wherein said covalent bond is a non-peptide bond.

- 125 -

55. The vaccine composition of claim 46, wherein said at least one antigen or antigenic determinant is fused to said virus-like particle.

56. The composition of claim 46, wherein said antigen or antigenic determinant is a RANKL protein or RANKL fragment.

57. The vaccine composition of claim 46, wherein said antigen or antigenic determinant is a human RANKL protein or a human RANKL fragment

58. The vaccine composition of claim 46, wherein said antigen or antigenic determinant has an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:79;
- (b) the amino acid sequence of SEQ ID NO:80;
- (c) the amino acid sequence of SEQ ID NO:81;
- (d) the amino acid sequence of SEQ ID NO:82;
- (e) the amino acid sequence of SEQ ID NO:83;
- (f) the amino acid sequence of SEQ ID NO:84
- (g) the amino acid sequence of SEQ ID NO:100
- (h) the amino acid sequence of SEQ ID NO:101;
- (i) the amino acid sequence of a fragment of any of SEQ ID NO:79-84, 100, 101.

59. The vaccine composition of claim 46, wherein said antigen or antigenic determinant is a RANKL peptide.

60. The vaccine composition of claim 59, wherein said RANKL peptide is a human RANKL peptide.

- 126 -

61. The vaccine composition of claim 46, wherein said antigen or antigenic determinant is a RANKL peptide comprising at least one antigenic site of a RANKL protein.

62. The vaccine composition of claim 46, wherein said antigen or antigenic determinant is a RANKL peptide comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:87;
- (b) the amino acid sequence of SEQ ID NO:88;
- (c) the amino acid sequence of SEQ ID NO:89;
- (d) the amino acid sequence of SEQ ID NO:90;
- (e) the amino acid sequence of SEQ ID NO:91;
- (f) the amino acid sequence of SEQ ID NO:92;
- (f) the amino acid sequence of SEQ ID NO:93;
- (k) the amino acid sequence of a fragment of any of SEQ ID NO:87-93.

63. The vaccine composition of claim 46, wherein said antigen or antigenic determinant further comprises at least one second attachment site selected from the group consisting of:

- (i) an attachment site not naturally occurring with said antigen or antigenic determinant; and
- (ii) an attachment site naturally occurring with said antigen or antigenic determinant.

64. The vaccine composition of claim 63, wherein said antigen or antigenic determinant with said at least second attachment site comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:104;
- (b) the amino acid sequence of SEQ ID NO:105;
- (c) the amino acid sequence of SEQ ID NO:106;

- 127 -

- (d) the amino acid sequence of SEQ ID NO:107;
- (e) the amino acid sequence of SEQ ID NO:108;
- (f) the amino acid sequence of SEQ ID NO:109;
- (g) the amino acid sequence of SEQ ID NO:110;
- (h) the amino acid sequence of a fragment of any of SEQ ID NO:104-110.

65. A process for producing a composition of claim 1 comprising:

- (a) providing a virus-like particle; and
- (b) providing at least one antigen or antigenic determinant, wherein said antigen or said antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide;
- (c) combining said virus-like particle and said at least one antigen or antigenic determinant so that said at least one antigen or antigenic determinant is bound to said virus-like particle.

66. A process for producing a composition of claim 22 comprising:

- (a) providing a core particle with at least one first attachment site;
- (b) providing at least one antigen or antigenic determinant with at least one second attachment site, wherein said antigen or antigenic determinant is a RANKL protein, RANKL fragment or RANKL peptide, and wherein said second attachment site being selected from the group consisting of:
 - (i) an attachment site not naturally occurring with said antigen or antigenic determinant; and
 - (ii) an attachment site naturally occurring with said antigen or antigenic determinant; and

wherein said second attachment site is capable of association to said first attachment site; and

- 128 -

- (c) combining said core particle and said at least one antigen or antigenic determinant, wherein said antigen or antigenic determinant and said core particle interact through said association to form an ordered and repetitive antigen array.

67. A method of immunization comprising administering the composition of claim 1 to an animal or human.

68. The method of immunization of claim 67, wherein said antigen or antigenic determinant is a self-antigen.

69. The method of immunization of claim 67, wherein said animal is a human, and wherein said antigen or antigenic determinant is a human RANKL protein, RANKL fragment or RANKL peptide.

70. Composition of claim 1 for use as a medicament.

71. Composition of claim 22 for use as a medicament.


72. Use of a composition of claim 1 for the manufacture of a medicament for treatment of bone diseases.

73. Use of a composition of claim 22 for the manufacture of a medicament for treatment of bone diseases.

74. Use of a composition of claim 1, wherein said composition is used in combination with at least one medicament suitable to treat bone diseases.

- 129 -

75. Use of a composition of claim 22, wherein said composition is used in combination with at least one medicament suitable to treat bone diseases.

**FIG. 1**





FIG 2A

BEST AVAILABLE COPY

**FIG 2B**

BEST AVAILABLE COPY







FIG 2C

BEST AVAILABLE COPY

**FIG 3A**

BEST AVAILABLE COPY

**FIG 3B****FIG 3C**

IgG against C-RANKL





FIG. 4

**FIG 5A**




FIG 5B

FIG. 6A






FIG. 6B



BEST AVAILABLE COPY

FIG. 6C



1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16

BEST AVAILABLE COPY




FIG. 7B




FIG. 7A

BEST AVAILABLE COPY

-1-

SEQUENCE LISTING

SEQ ID NO: 1

Met Lys Lys Thr Leu Leu Gly Ser Leu Ile Leu Leu Ala Phe Ala Gly
 1 5 10 15

Asn Val Gln Ala Ala Ala Asn Ala Asp Thr Ser Gly Thr Val Thr Phe
 20 25 30

Phe Gly Lys Val Val Glu Asn Thr Cys Gln Val Asn Gln Asp Ser Glu
 35 40 45

Tyr Glu Cys Asn Leu Asn Asp Val Gly Lys Asn His Leu Ser Gln Gln
 50 55 60

Gly Tyr Thr Ala Met Gln Thr Pro Phe Thr Ile Thr Leu Glu Asn Cys
 65 70 75 80

Asn Val Thr Thr Thr Asn Asn Lys Pro Lys Ala Thr Lys Val Gly Val
 85 90 95

Tyr Phe Tyr Ser Trp Glu Ile Ala Asp Lys Asp Asn Lys Tyr Thr Leu
 100 105 110

Lys Asn Ile Lys Glu Asn Thr Gly Thr Asn Asp Ser Ala Asn Lys Val
 115 120 125

Asn Ile Gln Leu Leu Glu Asp Asn Gly Thr Ala Glu Ile Lys Val Val
 130 135 140

Gly Lys Thr Thr Thr Asp Phe Thr Ser Glu Asn His Asn Gly Ala Gly
 145 150 155 160

Ala Asp Pro Val Ala Thr Asn Lys His Ile Ser Ser Leu Thr Pro Leu
 165 170 175

Asn Asn Gln Asn Ser Ile Asn Leu His Tyr Ile Ala Gln Tyr Tyr Ala
 180 185 190

Thr Gly Val Ala Glu Ala Gly Lys Val Pro Ser Ser Val Asn Ser Gln
 195 200 205

Ile Ala Tyr Glu
 210

-2-

SEQ ID NO: 2

Met Lys Ala Gln Met Gln Lys Gly Phe Thr Leu Ile Glu Leu Met Ile
 1 5 10 15

Val Val Ala Ile Ile Gly Ile Leu Ala Ala Ile Ala Leu Pro Ala Tyr
 20 25 30

Gln Asp Tyr Thr Val Arg Ser Asn Ala Ala Ala Ala Leu Ala Glu Ile
 35 40 45

Thr Pro Gly Lys Ile Gly Phe Glu Gln Ala Ile Asn Glu Gly Lys Thr
 50 55 60

Pro Ser Leu Thr Ser Thr Asp Glu Gly Tyr Ile Gly Ile Thr Asp Ser
 65 70 75 80

Thr Ser Tyr Cys Asp Val Asp Leu Asp Thr Ala Ala Asp Gly His Ile
 85 90 95

Glu Cys Thr Ala Lys Gly Gly Asn Ala Gly Lys Phe Asp Gly Lys Thr
 100 105 110

Ile Thr Leu Asn Arg Thr Ala Asp Gly Glu Trp Ser Cys Ala Ser Thr
 115 120 125

Leu Asp Ala Lys Tyr Lys Pro Gly Lys Cys Ser
 130 135

SEQ ID NO: 3

Met Thr Lys Phe Val Thr Arg Phe Leu Lys Asp Glu Ser Gly Ala Thr
 1 5 10 15

Ala Ile Glu Tyr Gly Leu Ile Val Ala Leu Ile Ala Val Val Ile Val
 20 25 30

Thr Ala Val Thr Thr Leu Gly Thr Asn Leu Arg Thr Ala Phe Thr Lys
 35 40 45

Ala Gly Ala Ala Val Ser Thr Ala Ala Gly Thr
 50 55

-3-

SEQ ID NO: 4

Met Ala Val Val Ser Phe Gly Val Asn Ala Ala Pro Thr Ile Pro Gln
1 5 10 15

Gly Gln Gly Lys Val Thr Phe Asn Gly Thr Val Val Asp Ala Pro Cys
20 25 30

Ser Ile Ser Gln Lys Ser Ala Asp Gln Ser Ile Asp Phe Gly Gln Leu
35 40 45

Ser Lys Ser Phe Leu Glu Ala Gly Gly Val Ser Lys Pro Met Asp Leu
50 55 60

Asp Ile Glu Leu Val Asn Cys Asp Ile Thr Ala Phe Lys Gly Gly Asn
65 70 75 80

Gly Ala Gln Lys Gly Thr Val Lys Leu Ala Phe Thr Gly Pro Ile Val
85 90 95

Asn Gly His Ser Asp Glu Leu Asp Thr Asn Gly Gly Thr Gly Thr Ala
100 105 110

Ile Val Val Gln Gly Ala Gly Lys Asn Val Val Phe Asp Gly Ser Glu
115 120 125

Gly Asp Ala Asn Thr Leu Lys Asp Gly Glu Asn Val Leu His Tyr Thr
130 135 140

Ala Val Val Lys Lys Ser Ser Ala Val Gly Ala Ala Val Thr Glu Gly
145 150 155 160

Ala Phe Ser Ala Val Ala Asn Phe Asn Leu Thr Tyr Gln
165 170

SEQ ID NO: 5

Met Ala Val Val Ser Phe Gly Val Asn Ala Ala Pro Thr Ile Pro Gln
1 5 10 15

Gly Gln Gly Lys Val Thr Phe Asn Gly Thr Val Val Asp Ala Pro Cys
20 25 30

-4-

Ser Ile Ser Gln Lys Ser Ala Asp Gln Ser Ile Asp Phe Gly Gln Leu
 35 40 45

Ser Lys Ser Phe Leu Glu Ala Gly Gly Val Ser Lys Pro Met Asp Leu
 50 55 60

Asp Ile Glu Leu Val Asn Cys Asp Ile Thr Ala Phe Lys Gly Gly Asn
 65 70 75 80

Gly Ala Gln Lys Gly Thr Val Lys Leu Ala Phe Thr Gly Pro Ile Val
 85 90 95

Asn Gly His Ser Asp Glu Leu Asp Thr Asn Gly Gly Thr Gly Thr Ala
 100 105 110

Ile Val Val Gln Gly Ala Gly Lys Asn Val Val Phe Asp Gly Ser Glu
 115 120 125

Gly Asp Ala Asn Thr Leu Lys Asp Gly Glu Asn Val Leu His Tyr Thr
 130 135 140

Ala Val Val Lys Lys Ser Ser Ala Val Gly Ala Ala Val Thr Glu Gly
 145 150 155 160

Ala Phe Ser Ala Val Ala Asn Phe Asn Leu Thr Tyr Gln
 165 170

SEQ ID NO: 6

Met Lys Ile Lys Thr Leu Ala Ile Val Val Leu Ser Ala Leu Ser Leu
 1 5 10 15

Ser Ser Thr Ala Ala Leu Ala Ala Thr Thr Val Asn Gly Gly Thr
 20 25 30

Val His Phe Lys Gly Glu Val Val Asn Ala Ala Cys Ala Val Asp Ala
 35 40 45

Gly Ser Val Asp Gln Thr Val Gln Leu Gly Gln Val Arg Thr Ala Ser
 50 55 60

Leu Ala Gln Glu Gly Ala Thr Ser Ser Ala Val Gly Phe Asn Ile Gln
 65 70 75 80

-5-

Leu Asn Asp Cys Asp Thr Asn Val Ala Ser Lys Ala Ala Val Ala Phe
 85 90 95

Leu Gly Thr Ala Ile Asp Ala Gly His Thr Asn Val Leu Ala Leu Gln
 100 105 110

Ser Ser Ala Ala Gly Ser Ala Thr Asn Val Gly Val Gln Ile Leu Asp
 115 120 125

Arg Thr Gly Ala Ala Leu Thr Leu Asp Gly Ala Thr Phe Ser Ser Glu
 130 135 140

Thr Thr Leu Asn Asn Gly Thr Asn Thr Ile Pro Phe Gln Ala Arg Tyr
 145 150 155 160

Phe Ala Gly Ala Ala Thr Pro Gly Ala Ala Asn Ala Asp Ala Thr Phe
 165 170 175

Lys Val Gln Tyr Gln
 180

SEQ ID NO: 7

Met Ala Val Val Ser Phe Gly Val Asn Ala Ala Pro Thr Thr Pro Gln
 1 5 10 15

Gly Gln Gly Arg Val Thr Phe Asn Gly Thr Val Val Asp Ala Pro Cys
 20 25 30

Ser Ile Ser Gln Lys Ser Ala Asp Gln Ser Ile Asp Phe Gly Gln Leu
 35 40 45

Ser Lys Ser Phe Leu Ala Asn Asp Gly Gln Ser Lys Pro Met Asn Leu
 50 55 60

Asp Ile Glu Leu Val Asn Cys Asp Ile Thr Ala Phe Lys Asn Gly Asn
 65 70 75 80

Ala Lys Thr Gly Ser Val Lys Leu Ala Phe Thr Gly Pro Thr Val Ser
 85 90 95

Gly His Pro Ser Glu Leu Ala Thr Asn Gly Gly Pro Gly Thr Ala Ile
 100 105 110

-6-

Met Ile Gln Ala Ala Gly Lys Asn Val Pro Phe Asp Gly Thr Glu Gly
 115 120 125

Asp Pro Asn Leu Leu Lys Asp Gly Asp Asn Val Leu His Tyr Thr Thr
 130 135 140

Val Gly Lys Lys Ser Ser Asp Gly Asn Ala Gln Ile Thr Glu Gly Ala
 145 150 155 160

Phe Ser Gly Val Ala Thr Phe Asn Leu Ser Tyr Gln
 165 170

SEQ ID NO: 8

Met Lys Ile Lys Thr Leu Ala Ile Val Val Leu Ser Ala Leu Ser Leu
 1 5 10 15

Ser Ser Thr Thr Ala Leu Ala Ala Ala Thr Thr Val Asn Gly Gly Thr
 20 25 30

Val His Phe Lys Gly Glu Val Val Asn Ala Ala Cys Ala Val Asp Ala
 35 40 45

Gly Ser Val Asp Gln Thr Val Gln Leu Gly Gln Val Arg Thr Ala Ser
 50 55 60

Leu Ala Gln Glu Gly Ala Thr Ser Ser Ala Val Gly Phe Asn Ile Gln
 65 70 75 80

Leu Asn Asp Cys Asp Thr Asn Val Ala Ser Lys Ala Ala Val Ala Phe
 85 90 95

Leu Gly Thr Ala Ile Asp Ala Gly His Thr Asn Val Leu Ala Leu Gln
 100 105 110

Ser Ser Ala Ala Gly Ser Ala Thr Asn Val Gly Val Gln Ile Leu Asp
 115 120 125

Arg Thr Gly Ala Ala Leu Thr Leu Asp Gly Ala Thr Phe Ser Ser Glu
 130 135 140

Thr Thr Leu Asn Asn Gly Thr Asn Thr Ile Pro Phe Gln Ala Arg Tyr
 145 150 155 160

-7-

Phe Ala Thr Gly Ala Ala Thr Pro Gly Ala Ala Asn Ala Asp Ala Thr
 165 170 175

Phe Lys Val Gln Tyr Gln
 180

SEQ ID NO: 9
 acgtttctgt ggctcgacgc atcttctctca ttcttctctc caaaaaccac ctcatgcaat 60
 ataaacatct ataaataaag ataacaaata gaatattaag ccaacaaata aactgaaaaa 120
 gtttgctcgc gatgctttac ctctatgagt caaaatggcc ccaatgtttc atcttttggg 180
 ggaaactgtg cagtgttggc agtcaaactc gttgacaaac aaagtgtaca gaacgactgc 240
 ccatgtcgat ttagaaatag ttttttgaaa ggaaagcagc atgaaaatta aaactctggc 300
 aatcgttggt ctgtcggctc tgtccctcag ttctacgacg gctctggccg ctgccacgac 360
 ggtaaatggt gggaccgttc actttaaaagg ggaagttggt aacgccgctt gcgcagttga 420
 tgcaggctct gttgatcaaa ccgttcagtt aggacaggtt cgtaccgcat cgctggcaca 480
 ggaaggagca accagttctg ctgtcgggtt taacattcag ctgaatgatt gcgataccaa 540
 tgttgcattc aaagccgctg ttgccttttt aggtacggcg attgatgcgg gtcataccaa 600
 cgttctggct ctgcagagtt cagctgcggg tagcgcaaca aacgttggtg tgcagatcct 660
 ggacagaacg ggtgctgcgc tgacgttgga tgggtgcgaca tttagttcag aaacaaccct 720
 gaataacgga accaatacca ttccgttcca ggcgcgttat tttgcaaccg gggccgcaac 780
 cccgggtgct gctaatacgg atgcgacctt caaggttcag tatcaataac ctacctaggt 840
 tcagggacgt tca 853

SEQ ID NO: 10

Ala Lys Leu Glu Thr Val Thr Leu Gly Asn Ile Gly Lys Asp Gly Lys
 1 5 10 15

Gln Thr Leu Val Leu Asn Pro Arg Gly Val Asn Pro Thr Asn Gly Val
 20 25 30

Ala Ser Leu Ser Gln Ala Gly Ala Val Pro Ala Leu Glu Lys Arg Val
 35 40 45

-8-

Thr Val Ser Val Ser Gln Pro Ser Arg Asn Arg Lys Asn Tyr Lys Val
 50 55 60

Gln Val Lys Ile Gln Asn Pro Thr Ala Cys Thr Ala Asn Gly Ser Cys
 65 70 75 80

Asp Pro Ser Val Thr Arg Gln Ala Tyr Ala Asp Val Thr Phe Ser Phe
 85 90 95

Thr Gln Tyr Ser Thr Asp Glu Glu Arg Ala Phe Val Arg Thr Glu Leu
 100 105 110

Ala Ala Leu Leu Ala Ser Pro Leu Leu Ile Asp Ala Ile Asp Gln Leu
 115 120 125

Asn Pro Ala Tyr
 130

SEQ ID NO: 11

Met Ala Lys Leu Glu Thr Val Thr Leu Gly Asn Ile Gly Lys Asp Gly
 1 5 10 15

Lys Gln Thr Leu Val Leu Asn Pro Arg Gly Val Asn Pro Thr Asn Gly
 20 25 30

Val Ala Ser Leu Ser Gln Ala Gly Ala Val Pro Ala Leu Glu Lys Arg
 35 40 45

Val Thr Val Ser Val Ser Gln Pro Ser Arg Asn Arg Lys Asn Tyr Lys
 50 55 60

Val Gln Val Lys Ile Gln Asn Pro Thr Ala Cys Thr Ala Asn Gly Ser
 65 70 75 80

Cys Asp Pro Ser Val Thr Arg Gln Ala Tyr Ala Asp Val Thr Phe Ser
 85 90 95

Phe Thr Gln Tyr Ser Thr Asp Glu Glu Arg Ala Phe Val Arg Thr Glu
 100 105 110

Leu Ala Ala Leu Leu Ala Ser Pro Leu Leu Ile Asp Ala Ile Asp Gln
 115 120 125

-9-

Leu Asn Pro Ala Tyr Trp Thr Leu Leu Ile Ala Gly Gly Gly Ser Gly
 130 135 140

Ser Lys Pro Asp Pro Val Ile Pro Asp Pro Pro Ile Asp Pro Pro Pro
 145 150 155 160

Gly Thr Gly Lys Tyr Thr Cys Pro Phe Ala Ile Trp Ser Leu Glu Glu
 165 170 175

Val Tyr Glu Pro Pro Thr Lys Asn Arg Pro Trp Pro Ile Tyr Asn Ala
 180 185 190

Val Glu Leu Gln Pro Arg Glu Phe Asp Val Ala Leu Lys Asp Leu Leu
 195 200 205

Gly Asn Thr Lys Trp Arg Asp Trp Asp Ser Arg Leu Ser Tyr Thr Thr
 210 215 220

Phe Arg Gly Cys Arg Gly Asn Gly Tyr Ile Asp Leu Asp Ala Thr Tyr
 225 230 235 240

Leu Ala Thr Asp Gln Ala Met Arg Asp Gln Lys Tyr Asp Ile Arg Glu
 245 250 255

Gly Lys Lys Pro Gly Ala Phe Gly Asn Ile Glu Arg Phe Ile Tyr Leu
 260 265 270

Lys Ser Ile Asn Ala Tyr Cys Ser Leu Ser Asp Ile Ala Ala Tyr His
 275 280 285

Ala Asp Gly Val Ile Val Gly Phe Trp Arg Asp Pro Ser Ser Gly Gly
 290 295 300

Ala Ile Pro Phe Asp Phe Thr Lys Phe Asp Lys Thr Lys Cys Pro Ile
 305 310 315 320

Gln Ala Val Ile Val Val Pro Arg Ala
 325

SEQ ID NO: 12

Ala Ser Asn Phe Thr Gln Phe Val Leu Val Asn Asp Gly Gly Thr Gly
 1 5 10 15

-10-

Asn Val Thr Val Ala Pro Ser Asn Phe Ala Asn Gly Val Ala Glu Trp
20 25 30

Ile Ser Ser Asn Ser Arg Ser Gln Ala Tyr Lys Val Thr Cys Ser Val
35 40 45

Arg Gln Ser Ser Ala Gln Asn Arg Lys Tyr Thr Ile Lys Val Glu Val
50 55 60

Pro Lys Val Ala Thr Gln Thr Val Gly Gly Val Glu Leu Pro Val Ala
65 70 75 80

Ala Trp Arg Ser Tyr Leu Asn Met Glu Leu Thr Ile Pro Ile Phe Ala
85 90 95

Thr Asn Ser Asp Cys Glu Leu Ile Val Lys Ala Met Gln Gly Leu Leu
100 105 110

Lys Asp Gly Asn Pro Ile Pro Ser Ala Ile Ala Ala Asn Ser Gly Ile
115 120 125

Tyr

SEQ ID NO: 13

Met Ala Ser Asn Phe Glu Glu Phe Val Leu Val Asp Asn Gly Gly Thr
1 5 10 15

Gly Asp Val Lys Val Ala Pro Ser Asn Phe Ala Asn Gly Val Ala Glu
20 25 30

Trp Ile Ser Ser Asn Ser Arg Ser Gln Ala Tyr Lys Val Thr Cys Ser
35 40 45

Val Arg Gln Ser Ser Ala Asn Asn Arg Lys Tyr Thr Val Lys Val Glu
50 55 60

Val Pro Lys Val Ala Thr Gln Val Gln Gly Gly Val Glu Leu Pro Val
65 70 75 80

Ala Ala Trp Arg Ser Tyr Met Asn Met Glu Leu Thr Ile Pro Val Phe
85 90 95

-11-

Ala Thr Asn Asp Asp Cys Ala Leu Ile Val Lys Ala Leu Gln Gly Thr
 100 105 110

Phe Lys Thr Gly Asn Pro Ile Ala Thr Ala Ile Ala Ala Asn Ser Gly
 115 120 125

Ile Tyr
 130

SEQ ID NO: 14

Met Ala Thr Leu Arg Ser Phe Val Leu Val Asp Asn Gly Gly Thr Gly
 1 5 10 15

Asn Val Thr Val Val Pro Val Ser Asn Ala Asn Gly Val Ala Glu Trp
 20 25 30

Leu Ser Asn Asn Ser Arg Ser Gln Ala Tyr Arg Val Thr Ala Ser Tyr
 35 40 45

Arg Ala Ser Gly Ala Asp Lys Arg Lys Tyr Ala Ile Lys Leu Glu Val
 50 55 60

Pro Lys Ile Val Thr Gln Val Val Asn Gly Val Glu Leu Pro Gly Ser
 65 70 75 80

Ala Trp Lys Ala Tyr Ala Ser Ile Asp Leu Thr Ile Pro Ile Phe Ala
 85 90 95

Ala Thr Asp Asp Val Thr Val Ile Ser Lys Ser Leu Ala Gly Leu Phe
 100 105 110

Lys Val Gly Asn Pro Ile Ala Glu Ala Ile Ser Ser Gln Ser Gly Phe
 115 120 125

Tyr Ala
 130

SEQ ID NO: 15

Met Ala Lys Leu Asn Gln Val Thr Leu Ser Lys Ile Gly Lys Asn Gly
 1 5 10 15

-12-

Asp Gln Thr Leu Thr Leu Thr Pro Arg Gly Val Asn Pro Thr Asn Gly
20 25 30

Val Ala Ser Leu Ser Glu Ala Gly Ala Val Pro Ala Leu Glu Lys Arg
35 40 45

Val Thr Val Ser Val Ala Gln Pro Ser Arg Asn Arg Lys Asn Phe Lys
50 55 60

Val Gln Ile Lys Leu Gln Asn Pro Thr Ala Cys Thr Arg Asp Ala Cys
65 70 75 80

Asp Pro Ser Val Thr Arg Ser Ala Phe Ala Asp Val Thr Leu Ser Phe
85 90 95

Thr Ser Tyr Ser Thr Asp Glu Glu Arg Ala Leu Ile Arg Thr Glu Leu
100 105 110

Ala Ala Leu Leu Ala Asp Pro Leu Ile Val Asp Ala Ile Asp Asn Leu
115 120 125

Asn Pro Ala Tyr
130

SEQ ID NO: 16

Ala Lys Leu Asn Gln Val Thr Leu Ser Lys Ile Gly Lys Asn Gly Asp
1 5 10 15

Gln Thr Leu Thr Leu Thr Pro Arg Gly Val Asn Pro Thr Asn Gly Val
20 25 30

Ala Ser Leu Ser Glu Ala Gly Ala Val Pro Ala Leu Glu Lys Arg Val
35 40 45

Thr Val Ser Val Ala Gln Pro Ser Arg Asn Arg Lys Asn Phe Lys Val
50 55 60

Gln Ile Lys Leu Gln Asn Pro Thr Ala Cys Thr Arg Asp Ala Cys Asp
65 70 75 80

Pro Ser Val Thr Arg Ser Ala Phe Ala Asp Val Thr Leu Ser Phe Thr
85 90 95

-13-

Ser Tyr Ser Thr Asp Glu Glu Arg Ala Leu Ile Arg Thr Glu Leu Ala
 100 105 110

Ala Leu Leu Ala Asp Pro Leu Ile Val Asp Ala Ile Asp Asn Leu Asn
 115 120 125

Pro Ala Tyr Trp Ala Ala Leu Leu Val Ala Ser Ser Gly Gly Gly Asp
 130 135 140

Asn Pro Ser Asp Pro Asp Val Pro Val Val Pro Asp Val Lys Pro Pro
 145 150 155 160

Asp Gly Thr Gly Arg Tyr Lys Cys Pro Phe Ala Cys Tyr Arg Leu Gly
 165 170 175

Ser Ile Tyr Glu Val Gly Lys Glu Gly Ser Pro Asp Ile Tyr Glu Arg
 180 185 190

Gly Asp Glu Val Ser Val Thr Phe Asp Tyr Ala Leu Glu Asp Phe Leu
 195 200 205

Gly Asn Thr Asn Trp Arg Asn Trp Asp Gln Arg Leu Ser Asp Tyr Asp
 210 215 220

Ile Ala Asn Arg Arg Arg Cys Arg Gly Asn Gly Tyr Ile Asp Leu Asp
 225 230 235 240

Ala Thr Ala Met Gln Ser Asp Asp Phe Val Leu Ser Gly Arg Tyr Gly
 245 250 255

Val Arg Lys Val Lys Phe Pro Gly Ala Phe Gly Ser Ile Lys Tyr Leu
 260 265 270

Leu Asn Ile Gln Gly Asp Ala Trp Leu Asp Leu Ser Glu Val Thr Ala
 275 280 285

Tyr Arg Ser Tyr Gly Met Val Ile Gly Phe Trp Thr Asp Ser Lys Ser
 290 295 300

Pro Gln Leu Pro Thr Asp Phe Thr Gln Phe Asn Ser Ala Asn Cys Pro
 305 310 315 320

Val Gln Thr Val Ile Ile Ile Pro Ser
 325

-14-

SEQ ID NO: 17

Met Ala Ser Asn Phe Thr Gln Phe Val Leu Val Asp Asn Gly Gly Thr
 1 5 10 15

Gly Asp Val Thr Val Ala Pro Ser Asn Phe Ala Asn Gly Val Ala Glu
 20 25 30

Trp Ile Ser Ser Asn Ser Arg Ser Gln Ala Tyr Lys Val Thr Cys Ser
 35 40 45

Val Arg Gln Ser Ser Ala Gln Asn Arg Lys Tyr Thr Ile Lys Val Glu
 50 55 60

Val Pro Lys Val Ala Thr Gln Thr Val Gly Gly Val Glu Leu Pro Val
 65 70 75 80

Ala Ala Trp Arg Ser Tyr Leu Asn Met Glu Leu Thr Ile Pro Ile Phe
 85 90 95

Ala Thr Asn Ser Asp Cys Glu Leu Ile Val Lys Ala Met Gln Gly Leu
 100 105 110

Leu Lys Asp Gly Asn Pro Ile Pro Ser Ala Ile Ala Ala Asn Ser Gly
 115 120 125

Ile Tyr
 130

SEQ ID NO: 18

Met Ala Lys Leu Gln Ala Ile Thr Leu Ser Gly Ile Gly Lys Lys Gly
 1 5 10 15

Asp Val Thr Leu Asp Leu Asn Pro Arg Gly Val Asn Pro Thr Asn Gly
 20 25 30

Val Ala Ala Leu Ser Glu Ala Gly Ala Val Pro Ala Leu Glu Lys Arg
 35 40 45

Val Thr Ile Ser Val Ser Gln Pro Ser Arg Asn Arg Lys Asn Tyr Lys
 50 55 60

-15-

Val Gln Val Lys Ile Gln Asn Pro Thr Ser Cys Thr Ala Ser Gly Thr
65 70 75 80

Cys Asp Pro Ser Val Thr Arg Ser Ala Tyr Ser Asp Val Thr Phe Ser
85 90 95

Phe Thr Gln Tyr Ser Thr Val Glu Glu Arg Ala Leu Val Arg Thr Glu
100 105 110

Leu Gln Ala Leu Leu Ala Asp Pro Met Leu Val Asn Ala Ile Asp Asn
115 120 125

Leu Asn Pro Ala Tyr
130

SEQ ID NO: 19

Met Ala Lys Leu Gln Ala Ile Thr Leu Ser Gly Ile Gly Lys Asn Gly
1 5 10 15

Asp Val Thr Leu Asn Leu Asn Pro Arg Gly Val Asn Pro Thr Asn Gly
20 25 30

Val Ala Ala Leu Ser Glu Ala Gly Ala Val Pro Ala Leu Glu Lys Arg
35 40 45

Val Thr Ile Ser Val Ser Gln Pro Ser Arg Asn Arg Lys Asn Tyr Lys
50 55 60

Val Gln Val Lys Ile Gln Asn Pro Thr Ser Cys Thr Ala Ser Gly Thr
65 70 75 80

Cys Asp Pro Ser Val Thr Arg Ser Ala Tyr Ala Asp Val Thr Phe Ser
85 90 95

Phe Thr Gln Tyr Ser Thr Asp Glu Glu Arg Ala Leu Val Arg Thr Glu
100 105 110

Leu Lys Ala Leu Leu Ala Asp Pro Met Leu Ile Asp Ala Ile Asp Asn
115 120 125

-16-

Leu Asn Pro Ala Tyr
130

SEQ ID NO: 20

Met Ala Lys Leu Asn Lys Val Thr Leu Thr Gly Ile Gly Lys Ala Gly
1 5 10 15

Asn Gln Thr Leu Thr Leu Thr Pro Arg Gly Val Asn Pro Thr Asn Gly
20 25 30

Val Ala Ser Leu Ser Glu Ala Gly Ala Val Pro Ala Leu Glu Lys Arg
35 40 45

Val Thr Val Ser Val Ala Gln Pro Ser Arg Asn Arg Lys Asn Tyr Lys
50 55 60

Val Gln Ile Lys Leu Gln Asn Pro Thr Ala Cys Thr Lys Asp Ala Cys
65 70 75 80

Asp Pro Ser Val Thr Arg Ser Gly Ser Arg Asp Val Thr Leu Ser Phe
85 90 95

Thr Ser Tyr Ser Thr Glu Arg Glu Arg Ala Leu Ile Arg Thr Glu Leu
100 105 110

Ala Ala Leu Leu Lys Asp Asp Leu Ile Val Asp Ala Ile Asp Asn Leu
115 120 125

Asn Pro Ala Tyr Trp Ala Ala Leu Leu Ala Ala Ser Pro Gly Gly Gly
130 135 140

Asn Asn Pro Tyr Pro Gly Val Pro Asp Ser Pro Asn Val Lys Pro Pro
145 150 155 160

Gly Gly Thr Gly Thr Tyr Arg Cys Pro Phe Ala Cys Tyr Arg Arg Gly
165 170 175

Glu Leu Ile Thr Glu Ala Lys Asp Gly Ala Cys Ala Leu Tyr Ala Cys
180 185 190

Gly Ser Glu Ala Leu Val Glu Phe Glu Tyr Ala Leu Glu Asp Phe Leu
195 200 205

-17-

Gly Asn Glu Phe Trp Arg Asn Trp Asp Gly Arg Leu Ser Lys Tyr Asp
 210 215 220

Ile Glu Thr His Arg Arg Cys Arg Gly Asn Gly Tyr Val Asp Leu Asp
 225 230 235 240

Ala Ser Val Met Gln Ser Asp Glu Tyr Val Leu Ser Gly Ala Tyr Asp
 245 250 255

Val Val Lys Met Gln Pro Pro Gly Thr Phe Asp Ser Pro Arg Tyr Tyr
 260 265 270

Leu His Leu Met Asp Gly Ile Tyr Val Asp Leu Ala Glu Val Thr Ala
 275 280 285

Tyr Arg Ser Tyr Gly Met Val Ile Gly Phe Trp Thr Asp Ser Lys Ser
 290 295 300

Pro Gln Leu Pro Thr Asp Phe Thr Arg Phe Asn Arg His Asn Cys Pro
 305 310 315 320

Val Gln Thr Val Ile Val Ile Pro Ser Leu
 325 330

SEQ ID NO: 21

Ala Ser Asn Phe Thr Gln Phe Val Leu Val Asn Asp Gly Gly Thr Gly
 1 5 10 15

Asn Val Thr Val Ala Pro Ser Asn Phe Ala Asn Gly Val Ala Glu Trp
 20 25 30

Ile Ser Ser Asn Ser Arg Ser Gln Ala Tyr Lys Val Thr Cys Ser Val
 35 40 45

Arg Gln Ser Ser Ala Gln Asn Arg Lys Tyr Thr Ile Lys Val Glu Val
 50 55 60

Pro Lys Val Ala Thr Gln Thr Val Gly Gly Val Glu Leu Pro Val Ala
 65 70 75 80

Ala Trp Arg Ser Tyr Leu Asn Leu Glu Leu Thr Ile Pro Ile Phe Ala
 85 90 95

-18-

Thr Asn Ser Asp Cys Glu Leu Ile Val Lys Ala Met Gln Gly Leu Leu
 100 105 110

Lys Asp Gly Asn Pro Ile Pro Ser Ala Ile Ala Ala Asn Ser Gly Ile
 115 120 125

Tyr

SEQ ID NO: 22

Met Ser Lys Thr Ile Val Leu Ser Val Gly Glu Ala Thr Arg Thr Leu
 1 5 10 15

Thr Glu Ile Gln Ser Thr Ala Asp Arg Gln Ile Phe Glu Glu Lys Val
 20 25 30

Gly Pro Leu Val Gly Arg Leu Arg Leu Thr Ala Ser Leu Arg Gln Asn
 35 40 45

Gly Ala Lys Thr Ala Tyr Arg Val Asn Leu Lys Leu Asp Gln Ala Asp
 50 55 60

Val Val Asp Cys Ser Thr Ser Val Cys Gly Glu Leu Pro Lys Val Arg
 65 70 75 80

Tyr Thr Gln Val Trp Ser His Asp Val Thr Ile Val Ala Asn Ser Thr
 85 90 95

Glu Ala Ser Arg Lys Ser Leu Tyr Asp Leu Thr Lys Ser Leu Val Ala
 100 105 110

Thr Ser Gln Val Glu Asp Leu Val Val Asn Leu Val Pro Leu Gly Arg
 115 120 125

SEQ ID NO: 23

Ala Lys Leu Glu Thr Val Thr Leu Gly Asn Ile Gly Arg Asp Gly Lys
 1 5 10 15

Gln Thr Leu Val Leu Asn Pro Arg Gly Val Asn Pro Thr Asn Gly Val
 20 25 30

-19-

Ala Ser Leu Ser Gln Ala Gly Ala Val Pro Ala Leu Glu Lys Arg Val
 35 40 45

Thr Val Ser Val Ser Gln Pro Ser Arg Asn Arg Lys Asn Tyr Lys Val
 50 55 60

Gln Val Lys Ile Gln Asn Pro Thr Ala Cys Thr Ala Asn Gly Ser Cys
 65 70 75 80

Asp Pro Ser Val Thr Arg Gln Lys Tyr Ala Asp Val Thr Phe Ser Phe
 85 90 95

Thr Gln Tyr Ser Thr Asp Glu Glu Arg Ala Phe Val Arg Thr Glu Leu
 100 105 110

Ala Ala Leu Leu Ala Ser Pro Leu Leu Ile Asp Ala Ile Asp Gln Leu
 115 120 125

Asn Pro Ala Tyr
 130

SEQ ID NO: 24

Ala Lys Leu Glu Thr Val Thr Leu Gly Lys Ile Gly Lys Asp Gly Lys
 1 5 10 15

Gln Thr Leu Val Leu Asn Pro Arg Gly Val Asn Pro Thr Asn Gly Val
 20 25 30

Ala Ser Leu Ser Gln Ala Gly Ala Val Pro Ala Leu Glu Lys Arg Val
 35 40 45

Thr Val Ser Val Ser Gln Pro Ser Arg Asn Arg Lys Asn Tyr Lys Val
 50 55 60

Gln Val Lys Ile Gln Asn Pro Thr Ala Cys Thr Ala Asn Gly Ser Cys
 65 70 75 80

Asp Pro Ser Val Thr Arg Gln Lys Tyr Ala Asp Val Thr Phe Ser Phe
 85 90 95

Thr Gln Tyr Ser Thr Asp Glu Glu Arg Ala Phe Val Arg Thr Glu Leu
 100 105 110

-20-

Ala Ala Leu Leu Ala Ser Pro Leu Leu Ile Asp Ala Ile Asp Gln Leu
 115 120 125

Asn Pro Ala Tyr
 130

SEQ ID NO: 25

Ala Arg Leu Glu Thr Val Thr Leu Gly Asn Ile Gly Arg Asp Gly Lys
 1 5 10 15

Gln Thr Leu Val Leu Asn Pro Arg Gly Val Asn Pro Thr Asn Gly Val
 20 25 30

Ala Ser Leu Ser Gln Ala Gly Ala Val Pro Ala Leu Glu Lys Arg Val
 35 40 45

Thr Val Ser Val Ser Gln Pro Ser Arg Asn Arg Lys Asn Tyr Lys Val
 50 55 60

Gln Val Lys Ile Gln Asn Pro Thr Ala Cys Thr Ala Asn Gly Ser Cys
 65 70 75 80

Asp Pro Ser Val Thr Arg Gln Lys Tyr Ala Asp Val Thr Phe Ser Phe
 85 90 95

Thr Gln Tyr Ser Thr Asp Glu Glu Arg Ala Phe Val Arg Thr Glu Leu
 100 105 110

Ala Ala Leu Leu Ala Ser Pro Leu Leu Ile Asp Ala Ile Asp Gln Leu
 115 120 125

Asn Pro Ala Tyr
 130

SEQ ID NO: 26

Ala Lys Leu Glu Thr Val Thr Leu Gly Asn Ile Gly Lys Asp Gly Arg
 1 5 10 15

Gln Thr Leu Val Leu Asn Pro Arg Gly Val Asn Pro Thr Asn Gly Val
 20 25 30

-21-

Ala Ser Leu Ser Gln Ala Gly Ala Val Pro Ala Leu Glu Lys Arg Val
35 40 45

Thr Val Ser Val Ser Gln Pro Ser Arg Asn Arg Lys Asn Tyr Lys Val
50 55 60

Gln Val Lys Ile Gln Asn Pro Thr Ala Cys Thr Ala Asn Gly Ser Cys
65 70 75 80

Asp Pro Ser Val Thr Arg Gln Lys Tyr Ala Asp Val Thr Phe Ser Phe
85 90 95

Thr Gln Tyr Ser Thr Asp Glu Glu Arg Ala Phe Val Arg Thr Glu Leu
100 105 110

Ala Ala Leu Leu Ala Ser Pro Leu Leu Ile Asp Ala Ile Asp Gln Leu
115 120 125

Asn Pro Ala Tyr
130

SEQ ID NO: 27

Ala Arg Leu Glu Thr Val Thr Leu Gly Asn Ile Gly Lys Asp Gly Arg
1 5 10 15

Gln Thr Leu Val Leu Asn Pro Arg Gly Val Asn Pro Thr Asn Gly Val
20 25 30

Ala Ser Leu Ser Gln Ala Gly Ala Val Pro Ala Leu Glu Lys Arg Val
35 40 45

Thr Val Ser Val Ser Gln Pro Ser Arg Asn Arg Lys Asn Tyr Lys Val
50 55 60

Gln Val Lys Ile Gln Asn Pro Thr Ala Cys Thr Ala Asn Gly Ser Cys
65 70 75 80

Asp Pro Ser Val Thr Arg Gln Lys Tyr Ala Asp Val Thr Phe Ser Phe
85 90 95

-22-

Thr Gln Tyr Ser Thr Asp Glu Glu Arg Ala Phe Val Arg Thr Glu Leu
 100 105 110

Ala Ala Leu Leu Ala Ser Pro Leu Leu Ile Asp Ala Ile Asp Gln Leu
 115 120 125

Asn Pro Ala Tyr
 130

SEQ ID NO: 28

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
 1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
 20 25 30

Thr Ala Ser Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
 35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
 50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Asn Asn Leu Glu Asp Pro Ala
 65 70 75 80

Ser Arg Asp Leu Val Val Asn Tyr Val Asn Thr Asn Met Gly Leu Lys
 85 90 95

Ile Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
 100 105 110

Glu Thr Val Leu Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
 115 120 125

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
 130 135 140

Glu Thr Thr Val Val Arg Arg Arg Asp Arg Gly Arg Ser Pro Arg Arg
 145 150 155 160

Arg Thr Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg
 165 170 175

-23-

Arg Ser Gln Ser Arg Glu Ser Gln Cys
 180 185

SEQ ID NO: 29

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
 1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
 20 25 30

Thr Ala Ser Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
 35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
 50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Ile
 65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
 85 90 95

Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
 100 105 110

Glu Thr Val Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
 115 120 125

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
 130 135 140

Glu Thr Thr Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
 145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Ser
 165 170 175

Gln Ser Arg Gly Ser Gln Cys
 180

-24-

SEQ ID NO: 30

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
20 25 30

Thr Ala Ser Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Thr
65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
85 90 95

Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
100 105 110

Glu Thr Val Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
115 120 125

Pro Pro Ala Tyr Arg Pro Thr Asn Ala Pro Ile Leu Ser Thr Leu Pro
130 135 140

Glu Thr Cys Val Ile Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser
165 170 175

Gln Ser Arg Gly Ser Gln Cys
180

SEQ ID NO: 31

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
1 5 10 15

-25-

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
85 90 95

Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Ile Ser Arg Asp
100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
195 200 205

Glu Ser Gln Cys
210

SEQ ID NO: 32

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
1 5 10 15

-26-

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Asn Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Ile Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

SEQ ID NO: 33

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
 1 5 10 15

-27-

Ser Phe Leu Pro Thr Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
20 25 30

Thr Ala Ser Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala
65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
85 90 95

Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
100 105 110

Glu Thr Val Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
115 120 125

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
130 135 140

Glu Thr Cys Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser
165 170 175

Gln Ser Arg Glu Ser Gln Cys
180

SEQ ID NO: 34

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
35 40 45

-28-

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp Leu Met Thr
85 90 95

Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Val Ser Arg Asp
100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Val Gly Leu Lys Phe Arg Gln
115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
195 200 205

Glu Ser Gln Cys
210

SEQ ID NO: 35

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Asp Met Asp Ile
20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
35 40 45

-29-

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp Leu Met Thr
85 90 95

Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Val Ser Arg Asp
100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Val Gly Leu Lys Phe Arg Gln
115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
195 200 205

Glu Ser Gln Cys
210

SEQ ID NO: 36

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
35 40 45

-30-

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro Gln
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Ile Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

SEQ ID NO: 37

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

-31-

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Lys Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Gly Ser Gln Cys
 210

SEQ ID NO: 38

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
 1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
 20 25 30

Thr Ala Ser Ala Leu Phe Arg Asp Ala Leu Glu Ser Pro Glu His Cys
 35 40 45

-32-

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Ala
65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
85 90 95

Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
100 105 110

Asp Thr Val Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
115 120 125

Pro Pro Ala Tyr Arg Pro Ser Asn Ala Pro Ile Leu Ser Thr Leu Pro
130 135 140

Glu Thr Cys Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser
165 170 175

Gln Ser Arg Glu Ser Gln Cys
180

SEQ ID NO: 39

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
20 25 30

Thr Ala Ser Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
50 55 60

-33-

Leu Met Thr Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala
65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
85 90 95

Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
100 105 110

Glu Thr Val Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
115 120 125

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
130 135 140

Glu Thr Thr Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Ser
165 170 175

Gln Ser Arg Glu Ser Gln Cys
180

SEQ ID NO: 40

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
65 70 75 80

His Thr Ala Leu Arg His Ala Ile Leu Cys Trp Gly Asp Leu Arg Thr
85 90 95

-34-

Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Ile Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

SEQ ID NO: 41

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Asp Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Phe Arg Asp Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
 85 90 95

-35-

Leu Ala Thr Trp Val Gly Ala Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Gln Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Cys
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

SEQ ID NO: 42

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
 1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
 20 25 30

Thr Ala Ser Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
 35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
 50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala
 65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
 85 90 95

-36-

Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
 100 105 110

Glu Thr Val Leu Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
 115 120 125

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
 130 135 140

Glu Thr Thr Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
 145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser
 165 170 175

Gln Ser Arg Glu Ser Gln Cys
 180

SEQ ID NO: 43

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp Leu Met Ser
 85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ile Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

-37-

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

SEQ ID NO: 44

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
 1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
 20 25 30

Thr Ala Ser Ala Leu Tyr Arg Asp Ala Leu Glu Ser Pro Glu His Cys
 35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
 50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala
 65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
 85 90 95

Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
 100 105 110

Glu Thr Val Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
 115 120 125

-38-

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
 130 135 140

Glu Thr Thr Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
 145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser
 165 170 175

Gln Ser Arg Glu Ser Gln Cys
 180

SEQ ID NO: 45

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
 1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
 20 25 30

Thr Ala Ser Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
 35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp
 50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala
 65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
 85 90 95

Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
 100 105 110

Glu Thr Val Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
 115 120 125

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
 130 135 140

Glu Thr Thr Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
 145 150 155 160

-39-

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser
 165 170 175

Gln Ser Arg Glu Ser Gln Cys
 180

SEQ ID NO: 46

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
 1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
 20 25 30

Thr Ala Ser Ala Leu Tyr Arg Asp Ala Leu Glu Ser Pro Glu His Cys
 35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
 50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Ala Asn Leu Glu Asp Pro Ala
 65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
 85 90 95

Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
 100 105 110

Glu Thr Val Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
 115 120 125

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
 130 135 140

Glu Thr Thr Val Val Arg Arg Arg Gly Arg Thr Pro Arg Arg Arg Thr
 145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser
 165 170 175

Gln Ser Arg Glu Ser Gln Cys
 180

-40-

SEQ ID NO: 47

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Asp Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

-41-

SEQ ID NO: 48

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

-42-

SEQ ID NO: 49

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Thr Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ala Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

-43-

SEQ ID NO: 50

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Phe Glu Cys Ser Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Ile Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

-44-

SEQ ID NO: 51 (Xaa at 28 can be any amino acid)

Met	Gln	Leu	Phe	His	Leu	Cys	Leu	Ile	Ile	Ser	Cys	Ser	Cys	Pro	Thr
1				5					10					15	

Val	Gln	Ala	Ser	Lys	Leu	Cys	Leu	Gly	Trp	Leu	Xaa	Asp	Met	Asp	Ile
			20					25					30		

Asp	Pro	Tyr	Lys	Glu	Phe	Gly	Ala	Thr	Val	Glu	Leu	Leu	Ser	Phe	Leu
		35					40						45		

Pro	Ser	Asp	Phe	Phe	Pro	Ser	Val	Arg	Asp	Leu	Leu	Asp	Thr	Ala	Ser
	50					55					60				

Ala	Leu	Tyr	Arg	Glu	Ala	Leu	Glu	Ser	Pro	Glu	His	Cys	Ser	Pro	His
65					70					75					80

His	Thr	Ala	Leu	Arg	Gln	Ala	Ile	Leu	Cys	Trp	Gly	Asp	Leu	Ile	Thr
			85						90					95	

Leu	Ser	Thr	Trp	Val	Gly	Gly	Asn	Leu	Glu	Asp	Pro	Thr	Ser	Arg	Asp
			100					105						110	

Leu	Val	Val	Ser	Tyr	Val	Asn	Thr	Asn	Met	Gly	Leu	Lys	Phe	Arg	Gln
		115					120					125			

Leu	Leu	Trp	Phe	His	Ile	Ser	Cys	Leu	Thr	Phe	Gly	Arg	Glu	Thr	Val
	130					135					140				

Ile	Glu	Tyr	Leu	Val	Ser	Phe	Gly	Val	Trp	Ile	Arg	Thr	Pro	Pro	Ala
145					150					155					160

Tyr	Arg	Pro	Pro	Asn	Ala	Pro	Ile	Leu	Ser	Thr	Leu	Pro	Glu	Thr	Thr
				165					170					175	

Val	Val	Arg	Arg	Arg	Gly	Arg	Ser	Pro	Arg	Arg	Arg	Thr	Pro	Ser	Pro
			180					185					190		

Arg	Arg	Arg	Arg	Ser	Gln	Ser	Pro	Arg	Arg	Arg	Arg	Thr	Gln	Ser	Arg
		195					200					205			

-45-

Glu Ser Gln Cys
210

SEQ ID NO: 52

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Asn Ala Ser
50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
195 200 205

-46-

Glu Ser Gln Cys
210

SEQ ID NO: 53

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
115 120 125

Leu Leu Trp Phe His Ile Cys Cys Leu Thr Phe Gly Arg Glu Thr Val
130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
180 185 190

-47-

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

SEQ ID NO: 54

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

-48-

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Pro Gln Cys
 210

SEQ ID NO: 55

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Ser Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

-49-

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

SEQ ID NO: 56

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Leu Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

-50-

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

SEQ ID NO: 57

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Lys Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

-51-

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

SEQ ID NO: 58

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ala
 50 55 60

Ala Leu Tyr Arg Asp Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Thr Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Leu Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

-52-

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

SEQ ID NO: 59

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Ser Met Glu Leu Leu
 1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Tyr Pro Ser Val Arg Asp Leu Leu Asp
 20 25 30

Thr Ala Ser Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
 35 40 45

Thr Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
 50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Gly Asn Leu Gln Asp Pro Thr
 65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
 85 90 95

Phe Arg Gln Leu Leu Trp Phe His Val Ser Cys Leu Thr Phe Gly Arg
 100 105 110

Glu Thr Val Val Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
 115 120 125

Pro Gln Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
 130 135 140

Glu Thr Cys Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
 145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser
 165 170 175

-53-

Gln Ser Arg Glu Ser Gln Cys
180

SEQ ID NO: 60

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
20 25 30

Thr Ala Ser Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
35 40 45

Ser Pro His His Thr Ala Leu Arg His Val Phe Leu Cys Trp Gly Asp
50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Thr
65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
85 90 95

Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
100 105 110

Glu Thr Val Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
115 120 125

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
130 135 140

Glu Thr Thr Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser
165 170 175

Gln Ser Arg Glu Ser Gln Cys
180

-54-

SEQ ID NO: 61

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp Leu Thr Thr
 85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

-55-

SEQ ID NO: 62

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15
 Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30
 Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45
 Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60
 Ala Leu Tyr Arg Asp Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80
 His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
 85 90 95
 Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110
 Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125
 Leu Leu Trp Phe His Ile Ser Cys Leu Ile Phe Gly Arg Glu Thr Val
 130 135 140
 Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160
 Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175
 Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190
 Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205
 Glu Ser Gln Cys
 210

-56-

SEQ ID NO: 63

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
 1 5 10 15
 Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
 20 25 30
 Thr Ala Ser Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
 35 40 45
 Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp
 50 55 60
 Leu Met Thr Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Val
 65 70 75 80
 Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Val Gly Leu Lys
 85 90 95
 Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
 100 105 110
 Glu Thr Val Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
 115 120 125
 Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
 130 135 140
 Glu Thr Thr Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
 145 150 155 160
 Pro Ser Pro Ala Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser
 165 170 175
 Gln Ser Arg Glu Ser Gln Cys
 180

SEQ ID NO: 64

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

-57-

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30
 Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45
 Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60
 Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80
 His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp Leu Met Asn
 85 90 95
 Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Val Ser Arg Asp
 100 105 110
 Leu Val Val Gly Tyr Val Asn Thr Thr Val Gly Leu Lys Phe Arg Gln
 115 120 125
 Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140
 Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160
 Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175
 Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190
 Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205
 Glu Ser Gln Cys
 210

SEQ ID NO: 65

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
 1 5 10 15

-58-

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
 20 25 30

Thr Ala Ser Ala Leu Tyr Arg Asp Ala Leu Glu Ser Pro Glu His Cys
 35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp
 50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala
 65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
 85 90 95

Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
 100 105 110

Glu Thr Val Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
 115 120 125

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
 130 135 140

Glu Thr Thr Val Val Arg Arg Arg Gly Arg Thr Pro Arg Arg Arg Thr
 145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser
 165 170 175

Gln Ser Arg Glu Ser Gln Cys
 180

SEQ ID NO: 66

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
 1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
 20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

-59-

Pro Ser Asp Phe Phe Pro Ser Val Arg Ala Leu Leu Asp Thr Ala Ser
50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
115 120 125

Ile Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
195 200 205

Glu Ser Gln Cys
210

SEQ ID NO: 67

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
35 40 45

-60-

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp Leu Met Thr
85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Thr Arg Asp
100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Val Gly Leu Lys Phe Arg Gln
115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
195 200 205

Glu Ser Gln Cys
210

SEQ ID NO: 68

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
35 40 45

-61-

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
65 70 75 80

His Thr Ala Leu Arg Gln Arg Ile Leu Cys Trp Gly Glu Leu Met Thr
85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Thr Arg Ser Gln Ser Arg
195 200 205

Glu Ser Gln Cys
210

SEQ ID NO: 69

Met Gln Leu Phe His Leu Cys Leu Val Ile Ser Cys Ser Cys Pro Thr
1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
20 25 30

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
35 40 45

-62-

Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ala
50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu Leu Met Thr
85 90 95

Leu Ala Thr Trp Val Gly Asn Asn Leu Glu Asp Pro Ala Ser Arg Asp
100 105 110

Leu Val Val Asn Tyr Val Asn Thr Asn Met Gly Leu Lys Ile Arg Gln
115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
130 135 140

Leu Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
195 200 205

Glu Ser Gln Cys
210

SEQ ID NO: 70

Met Gln Leu Phe His Leu Cys Leu Ile Ile Ser Cys Ser Cys Pro Thr
1 5 10 15

Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Gly Met Asp Ile
20 25 30

-63-

Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu Ser Phe Leu
 35 40 45

Pro Ser Ala Phe Phe Pro Ser Val Arg Asp Leu Leu Asp Thr Ala Ser
 50 55 60

Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys Ser Pro His
 65 70 75 80

His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp Leu Met Thr
 85 90 95

Leu Ala Thr Trp Val Gly Val Asn Leu Glu Asp Pro Ala Ser Arg Asp
 100 105 110

Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys Phe Arg Gln
 115 120 125

Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg Glu Thr Val
 130 135 140

Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Pro Ala
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu Thr Thr
 165 170 175

Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro
 180 185 190

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Arg
 195 200 205

Glu Ser Gln Cys
 210

SEQ ID NO: 71

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
 1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
 20 25 30

-64-

Thr Ala Ala Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Asn Asn Leu Glu Asp Pro Ala
65 70 75 80

Ser Arg Asp Leu Val Val Asn Tyr Val Asn Thr Asn Met Gly Leu Lys
85 90 95

Ile Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
100 105 110

Glu Thr Val Leu Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
115 120 125

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
130 135 140

Glu Thr Thr Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Ser
165 170 175

Gln Ser Arg Glu Ser Gln Cys
180

SEQ ID NO: 72

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
20 25 30

Thr Ala Ser Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
50 55 60

-65-

Leu Met Thr Leu Ala Thr Trp Val Gly Gly Asn Leu Glu Asp Pro Ile
65 70 75 80

Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Met Gly Leu Lys
85 90 95

Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
100 105 110

Glu Thr Val Ile Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
115 120 125

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
130 135 140

Glu Thr Cys Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr
145 150 155 160

Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser
165 170 175

Gln Ser Arg Gly Ser Gln Cys
180

SEQ ID NO: 73

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ser Ser Tyr Gln Leu Leu
1 5 10 15

Asn Phe Leu Pro Leu Asp Phe Phe Pro Asp Leu Asn Ala Leu Val Asp
20 25 30

Thr Ala Thr Ala Leu Tyr Glu Glu Glu Leu Thr Gly Arg Glu His Cys
35 40 45

Ser Pro His His Thr Ala Ile Arg Gln Ala Leu Val Cys Trp Asp Glu
50 55 60

Leu Thr Lys Leu Ile Ala Trp Met Ser Ser Asn Ile Thr Ser Glu Gln
65 70 75 80

-66-

Val Arg Thr Ile Ile Val Asn His Val Asn Asp Thr Trp Gly Leu Lys
85 90 95

Val Arg Gln Ser Leu Trp Phe His Leu Ser Cys Leu Thr Phe Gly Gln
100 105 110

His Thr Val Gln Glu Phe Leu Val Ser Phe Gly Val Trp Ile Arg Thr
115 120 125

Pro Ala Pro Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
130 135 140

Glu His Thr Val Ile Arg Arg Arg Gly Gly Ala Arg Ala Ser Arg Ser
145 150 155 160

Pro Arg Arg Arg Thr Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro
165 170 175

Arg Arg Arg Arg Ser Gln Ser Pro Ser Thr Asn Cys
180 185

SEQ ID NO: 74

Met Tyr Leu Phe His Leu Cys Leu Val Phe Ala Cys Val Pro Cys Pro
1 5 10 15

Thr Val Gln Ala Ser Lys Leu Cys Leu Gly Trp Leu Trp Asp Met Asp
20 25 30

Ile Asp Pro Tyr Lys Glu Phe Gly Ser Ser Tyr Gln Leu Leu Asn Phe
35 40 45

Leu Pro Leu Asp Phe Phe Pro Asp Leu Asn Ala Leu Val Asp Thr Ala
50 55 60

Ala Ala Leu Tyr Glu Glu Glu Leu Thr Gly Arg Glu His Cys Ser Pro
65 70 75 80

His His Thr Ala Ile Arg Gln Ala Leu Val Cys Trp Glu Glu Leu Thr
85 90 95

Arg Leu Ile Thr Trp Met Ser Glu Asn Thr Thr Glu Glu Val Arg Arg
100 105 110

-67-

Ile Ile Val Asp His Val Asn Asn Thr Trp Gly Leu Lys Val Arg Gln
 115 120 125

Thr Leu Trp Phe His Leu Ser Cys Leu Thr Phe Gly Gln His Thr Val
 130 135 140

Gln Glu Phe Leu Val Ser Phe Gly Val Trp Ile Arg Thr Pro Ala Pro
 145 150 155 160

Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro Glu His Thr
 165 170 175

Val Ile Arg Arg Arg Gly Gly Ser Arg Ala Ala Arg Ser Pro Arg Arg
 180 185 190

Arg Thr Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg
 195 200 205

Arg Ser Gln Ser Pro Ala Ser Asn Cys
 210 215

SEQ ID NO: 75

Met Asp Val Asn Ala Ser Arg Ala Leu Ala Asn Val Tyr Asp Leu Pro
 1 5 10 15

Asp Asp Phe Phe Pro Lys Ile Glu Asp Leu Val Arg Asp Ala Lys Asp
 20 25 30

Ala Leu Glu Pro Tyr Trp Lys Ser Asp Ser Ile Lys Lys His Val Leu
 35 40 45

Ile Ala Thr His Phe Val Asp Leu Ile Glu Asp Phe Trp Gln Thr Thr
 50 55 60

Gln Gly Met His Glu Ile Ala Glu Ala Ile Arg Ala Val Ile Pro Pro
 65 70 75 80

Thr Thr Ala Pro Val Pro Ser Gly Tyr Leu Ile Gln His Asp Glu Ala
 85 90 95

Glu Glu Ile Pro Leu Gly Asp Leu Phe Lys Glu Gln Glu Glu Arg Ile
 100 105 110

-68-

Val Ser Phe Gln Pro Asp Tyr Pro Ile Thr Ala Arg Ile His Ala His
115 120 125

Leu Lys Ala Tyr Ala Lys Ile Asn Glu Glu Ser Leu Asp Arg Ala Arg
130 135 140

Arg Leu Leu Trp Trp His Tyr Asn Cys Leu Leu Trp Gly Glu Ala Thr
145 150 155 160

Val Thr Asn Tyr Ile Ser Arg Leu Arg Thr Trp Leu Ser Thr Pro Glu
165 170 175

Lys Tyr Arg Gly Arg Asp Ala Pro Thr Ile Glu Ala Ile Thr Arg Pro
180 185 190

Ile Gln Val Ala Gln Gly Gly Arg Lys Thr Ser Thr Ala Thr Arg Lys
195 200 205

Pro Arg Gly Leu Glu Pro Arg Arg Arg Lys Val Lys Thr Thr Val Val
210 215 220

Tyr Gly Arg Arg Arg Ser Lys Ser Arg Glu Arg Arg Ala Ser Ser Pro
225 230 235 240

Gln Arg Ala Gly Ser Pro Leu Pro Arg Ser Ser Ser Ser His His Arg
245 250 255

Ser Pro Ser Pro Arg Lys
260

SEQ ID NO: 76

Met Trp Asp Leu Arg Leu His Pro Ser Pro Phe Gly Ala Ala Cys Gln
1 5 10 15

Gly Ile Phe Thr Ser Ser Leu Leu Leu Phe Leu Val Thr Val Pro Leu
20 25 30

Val Cys Thr Ile Val Tyr Asp Ser Cys Leu Cys Met Asp Ile Asn Ala
35 40 45

-69-

Ser Arg Ala Leu Ala Asn Val Tyr Asp Leu Pro Asp Asp Phe Phe Pro
50 55 60

Lys Ile Asp Asp Leu Val Arg Asp Ala Lys Asp Ala Leu Glu Pro Tyr
65 70 75 80

Trp Arg Asn Asp Ser Ile Lys Lys His Val Leu Ile Ala Thr His Phe
85 90 95

Val Asp Leu Ile Glu Asp Phe Trp Gln Thr Thr Gln Gly Met His Glu
100 105 110

Ile Ala Glu Ala Leu Arg Ala Ile Ile Pro Ala Thr Thr Ala Pro Val
115 120 125

Pro Gln Gly Phe Leu Val Gln His Glu Glu Ala Glu Glu Ile Pro Leu
130 135 140

Gly Glu Leu Phe Arg Tyr Gln Glu Glu Arg Leu Thr Asn Phe Gln Pro
145 150 155 160

Asp Tyr Pro Val Thr Ala Arg Ile His Ala His Leu Lys Ala Tyr Ala
165 170 175

Lys Ile Asn Glu Glu Ser Leu Asp Arg Ala Arg Arg Leu Leu Trp Trp
180 185 190

His Tyr Asn Cys Leu Leu Trp Gly Glu Pro Asn Val Thr Asn Tyr Ile
195 200 205

Ser Arg Leu Arg Thr Trp Leu Ser Thr Pro Glu Lys Tyr Arg Gly Lys
210 215 220

Asp Ala Pro Thr Ile Glu Ala Ile Thr Arg Pro Ile Gln Val Ala Gln
225 230 235 240

Gly Gly Arg Asn Lys Thr Gln Gly Val Arg Lys Ser Arg Gly Leu Glu
245 250 255

Pro Arg Arg Arg Arg Val Lys Thr Thr Ile Val Tyr Gly Arg Arg Arg
260 265 270

Ser Lys Ser Arg Glu Arg Arg Ala Pro Thr Pro Gln Arg Ala Gly Ser
275 280 285

Pro Leu Pro Arg Thr Ser Arg Asp His His Arg Ser Pro Ser Pro Arg
290 295 300

Glu
305

SEQ ID NO: 77

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
20 25 30

Thr Ala Ser Ala Leu Tyr Arg Glu Ala Leu Glu Ser Pro Glu His Cys
35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Glu
50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Asn Asn Leu Glu Asp Pro Ala
65 70 75 80

Ser Arg Asp Leu Val Val Asn Tyr Val Asn Thr Asn Met Gly Leu Lys
85 90 95

Ile Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr Phe Gly Arg
100 105 110

Glu Thr Val Leu Glu Tyr Leu Val Ser Phe Gly Val Trp Ile Arg Thr
115 120 125

Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser Thr Leu Pro
130 135 140

Glu Thr Thr Val Val Arg Arg Arg Asp Arg Gly Arg Ser Pro Arg Arg
145 150 155 160

Arg Thr Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg
165 170 175

Arg Ser Gln Ser Arg Glu Ser Gln Cys
180 185

SEQ ID NO: 78

Met Asp Ile Asp Pro Tyr Lys Glu Phe Gly Ala Thr Val Glu Leu Leu
1 5 10 15

Ser Phe Leu Pro Ser Asp Phe Phe Pro Ser Val Arg Asp Leu Leu Asp
20 25 30

Thr Ala Ala Ala Leu Tyr Arg Asp Ala Leu Glu Ser Pro Glu His Cys
35 40 45

Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp Gly Asp
50 55 60

Leu Met Thr Leu Ala Thr Trp Val Gly Thr Asn Leu Glu Asp Gly Gly
65 70 75 80

Lys Gly Gly Ser Arg Asp Leu Val Val Ser Tyr Val Asn Thr Asn Val
85 90 95

Gly Leu Lys Phe Arg Gln Leu Leu Trp Phe His Ile Ser Cys Leu Thr
100 105 110

Phe Gly Arg Glu Thr Val Leu Glu Tyr Leu Val Ser Phe Gly Val Trp
115 120 125

Ile Arg Thr Pro Pro Ala Tyr Arg Pro Pro Asn Ala Pro Ile Leu Ser
130 135 140

Thr Leu Pro Glu Thr Thr Val Val
145 150

SEQ ID NO: 79

RANKL_human TrEMBL:O14788 full-length

MRRASRDYTKYLRGSEEMGGGPGAPHEGPLHAPPPAPHPQPPAASRSMFVALLGLGLGQV
VCSVALFFYFRAQMDPNRISEDGTHCIYRILRLHENADFQDTTLESQDTKLIPDSCRRIK
QAFQGAVQKELQHIVGSQHIRAEKAMVDGSWLDLAKRSKLEAQPPFAHLTINATDIPSGSH
KVSLSWYHDRGWAKISNMTFSNGKLIVNQDGFYYLYANICFRHHETSGDLATEYLQLMV
YVTKTSIKIPSSHTLMKGGSTKYWSGNSEFHFYSINVGGFFKLRSGEISIEVSNPSLLD
PDQDATYFGAFKVRDID

SEQ ID NO: 80

RANKL_human: TrEMBL:O14788 spliced isoform 1

MFVALLGLGLGQVCSVALFFYFRAQMDPNRISEDGTHCIYRILRLHENADFQDTTLESQ
DTKLIPDSCRRIKQAFQGAVQKELQHIVGSQHIRAEKAMVDGSWLDLAKRSKLEAQPPFAH
LTINATDIPSGSHKVSLSWYHDRGWAKISNMTFSNGKLIVNQDGFYYLYANICFRHHET
SGDLATEYLQLMVYVTKTSIKIPSSHTLMKGGSTKYWSGNSEFHFYSINVGGFFKLRSGE
EISIEVSNPSLLDPDQDATYFGAFKVRDID

SEQ ID NO: 81

RANKL_human: TrEMBL:O14788 spliced isoform 2

MDPNRISEDG THCIYRILRL HENADFQDTT LESQDTKLIP DSCRRIKQAF QGAVQKELQH
IVGSQHIRAE KAMVDGSWLD LAKRSKLEAQ PFAHLTINAT DIPSGSHKVS LSSWYHDRGW
AKISNMTFSN GKLVNQDGF YYLYANICFR HHETSGDLAT EYLQLMVYVT KTSIKIPSSH
TLMKGGSTKY WSGNSEFHFY SINVGFFKL RSGEISIEV SNPSLLDPDQ DATYFGAFKV
RDID

SEQ ID NO: 82

RANKL_human: TrEMBL:O14788: extracellular region

YFRAQMDPNRIS EDGTHCIYRI LRLHENADFQ DTTLESQDTK LIPDSCRRRIK QAFQGA VQKE LQHIVGSQHI
RAEKAMVDGS WLDLAKRSKL EAQPF AHLTI NATDIPSGSH KVSLS SWYHD RGWAKISNMT FSNGKLIVNQ
DGFYYLYANI CFRHHETSGD LATEYLQLMV YVTKTSIKIP SSHTLMKGG S TKYWSGNSEF HFYSINVGGF
FKLRS GEEIS IEVSNPSLLD PDQDATYFGA FKVRDID

SEQ ID NO: 83

RANKL_human: TrEMBL:O14788: region cleaved by a TACE-like protease

IRAEKAMVDG SWLDLAKRSK LEAQPF AHLT INATDIPSGS HKVSLSSWYH DRGWAKISNM
TFSNGKLIVN QDGFYYLYAN ICFRHHETSG DLATEYLQLM VYVTKTSIKI PSSHTLMKGG
STKYWSGNSE FHFYSINVGG FFKLRS GEEI SIEVSNPSLL DPDQDATYFG AFKVRDID

SEQ ID NO: 84

RANKL_human: TrEMBL:O14788: TNFa-homolog region

KLEAQPF AHLTI NATDIPSGSH KVSLS SWYHD RGWAKISNMT FSNGKLIVNQ DGFYYLYANI
CFRHHETSGD LATEYLQLMV YVTKTSIKIP SSHTLMKGG S TKYWSGNSEF HFYSINVGGF
FKLRS GEEIS IEVSNPSLLD PDQDATYFGA FKVRDID

SEQ ID NO: 85

RANKL_mouse: TrEMBL:O35235: extracellular domain

YFRAQMDPNRI SEDSTHCFYR ILRLHENAGL QDSTLESED TLPDSCRRMKQ AFQGA VQKEL QHIVGPQRFS
GAPAMMEGSW LDVAQRGKPE AQPF AHLTIN AASIPSGSHK VTLSSWYHDR GWAKISNMTL SNGKL RVNQD
GFYYLYANIC FRHHETSGSV PTDYLQLMVY VVKTSIKIPS SHNLMKGGST KNWSGNSEFH FYSINVGGFF
KLRAGEEISI QVSNPSLLDP DQDATYFGAF KVQDID

SEQ ID NO: 86

RANKL_mouse spliced isoforms: TrEMBL:Q9JJK8

MKQAFQGA VQ KELQHIVGPQ RFSGAPAMME GSWLDVAQRG KPEAQPF AHL TINAASIPSG
SHKVTLSSWY HDRGWAKISN MTL SNGKL RV NQDGFYYLYA NICFRHHETS GSVPTDYLQL
MVYVVKTSIK IPSSHNL MKG GSTKNWSGNS EFHFYSINVG GFFKL RAGEE ISIQVSNPSL
LDPDQDATYF GAFKVQDID

SEQ ID NO: 87

RANKL peptides AA`` loop
NATDIPSGSHKVSLS SWYHDRGWA

SEQ ID NO: 88

RANKL peptides DE loop
SIKIPSSH

SEQ ID NO: 89

RANKL peptide b-strand D
YLQLMVVTKT

SEQ ID NO: 90

RANKL peptide CD loop
HETSGDLATE

SEQ ID NO: 91

RANKL peptide EF loop
KYWSGNSEPHF

SEQ ID NO: 92

RANKL peptide B strand-B`B loop
TFSNGKLI

SEQ ID NO: 93

RANKL peptide GH loop
DPDQDA

SEQ ID NO: 94

GST-PS-C-RANKL cDNA sequence (mouse RANKL)

1 atgtcccctatactagggttattggaaaattaagggccttgtgcaaccactcgacttcttttgggaatatcttgaa
76 gaaaaatatgaagagcatttgtatgagcgcgatgaaggtgataaatggcgaaacaaaaagtttgaattggggttg
151 gagtttcccaatcttccttattatattgatgggtgatgttaaattaacacagtctatggccatcatacgttatat
a
226 gctgacaagcacacaacatgttggtggttggtccaaaagagcgtgcagagatttcaatgcttgaaggagcgggtttt
g
301 gatattagatacgggtgttttcgagaattgcatatagtaaagactttgaaactctcaaagttgattttcttagcaa
g
376 ctacctgaaatgctgaaaatgttcgaagatcgtttatgtcataaaacatatttaaagtgtgatcatgtaaccca
t
451 cctgacttcatgttgtatgacgctcttgatgttgttttatacatggaccaatgtgcctggatgcgttcccaaa
a
526 ttagtttgttttaaaaaacgtattgaagctatcccacaaattgataagttacttgaaatccagcaagtatatagc
a
601 tggcctttgcagggctggcaagccacgtttggtggtggcgaccatcctccaaaatcggatctggaagttctgtt
c
676 cagGGGCCCCGGGTGCGGCGGTGGCCATCATCACCACCATCACCAGCGCTTCTCAGGAGCTCCAGCTATGATGGA
A
751 GGCTCATGGTTGGATGTGGCCAGCGAGGCAAGCCTGAGGCCAGCCATTTGCACACCTCACCATCAATGCTGC
C
826 AGCATCCCATCGGGTTCCCATAAAGTCACTCTGTCTCTTGGTACCACGATCGAGGCTGGGCCAAGATCTCTAA
C
901 ATGACGTTAAGCAACGGAAAACTAAGGGTTAACCAAGATGGCTTCTATTACCTGTACGCCAACATTTGCTTTTCG
G
976 CATCATGAAACATCGGGAAGCGTACCTACAGACTATCTTCAGCTGATGGTGTATGTCGTTAAAACCAGCATCAA
A

1051 ATCCCAAGTTCTCATAACCTGATGAAAGGAGGGAGCACGAAAACTGGTCGGGCAATTCTGAATTCCACTTTT
AT
1126 TCCATAAATGTTGGGGGATTTTCAAGCTCCGAGCTGGTGAAGAAATTAGCATTACAGGTGTCCAACCCCTTCCC
TG
1201 CTGGATCCGGATCAAGATGCGACGTACTTTGGGGCTTTCAAAGTTCAGGACATAGACTAACTCGAGCGG

SEQ ID NO: 95

GST-PS-C-RANKL protein sequence (mouse RANKL)

1 M S P I L G Y W K I K G L V Q P T R L L L E Y L
26 E K Y E E H L Y E R D E G D K W R N K K F E L G L
51 E F P N L P Y Y I D G D V K L T Q S M A I I R Y I
76 A D K H N M L G G C P K E R A E I S M L E G A V L
101 D I R Y G V S R I A Y S K D F E T L K V D F L S K
126 L P E M L K M F E D R L C H K T Y L N G D H V T H
151 P D F M L Y D A L D V V L Y M D P M C L D A F P K
176 L V C F K K R I E A I P Q I D K Y L K S S K Y I A
201 W P L Q G W Q A T F G G G D H P P K S D L E V L F
226 Q G P G C G G G H H H H H H Q R F S G A P A M M E
251 G S W L D V A Q R G K P E A Q P F A H L T I N A A
276 S I P S G S H K V T L S S W Y H D R G W A K I S N
301 M T L S N G K L R V N Q D G F Y Y L Y A N I C F R
326 H H E T S G S V P T D Y L Q L M V Y V V K T S I K
351 I P S S H N L M K G G S T K N W S G N S E F H F Y
376 S I N V G G F F K L R A G E E I S I Q V S N P S L
401 L D P D Q D A T Y F G A F K V Q D I D *

SEQ ID NO: 96

RANKL-C protein sequence (mouse RANKL)

G P G C G G G H H H H H H Q R F S G A P A M M E
G S W L D V A Q R G K P E A Q P F A H L T I N A A

S I P S G S H K V T L S S W Y H D R G W A K I S N

M T L S N G K L R V N Q D G F Y Y L Y A N I C F R

H H E T S G S V P T D Y L Q L M V Y V V K T S I K

I P S S H N L M K G G S T K N W S G N S E F H F Y

S I N V G G F F K L R A G E E I S I Q V S N P S L

L D P D Q D A T Y F G A F K V Q D I D *

SEQ ID NO: 97

GST-EK-RANKL-C cDNA sequence (mouse RANKL)

```
1   ATGGATCCTATACTAGGTTATTGGAAAAATTAAGGGCCTTGTGCAACCCACTCGACTTCTT
61  TTGGAATATCTTGAAGAAAAATATGAAGAGCATTTGTATGAGCGCGATGAAGGTGATAAA
121 TGGCGAAACAAAAAGTTTGAATTGGGTTTGGAGTTTCCCAATCTTCCTTATTATATTGAT
181 GGTGATGTTAAATTAACACAGTCTATGGCCATCATACGTTATATAGCTGACAAGCACAAC
241 ATGTTGGGTGGTGTGCCAAAAGAGCGTGCAGAGATTTCAATGCTTGAAGGAGCGGTTTTG
301 GATATTAGATACGGTGTTCGAGAATTGCATATAGTAAAGACTTTGAAACTCTCAAAGTT
361 GATTTTCTTAGCAAGCTACCTGAAATGCTGAAAATGTTTGAAGATCGTTTATGTCATAAA
421 ACATATTTAAATGGTGATCATGTAACCCATCCTGACTTCATGTTGTATGACGCTCTTGAT
481 GTTGTTTTATACATGGACCCAATGTGCCTGGATGCGTTCCCAAAATTAGTTTGTMTTAAA
541 AAACGTATTGAAGCTATCCCAAAAATTGATAAGTACTTGAAATCCAGCAAGTATATAGCA
601 TGGCCTTTGCAGGGCTGGCAAGCCACGTTTGGTGGTGGCGACCATCTCCAAAAGCTAGT
661 ATGACTGGTGGACAGCAAATGGGTCGGGATCTGTACGACGATGACGATAAGCTAGCAAAG
721 CCTGAGGCCAGCCATTTCACACCTCACCATCAATGCTGCCAGCATCCCATCGGGTTCC
781 CATAAAGTCACTCTGTCTCTTGGTACCACGATCGAGGCTGGGCCAAGATCTCTAACATG
841 ACGTTAAGCAACGGAATAAGGGTTAAACCAAGATGGCTTCTATTACCTGTACGCCAAC
901 ATTTGCTTTTCGGCATCATGAAACATCGGGAAGCGTACCTACAGACTATCTTCAGCTGATG
961 GTGTATGTCGTTAAACCAGCATCAAAATCCCAAGTTCTCATAACCTGATGAAAGGAGGG
1021 AGCACGAAAAACTGGTTCGGGCAATTCTGAATTCCACTTTTATTCCATAAATGTTGGGGGA
1081 TTTTTCAGCTCCGAGCTGGTGAAGAAATTAGCATTCAGGTGTCCAACCCTTCCCTGCTG
1161 GATCCGGATCAAGATGCGACGTACTTTGGGGCTTTCAAAGTTCAGGACATAGACCTCGAG
1221 GGTGGTGGTGGTGGTTGCGGT
```

SEQ ID NO: 98

GST-EK-RANKL-C protein sequence (mouse RANKL)

```
1   MDPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELGLEFPNLPYYID
61  GDVKLTQSMAIIRYIADKHNMLGGCPKERAIEISMLEGAVLDIRYGVSR IAYSKDFETLKV
121 DFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALDVVLYMDPMCLDAFPKLVCFK
181 KRIEAIQIDKYLKSSKYIAWPLQGWQATFGGGDHPKASMTGGQQMGRDLYDDDDKLAK
241 PEAQPF AHLTINAASIPSGSHKVTLSWYHDRGWAKISNMTLSNGKLRVNQDGFYYLYAN
301 ICFRHHETSGSVPTDYLQLMVYVVKTSIKIPSSHNLMKGGSTKNWSGNSEFHFYSINVG
361 FFKLRAGEEISIQVSNPSLLDPDQDATYFGAFKVQDIDLEGGGGGCG
```

SEQ ID NO: 99

RANKL-C protein sequence (mouse RANKL)

LAKPEAQPF AHLTINAASIPSGSHKVTLSWYHDRGWAKISNMTLSNGKLRVNQDGFYYLYAN
ICFRHHETSGSVPTDYLQLMVYVVKTSIKIPSSHNLMKGGSTKNWSGNSEFHFYSINVGG
FFKLRAGEEISIQVSNPSLLDPDQDATYFGAFKVQDIDLEGGGGGCG

SEQ ID NO: 100

Human-C-RANKL corresponding to mouse C-RANKL 96

QHIRAEKAMVDGSWDLAKRSKLEAQPF AHLTINATDIPSGSHKVSLSWYHDRGWAKISNMTFSNGKLIVNQDGFYY
LYANICFRHHETSGDLATEYLQLMVYVVKTSIKIPSSHTLMKGGSTKYWSGNSEFHFYSINVGGFFKLRSGEESIEV
SNPSLLDPDQDATYFGAFKVRDID

SEQ ID NO: 101

Human-C-RANKL corresponding to mouse RANKL-C 99

LEAQPF AHLTINATDIPSGSHKVSLSWYHDRGWAKISNMTFSNGKLIVNQDGFYYLYANICFRHHETSGDLATEYLQ
LMVYVVKTSIKIPSSHTLMKGGSTKYWSGNSEFHFYSINVGGFFKLRSGEESIEVSNPSLLDPDQDATYFGAFKVRD
ID

SEQ ID NO: 102

Human-C-RANKL with linker corresponding to mouse C-RANKL 96

CGGGGGGQHIRAEKAMVDGSWDLAKRSKLEAQPF AHLTINATDIPSGSHKVSLSWYHDRGWAKISNMTFSNGKLIV
NQDGFYYLYANICFRHHETSGDLATEYLQLMVYVVKTSIKIPSSHTLMKGGSTKYWSGNSEFHFYSINVGGFFKLRS
GEESIEVSNPSLLDPDQDATYFGAFKVRDID

SEQ ID NO: 103

Human-C-RANKL with linker corresponding to mouse RANKL-C 99

LEAQPF AHLTINATDIPSGSHKVSLSWYHDRGWAKISNMTFSNGKLIVNQDGFYYLYANICFRHHETSGDLATEYLQ
LMVYVVKTSIKIPSSHTLMKGGSTKYWSGNSEFHFYSINVGGFFKLRSGEESIEVSNPSLLDPDQDATYFGAFKVRD
IDGGGGGCG

SEQ ID NO: 104

RANKL peptides AA`` loop
CGGNATDIPSGSHKVSLSWYHDRGWA

SEQ ID NO: 105

RANKL peptides DE loop
CGGSIKIPSSH

SEQ ID NO: 106

RANKL peptide b-strand D
CGGGYLQLMVYVTKT

SEQ ID NO: 107

RANKL peptide CD loop
CGGHETSGDLATE

SEQ ID NO: 108

RANKL peptide EF loop
CGGKYWSGNSEFHF

SEQ ID NO: 109

RANKL peptide B strand-B`B loop
CGGTFSNGKLI

SEQ ID NO: 110

RANKL peptide GH loop
CGGDPDQDA

SEQ ID NO: 111

CGAGCTCGCCCCCTGGCTTATCGAAATTAATACGACTCACTATAGGGAGACCGGAATTTCGA
GCTCGCCCCGGGGATCCTCTAGAATTTTCTGCGCACCCATCCCGGGTGGCGCCCAAAGTGA
GGAAATCACATGGCAAATAAGCCAATGCAACCGATCACATCTACAGCAAATAAAATTGT
GTGGTCGGATCCAACTCGTTTATCAACTACATTTTCAGCAAGTCTGTTACGCCAACGTGT
TAAAGTTGGTATAGCCGAACCTGAATAATGTTTCAGGTCAATATGTATCTGTTTATAAGCG
TCCTGCACCTAAACCGGAAGGTTGTGCAGATGCCTGTGTCAATTATGCCGAATGAAAACCA
ATCCATTTCGCACAGTGATTTTCAGGGTCAGCCGAAAACCTGGCTACCTTAAAAGCAGAATG
GGAAACTCACAAACGTAACGTTGACACACTCTTCGCGAGCGGCAACGCCGGTTTGGGTTT
CCTTGACCCCTACTGCGGCTATCGTATCGTCTGATACTACTGCTTAAGCTTGTATTCTATA
GTGTACCTAAATCGTATGTGTATGATACATAAGGTTATGTATTAATTGTAGCCGCGTTC
TAACGACAATATGTACAAGCCTAATTGTGTAGCATCTGGCTTACTGAAGCAGACCCCTATC
ATCTCTCTCGTAAACTGCCGTCAGAGTCGGTTTGGTTGGACGAACCTTCTGAGTTTCTGG
TAACGCCGTTCCGCACCCCGGAAATGGTCACCGAACCAATCAGCAGGGTCATCGCTAGCC
AGATCCTCTACGCCGGACGCATCGTGGCCGGCATCACCGCGCCACAGGTGCGGTTGCTG
GCGCCTATATCGCCGACATCACCGATGGGGAAGATCGGGCTCGCCACTTCGGGCTCATGA
GCGCTTGTTTCGGCGTGGGTATGGTGGCAGGCCCCGTGGCCGGGGGACTGTTGGGCGCCA
TCTCCTTGCAATGCACCATTCCTTGCGGCGGCGGTGCTCAACGGCCTCAACCTACTACTGG
CTGCTTCCCTAATGACAGGAGTCGCATAAGGGAGAGCGTCGATATGGTGCACCTCTCAGTAC
AATCTGCTCTGATGCCGCATAGTTAAGCCAACCTCCGCTATCGCTACGTGACTGGGTCATG
GCTGCGCCCCGACACCCGCCAACACCCGCTGACGCGCCCTGACGGGCTTGTCTGCTCCCG
GCATCCGCTTACAGACAAGCTGTGACCGTCTCCGGGAGCTGCATGTGTGACAGGTTTTCA
CCGTCATCACCGAAACGCGCGAGGCAGCTTGAAGACGAAAGGGCCTCGTGATACGCCTAT
TTTTATAGGTTAATGTCATGATAATAATGGTPTTCTTAGACGTCAGGTGGCACTTTTCGGG
GAAATGTGCGCGGAACCCCTATTTGTTTATTTTTCTAAATACATTCAAATATGTATCCGC
TCATGAGACAATAACCCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGATGATGAGTA
TTCAACATTTCCGTGTGCGCCCTTATTCCTTTTTCGCGGCATTTGCTTCTGTTTGTG
CTCACCCAGAAACGCTGGTGAAAGTAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGG
GTTACATCGAACTGGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAAC
GTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATTG
ACGCCGGGCAAGAGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTTGGTTGAGT
ACTCACAGTCACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTG
CTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCTGACAACGATCGGAGGAC
CGAAGGAGCTAACCCTTTTTTGCAACAACATGGGGGATCATGTAACCTGCCTTGATCGTT
GGGAACCGGAGCTGAATGAAGCCATACCAACGACGAGCGTGACACCACGATGCCGTAG
CAATGGCAACAACGTTGCGCAAACTATTAACCTGGCGAACTACTTACTCTAGCTTCCCGGC
AACAATTAATAGACTGGATGGAGCGGATAAAGTTGCAGGACCCTTCTGCGCTCGGCCC
TTCCGGCTGGCTGGTTTATTTGCTGATAAATCTGGAGCCGGTGAGCGTGGGTCTCGCGGTA
TCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGTAGTTATCTACACGACGG
GGAGTCAGGCAACTATGGATGAACGAAATAGACAGATCGCTGAGATAGGTGCCTCACTGA
TTAAGCATTGGTAACTGTGACCAAGTTTACTCATATATACTTTAGATTGATTTAAAC
TTCATTTTTAATTTAAAGGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAAA
TCCCTTAACGTGAGTTTTCTGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGAT
CTTCTTGAGATCCTTTTTTTCTGCGCGTAATCTGCTGCTTGCAACAAAAAACCCCGC
TACCAGCGGTGGTTTGTGTTGCCGGATCAAGAGCTACCAACTCTTTTCCGAAGGTAAC TG
GCTTCAGCAGAGCGCAGATACCAAACTGTCTTCTAGTGTAGCCGTAGTTAGGCCACC
ACTTCAAGAACTCTGTAGCACCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGG

CTGCTGCCAGTGGCGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATAGTTACCGG
ATAAGGCGCAGCGGTCGGGCTGAACGGGGGGTTCGTGCACACAGCCAGCTTGGAGCGAA
CGACCTACACCGAACTGAGATACCTACAGCGCGAGCATTGAGAAAGCGCCACGCTTCCCG
AAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGA
GGGAGCTTCCAGGGGGAAACGCCTGGTATCTTTATAGTCCTGTCGGGTTTCGCCACCTCT
GACTTGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGGCGGAGCCTATGGAAAACGCCA
GCAACGCGGCCTTTTTACGGTTCCTGGCCTTTTGTGCTGGCCTTTTGTCTACATGTTCTTTC
CTGCGTTATCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTGAGCTGATACCG
CTCGCCGCGAGCCGAACGACCGAGCGCAGCGAGTCAGTGAGCGAGGAAGCGGAAGAGCGCC
CAATACGCAAACCGCCTCTCCCCGCGCGTTGGCCGATTCAATTAATGCAGCTGTGGTGTCA
TGGTCGGTGATCGCCAGGGTGCCGACGCGCATCTCGACTGCATGGTGACCAATGCTTCT
GGCGTCAGGCAGCCATCGGAAGCTGTGGTATGGCCGTGCAGGTCGTAAATCACTGCATAA
TTCGTGTGCTCAAGGCGCACCTCCCGTTCGGATAATGTTTTTTCGCCCGACATCATAAC
GGTTCGGCAAATATCTGAAATGAGCTGTTGACAATTAATCATCGAACTAGTTAACTAG
TACGCAAGTTCACGTAAAAAGGTATCGCGGAATT

SEQ ID NO: 112

MANKPMQPITSTANKIVWSDPTLSTTFASLLRQRVKVGIAELNNVSGQYVSVYKRPAPKPEGCADACVIMPNNQS
IRTVISGSAENLATLKAETHKRNVDTLFASGNAGLGFLDPTAAIVSSDTTA

SEQ ID NO: 113

MANKTMQPITSTANKIVWSDPTLSTTFASLLRQRVKVGIAELNNVSGQYVSVYKRPAPKPEGCADACVIMPNNQS
IRTVISGSAENLATLKAETHKRNVDTLFASGNAGLGFLDPTAAIVSSDTTA

SEQ ID NO: 114

CGAGCTCGCCCCCTGGCTTATCGAAATTAATACGACTCACTATAGGGAGACCGGAATTCGA
GCTCGCCCGGGGATCCTCTAGATTAACCCAACGCGTAGGAGTCAGGCCATGGCAAATAAG
ACAATGCAACCGATCACATCTACAGCAAATAAAATTGTGTGGTCGGATCCAACTCGTTTA
TCAACTACATTTTCAGCAAGTCTGTTACGCCAACGTGTTAAAGTTGGTATAGCCGAACGTG
AATAATGTTTTCAGGTCAATATGTATCTGTTTTATAAGCGTCTTGCACCTAAACCGGAAGGT
TGTGCAGATGCCTGTGTCAATTATGCCGAATGAAAACCAATCCATTTCGCACAGTGATTTCA
GGGTCAGCCGAAAACCTTGGCTACCTTAAAAGCAGAATGGGAAACTCACAAACGTAACGTT
GACACACTCTTCGCGAGCGGCAACGCCGGTTTGGGTTTCCTTGACCCTACTGCGGCTATC
GTATCGTCTGATACTACTGCTTAAGCTTGTATTCTATAGTGTACCTAAATCGTATGTGT
ATGATACATAAGGTTATGTATTAATTGTAGCCGCGTTCTAACGACAATATGTACAAGCCT
AATTGTGTAGCATCTGGCTTACTGAAGCAGACCCATCATCTCTCTCGTAAACTGCCGTC
AGAGTCGGTTTTGGTTGGACGAACCTTCTGAGTTTTCTGGTAACGCCGTTCCGCACCCCGGA
AATGGTCACCGAACCAATCAGCAGGGTCATCGCTAGCCAGATCCTCTACGCCGGACGCAT
CGTGGCCGGCATACCGGCGCCACAGGTGCGGTTGCTGGCGCCTATATCGCCGACATCAC
CGATGGGGAAGATCGGGCTCGCCACTTCGGGCTCATGAGCGCTTGTTTCGGCGTGCGGTAT
GGTGGCAGGCCCCGTGGCCGGGGACTGTTGGGCGCCATCTCCTTGTCATGCACCATTCCT
TGCGGCGGCGGTGCTCAACGGCCTCAACCTACTACTGGGCTGCTTCCTAATGCAGGAGTC
GCATAAGGGAGAGCGTCGATATGGTGCACCTCTCAGTACAATCTGCTCTGATGCCGATAG
TTAAGCCAACTCCGCTATCGCTACGTGACTGGGTCATGGCTGCGCCCCGACACCCGCCAA
CACCCGCTGACGCGCCCTGACGGGCTTGCTGCTCCCGCATCCGCTTACAGACAAGCTG
TGACCGTCTCCGGGAGCTGCATGTGTGACAGGTTTTTACCCTCATCACCGAAACGCGCGA
GGCAGCTTGAAGACGAAAGGGCCTCGTGATACGCCATTTTTTATAGGTTAATGTCATGAT
AATAATGGTTTTCTTAGACGTACAGGTGGCACTTTTCGGGAAATGTGCGCGGAACCCCTAT
TTGTTTTATTTTCTAAATACATTCAAATATGTATCCGCTCATGAGACAATAACCCGTGATA

AATGCTTCAATAATATTGAAAAAGGAAGAGTATGAGTATTCAACATTTCCGTGTCGCCCT
TATTCCTTTTTTTCGGCATTTTTCCTTCCGTGTTTTTTCGCTCACCAGAAACGCTGGTGAA
AGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAA
CAGCGGTAAGATCCTTGAGAGTTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTT
TAAAGTTCTGCTATGTGGCGCGGTATTATCCCCTATTGACGCCGGGCAAGAGCAACTCGG
TCGCCGCATACACTATTCTCAGAAATGACTTGGTTGAGTACTCACCAGTCACAGAAAAGCA
TCTTACGGATGGCATGACAGTAAGAGAATTATGCACTGCTGCCATAACCATGAGTGATAA
CACTGCGGCCAACTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTT
GCACAACATGGGGGATCATGTAACCTGCCCTTGATCGTTGGGAACCGGAGCTGAATGAAGC
CATACCAAACGACGAGCGTGACACCACGATGCCCTGTAGCAATGGCAACAACGTTGCGCAA
ACTATTAACGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGA
GGCGGATAAAGTTGCAGGACCCTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTTATTGC
TGATAAATCTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTCAGCACTGGGGCCAGA
TGGAAGCCCTCCCGTATCGTAGTTATCTACACGACGGGAGTCAGGCAACTATGGATGA
ACGAAATAGACAGATCGCTGAGATAGGTGCCCTCACTGATTAAGCATTTGGTAACTGTCAGA
CCAAGTTTACTCATATATACTTTAGATTGATTTAAAACCTCATTTTTTAATTTAAAAGGAT
CTAGGTGAAGATCCTTTTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGTTTTTCGTT
CCACTGAGCGTCAGACCCCGTAGAAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCT
GCGCGTAATCTGCTGCTTGCAAAACAAAAAACACCGCTACCAGCGGTGGTTTTGTTGCC
GGATCAAGAGCTACCAACTCTTTTTCCGAAGGTAACCTGGCTTCAGCAGAGCGCAGATACC
AAATACGTCTCTTAGTGTAGCCGTAGTTAGGCCACCCTCAAGAATCTGTAGCACC
GCCATACATACCCTGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTC
GTGCTTTACCGGTTGGACTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTCCGGCTG
AACGGGGGGTTCTGTGCACACAGCCAGCTTGGAGCGAACGACCTACACCGAAGTGAAGATA
CCTACAGCGCGAGCATTTGAGAAAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTA
TCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGAAACGC
CTGGTATCTTTATAGTCTGTGCGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTG
ATGCTCGTCAAGGGGGCGGAGCCTATGGAACCAACCGCGGCTTTTTACGGTT
CCTGGCTTTTTGCTGGCTTTTTGCTCAGATGTTCTTTCTGCGTTATCCCTGATTCTGT
GGATAACCGTATTACCGCTTTGAGTGAGCTGATACCGCTCGCCGAGCCGAACGACCGA
GCGCAGCGAGTCAGTGAGCGAGGAAGCGGAAGAGCGCCCAATACGCAAAACCGCTCTCCC
CGCGCGTTGGCCGATTCAATTAATGCAGCTGTGGTGTGATGGTGGTGATCGCCAGGGTGC
CGACGCGCATCTCGACTGCATGGTGCACCAATGCTTCTGGCGTCAGGCAGCCATCGGAAG
CTGTGGTATGGCCGTGCAGGTGTAATCACTGCATAATTCGTGTGCTCAAGGCGCACT
CCCGTTCTGGATAATGTTTTTTGCGCCGACATCATAACGGTTCTGGCAAAATATTCTGAAA
TGAGCTGTTGACAATTAATCATCGAACTAGTTAACTAGTACGCAAGTTACGTAATAAAGG
GTATCGCGGAATT

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.